

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
18 July 2002 (18.07.2002)

PCT

(10) International Publication Number  
**WO 02/055548 A2**

- (51) International Patent Classification<sup>7</sup>: **C07K 14/005**
- (21) International Application Number: **PCT/EP02/00219**
- (22) International Filing Date: 11 January 2002 (11.01.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/260,699 11 January 2001 (11.01.2001) US  
60/315,768 30 August 2001 (30.08.2001) US
- (71) Applicant (for all designated States except US): **INNO-GENETICS N.V. [BE/BE]**; Industriepark Zwijnaarde 7, Box 4, B-9052 Ghent (BE).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **MAERTENS, Geert [BE/BE]**; Zilverparrenstraat 64, B-8310 Brugge (BE). **BOSMAN, Fons [BE/BE]**; Hulst 165, B-1745 Opwijk (BE). **BUYSE, Marie-Ange [BE/BE]**; E. Ronsestraat 23, B-9820 Merelbeke (BE).
- (74) Common Representative: **INNOGENETICS N.V.**; Industriepark Zwijnaarde 7, Box 4, B-9052 Ghent (BE).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
— without international search report and to be republished upon receipt of that report  
— with (an) indication(s) in relation to deposited biological material furnished under Rule 13bis separately from the description
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/055548 A2

(54) Title: **PURIFIED HEPATITIS C VIRUS ENVELOPE PROTEINS FOR DIAGNOSTIC AND THERAPEUTIC USE**

(57) Abstract: The present invention relates to a method for purifying recombinant HCV single or specific oligomeric envelope proteins selected from the group consisting of E1 and/or E2 and/or E1/E2, characterized in that upon lysing the transformed host cells to isolate the recombinantly expressed protein a disulphide bond cleavage or reduction step is carried out with a disulphide bond cleavage agent. The present invention also relates to a composition isolated by such a method. The present invention also relates to the diagnostic and therapeutic application of these compositions. Furthermore, the invention relates to the use of HCV E1 protein and peptides for prognosing and monitoring the clinical effectiveness and/or clinical outcome if HCV treatment.

PURIFIED HEPATITIS C VIRUS ENVELOPE PROTEINS FOR DIAGNOSTIC AND THERAPEUTIC USEField of the invention

5

The present invention relates to the general fields of recombinant protein expression, purification of recombinant proteins, synthetic peptides, diagnosis of HCV infection, prophylactic treatment against HCV infection and to the prognosis/monitoring of the clinical efficiency of treatment of an individual with chronic hepatitis, or the prognosis/monitoring of natural disease.

10

More particularly, the present invention relates to purification methods for hepatitis C virus envelope proteins, the use in diagnosis, prophylaxis or therapy of HCV envelope proteins purified according to the methods described in the present invention, the use of single or specific oligomeric E1 and/or E2 and/or E1/E2 envelope proteins in assays for monitoring disease, and/or diagnosis of disease, and/or treatment of disease. The invention also relates to epitopes of the E1 and/or E2 envelope proteins and monoclonal antibodies thereto, as well their use in diagnosis, prophylaxis or treatment.

15

Background of the invention

The E2 protein purified from cell lysates according to the methods described in the present invention reacts with approximately 95% of patient sera. This reactivity is similar to the reactivity obtained with E2 secreted from CHO cells (Spaete et al., 1992). However, the intracellularly expressed form of E2 may more closely resemble the native viral envelope protein because it contains high mannose carbohydrate motifs, whereas the E2 protein secreted from CHO cells is further modified with galactose and sialic acid sugar moieties. When the aminoterminal half of E2 is expressed in the baculovirus system, only about 13 to 21% of sera from several patient groups can be detected (Inoue et al., 1992). After expression of E2 from E. coli, the reactivity of HCV sera was even lower and ranged from 14 (Yokosuka et al., 1992) to 17% (Mita et al., 1992). About 75% of HCV sera (and 95% of chronic patients) are anti-E1 positive using the purified, vaccinia-expressed recombinant E1 protein of the present invention, in sharp contrast with the results of Kohara et al. (1992) and Hsu et al. (1993). Kohara et al. used a vaccinia-virus expressed E1 protein and detected anti-E1 antibodies in 7 to 23% of patients, while Hsu et al. only detected 14/50 (28%) sera using baculovirus-expressed E1.

20  
25  
30

These results show that not only a good expression system but also a good purification protocol are required to reach a high reactivity of the envelope proteins with human patient sera. This can be obtained using the proper expression system and/or purification protocols of the present invention which guarantee the conservation of the natural folding of the protein and the purification protocols of the present invention which

guarantee the elimination of contaminating proteins and which preserve the conformation, and thus the reactivity of the HCV envelope proteins. The amounts of purified HCV envelope protein needed for diagnostic screening assays are in the range of grams per year. For vaccine purposes, even higher amounts of envelope protein would be needed. Therefore, the vaccinia virus system may be used for selecting the best expression constructs and for limited upscaling, and large-scale expression and purification of single or specific oligomeric envelope proteins containing high-mannose carbohydrates may be achieved when expressed from several yeast strains. In the case of hepatitis B for example, manufacturing of HBsAg from mammalian cells was much more costly compared with yeast-derived hepatitis B vaccines.

10 Aims of the invention

It is an aim of the present invention to provide a new purification method for recombinantly expressed E1 and/or E2 and/or E1/E2 proteins such that said recombinant proteins are directly usable for diagnostic and vaccine purposes as single or specific oligomeric recombinant proteins free from contaminants instead of aggregates.

It is another aim of the present invention to provide compositions comprising purified (single or specific oligomeric) recombinant E1 and/or E2 and/or E1/E2 glycoproteins comprising conformational epitopes from the E1 and/or E2 domains of HCV.

It is yet another aim of the present invention to provide novel recombinant vector constructs for recombinantly expressing E1 and/or E2 and/or E1/E2 proteins, as well as host cells transformed with said vector constructs.

It is also an aim of the present invention to provide a method for producing and purifying recombinant HCV E1 and/or E2 and/or E1/E2 proteins.

It is also an aim of the present invention to provide diagnostic and immunogenic uses of the recombinant HCV E1 and/or E2 and/or E1/E2 proteins of the present invention, as well as to provide kits for diagnostic use, vaccines or therapeutics comprising any of the recombinant HCV E1 and/or E2 and/or E1/E2 proteins of the present invention.

It is further an aim of the present invention to provide for a new use of E1, E2, and/or E1/E2 proteins, or suitable parts thereof, for monitoring/prognosing the response to treatment of patients (e.g. with interferon) suffering from HCV infection.

It is also an aim of the present invention to provide for the use of the recombinant E1, E2, and/or E1/E2 proteins of the present invention in HCV screening and confirmatory antibody tests.

It is also an aim of the present invention to provide E1 and/or E2 peptides which can be used for diagnosis of HCV infection and for raising antibodies. Such peptides may also be used to isolate human

**PROOF OF SERVICE**

DATE	PLACE
SERVED	
SERVED ON (PRINT NAME)	MANNER OF SERVICE
SERVED BY (PRINT NAME)	TITLE

**DECLARATION OF SERVER**

I declare under penalty of perjury under the laws of the United States of America that the foregoing information contained in the Proof of Service is true and correct.

Executed on	DATE	SIGNATURE OF SERVER
		ADDRESS OF SERVER

Rule 45, Federal Rules of Civil Procedure, Parts C & D:

**(c) PROTECTION OF PERSONS SUBJECT TO SUBPOENAS.**

(1) A party or an attorney responsible for the issuance and service of a subpoena shall take reasonable steps to avoid imposing undue burden or expense on a person subject to that subpoena. The court on behalf of which the subpoena was issued shall enforce this duty and impose upon the party or attorney in breach of this duty an appropriate sanction, which may include, but is not limited to, lost earnings and reasonable attorney's fee.

(2)(A) A person commanded to produce and permit inspection and copying of designated books, papers, documents or tangible things, or inspection of premises need not appear in person at the place of production or inspection unless commanded to appear for deposition, hearing or trial.

(b) Subject to paragraph (d)(2) of this rule, a person commanded to produce and permit inspection and copying may, within 14 days after service of the subpoena or before the time specified for compliance if such time is less than 14 days after service, serve upon the party or attorney designated in the subpoena written objection to inspection or copying of any or all of the designated materials or of the premises. If objection is made, the party serving the subpoena shall not be entitled to inspect and copy the materials or inspect the premises except pursuant to an order of the court by which the subpoena was issued. If objection has been made, the party serving the subpoena may, upon notice to the person commanded to produce, move at any time for an order to compel the production. Such an order to compel production shall protect any person who is not a party or an officer of a party from significant expense resulting from the inspection and copying commanded.

(3)(A) On timely motion, the court by which a subpoena was issued shall quash or modify the subpoena if it

- (i) fails to allow reasonable time for compliance;
- (ii) requires a person who is not a party or an officer of a party to travel to a place more than 100 miles from the place where the person resides, is employed

or regularly transacts business in person, except that, subject to the provisions of clause (c)(3)(B)(iii) of this rule, such a person may in order to attend trial be commanded to travel from any such place within the state in which the trial is held, or

- (iii) requires disclosure of privileged or other protected matter and no exception or waiver applies, or
- (iv) subjects a person to undue burden.

**(B) If a subpoena**

(i) requires disclosure of a trade secret or other confidential research, development, or commercial information, or

(ii) requires disclosure of an unretained expert's opinion or information not describing specific events or occurrences in dispute and resulting from the expert's study made not at the request of any party, or

(iii) requires a person who is not a party or an officer of a party to incur substantial expense to travel more than 100 miles to attend trial, the court may, to protect a person subject to or affected by the subpoena, quash or modify the subpoena or, if the party in whose behalf the subpoena is issued shows a substantial need for the testimony or material that cannot be otherwise met without the undue hardship and assures that the person to whom the subpoena is addressed will be reasonably compensated, the court may order appearance or production only upon specified conditions.

**(d) DUTIES IN RESPONDING TO SUBPOENA.**

(1) A person responding to a subpoena to produce documents shall produce them as they are kept in the usual course of business or shall organize and label them to correspond with the categories in the demand.

(2) When information subject to a subpoena is withheld on a claim that is privileged or subject to protection as trial preparation materials the claim shall be made expressly and shall be supported by a description of the nature of the documents, communications, or things not produced that is sufficient to enable the demanding party to contest the claim.



monoclonal antibodies.

It is also an aim of the present invention to provide monoclonal antibodies, more particularly human monoclonal antibodies or mouse monoclonal antibodies which are humanized, which react specifically with E1 and/or E2 epitopes, either comprised in peptides or conformational epitopes comprised in recombinant proteins.

5 It is also an aim of the present invention to provide possible uses of anti-E1 or anti-E2 monoclonal antibodies for HCV antigen detection or for therapy of chronic HCV infection.

It is also an aim of the present invention to provide kits for monitoring/prognosing the response to treatment (e.g. with interferon) of patients suffering from HCV infection or monitoring/prognosing the outcome of the disease.

10 All the aims of the present invention are considered to have been met by the embodiments as set out below.

#### Definitions

15 The following definitions serve to illustrate the different terms and expressions used in the present invention.

The term 'hepatitis C virus single envelope protein' refers to a polypeptide or an analogue thereof (e.g. mimotopes) comprising an amino acid sequence (and/or amino acid analogues) defining at least one HCV epitope of either the E1 or the E2 region. These single envelope proteins in the broad sense of the word may be both monomeric or homo-oligomeric forms of recombinantly expressed envelope proteins. Typically, the sequences defining the epitope correspond to the amino acid sequence of either the E1 or the E2 region of HCV (either identically or via substitution of analogues of the native amino acid residue that do not destroy the epitope). In general, the epitope-defining sequence will be 3 or more amino acids in length, more typically, 5 or more amino acids in length, more typically 8 or more amino acids in length, and even more typically 10 or more amino acids in length. With respect to conformational epitopes, the length of the epitope-defining sequence can be subject to wide variations, since it is believed that these epitopes are formed by the three-dimensional shape of the antigen (e.g. folding). Thus, the amino acids defining the epitope can be relatively few in number, but widely dispersed along the length of the molecule being brought into the correct epitope conformation via folding. The portions of the antigen between the residues defining the epitope may not be critical to the conformational structure of the epitope. For example, deletion or substitution of these intervening sequences may not affect the conformational epitope provided sequences critical to epitope conformation are maintained (e.g. cysteines involved in disulfide bonding, glycosylation sites, etc.). A conformational epitope may also be formed by 2 or more essential regions of subunits of a homooligomer or heterooligomer.

30 The HCV antigens of the present invention comprise conformational epitopes from the E1 and/or E2

(envelope) domains of HCV. The E1 domain, which is believed to correspond to the viral envelope protein, is currently estimated to span amino acids 192-383 of the HCV polyprotein (Hijikata et al., 1991). Upon expression in a mammalian system (glycosylated), it is believed to have an approximate molecular weight of 35 kDa as determined via SDS-PAGE. The E2 protein, previously called NS1, is believed to span amino acids 384-809 or 384-746 (Grakoui et al., 1993) of the HCV polyprotein and to also be an envelope protein. Upon expression in a vaccinia system (glycosylated), it is believed to have an apparent gel molecular weight of about 72 kDa. It is understood that these protein endpoints are approximations (e.g. the carboxy terminal end of E2 could lie somewhere in the 730-820 amino acid region, e.g. ending at amino acid 730, 735, 740, 742, 744, 745, preferably 746, 747, 748, 750, 760, 770, 780, 790, 800, 809, 810, 820). The E2 protein may also be expressed together with the E1, P7 (aa 747-809), NS2 (aa 810-1026), NS4A (aa 1658-1711) or NS4B (aa 1712-1972). Expression together with these other HCV proteins may be important for obtaining the correct protein folding.

It is also understood that the isolates used in the examples section of the present invention were not intended to limit the scope of the invention and that any HCV isolate from type 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or any other new genotype of HCV is a suitable source of E1 and/or E2 sequence for the practice of the present invention.

The E1 and E2 antigens used in the present invention may be full-length viral proteins, substantially full-length versions thereof, or functional fragments thereof (e.g. fragments which are not missing sequence essential to the formation or retention of an epitope). Furthermore, the HCV antigens of the present invention can also include other sequences that do not block or prevent the formation of the conformational epitope of interest. The presence or absence of a conformational epitope can be readily determined through screening the antigen of interest with an antibody (polyclonal serum or monoclonal to the conformational epitope) and comparing its reactivity to that of a denatured version of the antigen which retains only linear epitopes (if any). In such screening using polyclonal antibodies, it may be advantageous to adsorb the polyclonal serum first with the denatured antigen and see if it retains antibodies to the antigen of interest.

The HCV antigens of the present invention can be made by any recombinant method that provides the epitope of interest. For example, recombinant intracellular expression in mammalian or insect cells is a preferred method to provide glycosylated E1 and/or E2 antigens in 'native' conformation as is the case for the natural HCV antigens. Yeast cells and mutant yeast strains (e.g. mnn 9 mutant (Kniskern et al., 1994) or glycosylation mutants derived by means of vanadate resistance selection (Ballou et al., 1991)) may be ideally suited for production of secreted high-mannose-type sugars; whereas proteins secreted from mammalian cells may contain modifications including galactose or sialic acids which may be undesirable for certain diagnostic or vaccine applications. However, it may also be possible and sufficient for certain applications, as it is known for proteins, to express the antigen in other recombinant hosts (such as *E. coli*) and renature the protein after recovery.

The term 'fusion polypeptide' intends a polypeptide in which the HCV antigen(s) are part of a single continuous chain of amino acids, which chain does not occur in nature. The HCV antigens may be connected directly to each other by peptide bonds or be separated by intervening amino acid sequences. The fusion polypeptides may also contain amino acid sequences exogenous to HCV.

5       The term 'solid phase' intends a solid body to which the individual HCV antigens or the fusion polypeptide comprised of HCV antigens are bound covalently or by noncovalent means such as hydrophobic adsorption.

10       The term 'biological sample' intends a fluid or tissue of a mammalian individual (e.g. an anthropoid, a human) that commonly contains antibodies produced by the individual, more particularly antibodies against HCV. The fluid or tissue may also contain HCV antigen. Such components are known in the art and include, without limitation, blood, plasma, serum, urine, spinal fluid, lymph fluid, secretions of the respiratory, intestinal or genitourinary tracts, tears, saliva, milk, white blood cells and myelomas. Body components include biological liquids. The term 'biological liquid' refers to a fluid obtained from an organism. Some biological fluids are used as a source of other products, such as clotting factors (e.g. Factor VIII;C), serum albumin, growth hormone and the like. In such cases, it is important that the source of biological fluid be free of contamination by virus such as HCV.

15       The term 'immunologically reactive' means that the antigen in question will react specifically with anti-HCV antibodies present in a body component from an HCV infected individual.

20       The term 'immune complex' intends the combination formed when an antibody binds to an epitope on an antigen.

25       'E1' as used herein refers to a protein or polypeptide expressed within the first 400 amino acids of an HCV polyprotein, sometimes referred to as the E, ENV or S protein. In its natural form it is a 35 kDa glycoprotein which is found in strong association with membranes. In most natural HCV strains, the E1 protein is encoded in the viral polyprotein following the C (core) protein. The E1 protein extends from approximately amino acid (aa) 192 to about aa 383 of the full-length polyprotein.

      The term 'E1' as used herein also includes analogs and truncated forms that are immunologically cross-reactive with natural E1, and includes E1 proteins of genotypes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or any other newly identified HCV type or subtype.

30       'E2' as used herein refers to a protein or polypeptide expressed within the first 900 amino acids of an HCV polyprotein, sometimes referred to as the NS1 protein. In its natural form it is a 72 kDa glycoprotein that is found in strong association with membranes. In most natural HCV strains, the E2 protein is encoded in the viral polyprotein following the E1 protein. The E2 protein extends from approximately amino acid position 384 to amino acid position 746, another form of E2 extends to amino acid position 809. The term 'E2' as used herein also includes analogs and truncated forms that are immunologically cross-reactive with natural E2. For example,

insertions of multiple codons between codon 383 and 384, as well as deletions of amino acids 384-387 have been reported by Kato et al. (1992).

'E1/E2' as used herein refers to an oligomeric form of envelope proteins containing at least one E1 component and at least one E2 component.

5       The term 'specific oligomeric' E1 and/or E2 and/or E1/E2 envelope proteins refers to all possible oligomeric forms of recombinantly expressed E1 and/or E2 envelope proteins which are not aggregates. E1 and/or E2 specific oligomeric envelope proteins are also referred to as homo-oligomeric E1 or E2 envelope proteins (see below).

10       The term 'single or specific oligomeric' E1 and/or E2 and/or E1/E2 envelope proteins refers to single monomeric E1 or E2 proteins (single in the strict sense of the word) as well as specific oligomeric E1 and/or E2 and/or E1/E2 recombinantly expressed proteins. These single or specific oligomeric envelope proteins according to the present invention can be further defined by the following formula  $(E1)_x(E2)_y$ , wherein x can be a number between 0 and 100, and y can be a number between 0 and 100, provided that x and y are not both 0. With x=1 and y=0 said envelope proteins include monomeric E1.

15       The term 'homo-oligomer' as used herein refers to a complex of E1 and/or E2 containing more than one E1 or E2 monomer, e.g. E1/E1 dimers, E1/E1/E1 trimers or E1/E1/E1/E1 tetramers and E2/E2 dimers, E2/E2/E2 trimers or E2/E2/E2/E2 tetramers, E1 pentamers and hexamers, E2 pentamers and hexamers or any higher-order homo-oligomers of E1 or E2 are all 'homo-oligomers' within the scope of this definition. The oligomers may contain one, two, or several different monomers of E1 or E2 obtained from different types or  
20       subtypes of hepatitis C virus including for example those described in an international application published under WO 94/25601 and European application No. 94870166.9 both by the present applicants. Such mixed oligomers are still homo-oligomers within the scope of this invention, and may allow more universal diagnosis, prophylaxis or treatment of HCV.

25       The term 'purified' as applied to proteins herein refers to a composition wherein the desired protein comprises at least 35% of the total protein component in the composition. The desired protein preferably comprises at least 40%, more preferably at least about 50%, more preferably at least about 60%, still more preferably at least about 70%, even more preferably at least about 80%, even more preferably at least about 90%, and most preferably at least about 95% of the total protein component. The composition may contain other compounds such as carbohydrates, salts, lipids, solvents, and the like, without affecting the determination of  
3.0       the percentage purity as used herein. An 'isolated' HCV protein intends an HCV protein composition that is at least 35% pure.

      The term 'essentially purified proteins' refers to proteins purified such that they can be used for in vitro diagnostic methods and as a therapeutic compound. These proteins are substantially free from cellular proteins, vector-derived proteins or other HCV viral components. Usually these proteins are purified to homogeneity (at

least 80% pure, preferably, 90%, more preferably 95%, more preferably 97%, more preferably 98%, more preferably 99%, even more preferably 99.5%, and most preferably the contaminating proteins should be undetectable by conventional methods like SDS-PAGE and silver staining.

5 The term 'recombinantly expressed' used within the context of the present invention refers to the fact that the proteins of the present invention are produced by recombinant expression methods be it in prokaryotes, or lower or higher eukaryotes as discussed in detail below.

10 The term 'lower eukaryote' refers to host cells such as yeast, fungi and the like. Lower eukaryotes are generally (but not necessarily) unicellular. Preferred lower eukaryotes are yeasts, particularly species within Saccharomyces, Schizosaccharomyces, Kluveromyces, Pichia (e.g. Pichia pastoris), Hansenula (e.g. Hansenula polymorpha), Yarrowia, Schwaniomyces, Schizosaccharomyces, Zygosaccharomyces and the like. Saccharomyces cerevisiae, S. carlsbergensis and K. lactis are the most commonly used yeast hosts, and are convenient fungal hosts.

The term 'prokaryotes' refers to hosts such as E.coli, Lactobacillus, Lactococcus, Salmonella, Streptococcus, Bacillus subtilis or Streptomyces. Also these hosts are contemplated within the present invention.

15 The term 'higher eukaryote' refers to host cells derived from higher animals, such as mammals, reptiles, insects, and the like. Presently preferred higher eukaryote host cells are derived from Chinese hamster (e.g. CHO), monkey (e.g. COS and Vero cells), baby hamster kidney (BHK), pig kidney (PK15), rabbit kidney 13 cells (RK13), the human osteosarcoma cell line 143 B, the human cell line HeLa and human hepatoma cell lines like Hep G2, and insect cell lines (e.g. Spodoptera frugiperda). The host cells may be provided in suspension or  
20 flask cultures, tissue cultures, organ cultures and the like. Alternatively the host cells may also be transgenic animals.

The term 'polypeptide' refers to a polymer of amino acids and does not refer to a specific length of the product; thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide. This term also does not refer to or exclude post-expression modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations and the like. Included within the definition are, for example, polypeptides  
25 containing one or more analogues of an amino acid (including, for example, unnatural amino acids, PNA, etc.), polypeptides with substituted linkages, as well as other modifications known in the art, both naturally occurring and non-naturally occurring.

The term 'recombinant polynucleotide or nucleic acid' intends a polynucleotide or nucleic acid of  
30 genomic, cDNA, semisynthetic, or synthetic-origin which, by virtue of its origin or manipulation : (1) is not associated with all or a portion of a polynucleotide with which it is associated in nature, (2) is linked to a polynucleotide other than that to which it is linked in nature, or (3) does not occur in nature.

The term 'recombinant host cells', 'host cells', 'cells', 'cell lines', 'cell cultures', and other such terms denoting microorganisms or higher eukaryotic cell lines cultured as unicellular entities refer to cells which can be

or have been, used as recipients for a recombinant vector or other transfer polynucleotide, and include the progeny of the original cell which has been transfected. It is understood that the progeny of a single parental cell may not necessarily be completely identical in morphology or in genomic or total DNA complement as the original parent, due to natural, accidental, or deliberate mutation.

- 5           The term 'replicon' is any genetic element, e.g., a plasmid, a chromosome, a virus, a cosmid, etc., that behaves as an autonomous unit of polynucleotide replication within a cell; i.e., capable of replication under its own control.

          The term 'vector' is a replicon further comprising sequences providing replication and/or expression of a desired open reading frame.

- 10           The term 'control sequence' refers to polynucleotide sequences which are necessary to effect the expression of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism; in prokaryotes, such control sequences generally include promoter, ribosomal binding site, and terminators; in eukaryotes, generally, such control sequences include promoters, terminators and, in some instances, enhancers. The term 'control sequences' is intended to include, at  
15           a minimum, all components whose presence is necessary for expression, and may also include additional components whose presence is advantageous, for example, leader sequences which govern secretion.

          The term 'promoter' is a nucleotide sequence which is comprised of consensus sequences which allow the binding of RNA polymerase to the DNA template in a manner such that mRNA production initiates at the normal transcription initiation site for the adjacent structural gene.

- 20           The expression 'operably linked' refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. A control sequence 'operably linked' to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

- An 'open reading frame' (ORF) is a region of a polynucleotide sequence which encodes a polypeptide  
25           and does not contain stop codons; this region may represent a portion of a coding sequence or a total coding sequence.

- A 'coding sequence' is a polynucleotide sequence which is transcribed into mRNA and/or translated into a polypeptide when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a translation start codon at the 5'-terminus and a translation stop codon at  
30           the 3'-terminus. A coding sequence can include but is not limited to mRNA, DNA (including cDNA), and recombinant polynucleotide sequences.

          As used herein, 'epitope' or 'antigenic determinant' means an amino acid sequence that is immunoreactive. Generally an epitope consists of at least 3 to 4 amino acids, and more usually, consists of at least 5 or 6 amino acids, sometimes the epitope consists of about 7 to 8, or even about 10 amino acids. As used

herein, an epitope of a designated polypeptide denotes epitopes with the same amino acid sequence as the epitope in the designated polypeptide, and immunologic equivalents thereof. Such equivalents also include strain, subtype (=genotype), or type(group)-specific variants, e.g. of the currently known sequences or strains belonging to genotypes 1a, 1b, 1c, 1d, 1e, 1f, 2a, 2b, 2c, 2d, 2e, 2f, 2g, 2h, 2i, 3a, 3b, 3c, 3d, 3e, 3f, 3g, 4a, 4b, 4c, 4d, 4e, 4f, 4g, 4h, 4i, 4j, 4k, 4l, 5a, 5b, 6a, 6b, 6c, 7a, 7b, 7c, 8a, 8b, 9a, 9b, 10a, or any other newly defined HCV (sub)type. It is to be understood that the amino acids constituting the epitope need not be part of a linear sequence, but may be interspersed by any number of amino acids, thus forming a conformational epitope.

The term 'immunogenic' refers to the ability of a substance to cause a humoral and/or cellular response, whether alone or when linked to a carrier, in the presence or absence of an adjuvant. 'Neutralization' refers to an immune response that blocks the infectivity, either partially or fully, of an infectious agent. A 'vaccine' is an immunogenic composition capable of eliciting protection against HCV, whether partial or complete. A vaccine may also be useful for treatment of an individual, in which case it is called a therapeutic vaccine.

The term 'therapeutic' refers to a composition capable of treating HCV infection.

The term 'effective amount' refers to an amount of epitope-bearing polypeptide sufficient to induce an immunogenic response in the individual to which it is administered, or to otherwise detectably immunoreact in its intended system (e.g., immunoassay). Preferably, the effective amount is sufficient to effect treatment, as defined above. The exact amount necessary will vary according to the application. For vaccine applications or for the generation of polyclonal antiserum / antibodies, for example, the effective amount may vary depending on the species, age, and general condition of the individual, the severity of the condition being treated, the particular polypeptide selected and its mode of administration, etc. It is also believed that effective amounts will be found within a relatively large, non-critical range. An appropriate effective amount can be readily determined using only routine experimentation. Preferred ranges of E1 and/or E2 and/or E1/E2 single or specific oligomeric envelope proteins for prophylaxis of HCV disease are 0.01 to 100 µg/dose, preferably 0.1 to 50 µg/dose. Several doses may be needed per individual in order to achieve a sufficient immune response and subsequent protection against HCV disease.

#### Detailed description of the invention

More particularly, the present invention contemplates a method for isolating or purifying recombinant HCV single or specific oligomeric envelope protein selected from the group consisting of E1 and/or E2 and/or E1/E2, characterized in that upon lysing the transformed host cells to isolate the recombinantly expressed protein a disulphide bond cleavage or reduction step is carried out with a disulphide bond cleaving agent.

The essence of these 'single or specific oligomeric' envelope proteins of the invention is that they are

free from contaminating proteins and that they are not disulphide bond linked with contaminants.

The proteins according to the present invention are recombinantly expressed in lower or higher eukaryotic cells or in prokaryotes. The recombinant proteins of the present invention are preferably glycosylated and may contain high-mannose-type, hybrid, or complex glycosylations. Preferentially said proteins are expressed from mammalian cell lines as discussed in detail in the Examples section, or in yeast such as in mutant yeast strains also as detailed in the Examples section.

The proteins according to the present invention may be secreted or expressed within components of the cell, such as the ER or the Golgi Apparatus. Preferably, however, the proteins of the present invention bear high-mannose-type glycosylations and are retained in the ER or Golgi Apparatus of mammalian cells or are retained in or secreted from yeast cells, preferably secreted from yeast mutant strains such as the *mn9* mutant (Kniskern et al., 1994), or from mutants that have been selected by means of vanadate resistance (Ballou et al., 1991).

Upon expression of HCV envelope proteins, the present inventors could show that some of the free thiol groups of cysteines not involved in intra- or inter-molecular disulphide bridges, react with cysteines of host or expression-system-derived (e.g. vaccinia) proteins or of other HCV envelope proteins (single or oligomeric), and form aspecific intermolecular bridges. This results in the formation of 'aggregates' of HCV envelope proteins together with contaminating proteins. It was also shown in WO 92/08734 that 'aggregates' were obtained after purification, but it was not described which protein interactions were involved. In patent application WO 92/08734, recombinant E1/E2 protein expressed with the vaccinia virus system were partially purified as aggregates and only found to be 70% pure, rendering the purified aggregates not useful for diagnostic, prophylactic or therapeutic purposes.

Therefore, a major aim of the present invention resides in the separation of single or specific-oligomeric HCV envelope proteins from contaminating proteins, and to use the purified proteins (> 95% pure) for diagnostic, prophylactic and therapeutic purposes. To those purposes, the present inventors have been able to provide evidence that aggregated protein complexes ('aggregates') are formed on the basis of disulphide bridges and non-covalent protein-protein interactions. The present invention thus provides a means for selectively cleaving the disulphide bonds under specific conditions and for separating the cleaved proteins from contaminating proteins which greatly interfere with diagnostic, prophylactic and therapeutic applications. The free thiol groups may be blocked (reversibly or irreversibly) in order to prevent the reformation of disulphide bridges, or may be left to oxidize and oligomerize with other envelope proteins (see definition homo-oligomer). It is to be understood that such protein oligomers are essentially different from the 'aggregates' described in WO 92/08734 and WO 94/01778, since the level of contaminating proteins is undetectable.

Said disulphide bond cleavage may also be achieved by:

(1) performic acid oxidation by means of cysteic acid in which case the cysteine residues are modified into



cysteic acid (Moore et al., 1963).

(2) Sulfitolysis ( $R-S-S-R \rightarrow 2 R-SO_3^-$ ) for example by means of sulphite ( $SO_3^{2-}$ ) together with a proper oxidant such as  $Cu^{2+}$  in which case the cysteine is modified into S-sulpho-cysteine (Bailey and Cole, 1959).

(3) Reduction by means of mercaptans, such as dithiothreitol (DTT),  $\beta$ -mercapto-ethanol, cysteine, glutathione Red,  $\epsilon$ -mercapto-ethylamine, or thioglycolic acid, of which DTT and  $\beta$ -mercapto-ethanol are commonly used (Cleland, 1964), is the preferred method of this invention because the method can be performed in a water environment and because the cysteine remains unmodified.

(4) Reduction by means of a phosphine (e.g.  $Bu_3P$ ) (Ruegg and Rudinger, 1977).

All these compounds are thus to be regarded as agents or means for cleaving disulphide bonds according to the present invention.

Said disulphide bond cleavage (or reducing) step of the present invention is preferably a partial disulphide bond cleavage (reducing) step (carried out under partial cleavage or reducing conditions).

A preferred disulphide bond cleavage or reducing agent according to the present invention is dithiothreitol (DTT). Partial reduction is obtained by using a low concentration of said reducing agent, i.e. for DTT for example in the concentration range of about 0.1 to about 50 mM; preferably about 0.1 to about 20 mM, preferably about 0.5 to about 10 mM, preferably more than 1 mM, more than 2 mM or more than 5 mM, more preferably about 1.5 mM, about 2.0 mM, about 2.5 mM, about 5 mM or about 7.5 mM.

Said disulphide bond cleavage step may also be carried out in the presence of a suitable detergent (as an example of a means for cleaving disulphide bonds or in combination with a cleaving agent) able to dissociate the expressed proteins, such as DecylPEG, EMPIGEN-BB, NP-40, sodium cholate, Triton X-100.


Said reduction or cleavage step (preferably a partial reduction or cleavage step) is carried out preferably in the presence of (with) a detergent. A preferred detergent according to the present invention is Empigen-BB. The amount of detergent used is preferably in the range of 1 to 10 %, preferably more than 3%, more preferably about 3.5% of a detergent such as Empigen-BB.

A particularly preferred method for obtaining disulphide bond cleavage employs a combination of a classical disulphide bond cleavage agent as detailed above and a detergent (also as detailed above). As contemplated in the Examples section, the particular combination of a low concentration of DTT (1.5 to 7.5 mM) and about 3.5 % of Empigen-BB is proven to be a particularly preferred combination of reducing agent and detergent for the purification of recombinantly expressed E1 and E2 proteins. Upon gel filtration chromatography, said partial reduction is shown to result in the production of possibly dimeric E1 protein and separation of this E1 protein from contaminating proteins that cause false reactivity upon use in immunoassays.

It is, however, to be understood that also any other combination of any reducing agent known in the art with any detergent or other means known in the art to make the cysteines better accessible is also within the scope of the present invention, insofar as said combination reaches the same goal of disulphide bridge cleavage

as the preferred combination exemplified in the present invention.

Apart from reducing the disulphide bonds, a disulphide bond cleaving means according to the present invention may also include any disulphide bridge exchanging agents (competitive agent being either organic or proteinaceous, see for instance Creighton, 1988) known in the art which allows the following type of reaction to occur:



\* R1, R2: compounds of protein aggregates

\* R3 SH: competitive agent (organic, proteinaceous)

The term 'disulphide bridge exchanging agent' is to be interpreted as including disulphide bond reforming as well as disulphide bond blocking agents.

The present invention also relates to methods for purifying or isolating HCV single or specific oligomeric envelope proteins as set out above further including the use of any SH group blocking or binding reagent known in the art such as chosen from the following list:

- Glutathion
- 15 - 5,5'-dithiobis-(2-nitrobenzoic acid) or bis-(3-carboxy-4-nitrophenyl)-disulphide (DTNB or Ellman's reagent) (Elmann, 1959)
- N-ethylmaleimide (NEM; Benesch et al., 1956)
- N-(4-dimethylamino-3,5-dinitrophenyl) maleimide or Tuppy's maleimide which provides a color to the protein
- 20 - P-chloromercuribenzoate (Grassetti et al., 1969)
- 4-vinylpyridine (Friedman and Krull, 1969) can be liberated after reaction by acid hydrolysis
- acrylonitrile, can be liberated after reaction by acid hydrolysis (Weil and Seibles, 1961)
- NEM-biotin (e.g. obtained from Sigma 81267)
- 2,2'-dithiopyridine (Grassetti and Murray, 1967)
- 25 - 4,4'-dithiopyridine (Grassetti and Murray, 1967)
- 6,6'-dithiodinicotinic acid (DTDNA; Brown and Cunningham, 1970)
- 2,2'-dithiobis-(5'-nitropyridine) (DTNP; US patent 3597160) or other dithiobis (heterocyclic derivative) compounds (Grassetti and Murray, 1969)

A survey of the publications cited shows that often different reagents for sulphhydryl groups will react with varying numbers of thiol groups of the same protein or enzyme molecule. One may conclude that this variation in reactivity of the thiol groups is due to the steric environment of these groups, such as the shape of the molecule and the surrounding groups of atoms and their charges, as well as to the size, shape and charge of the reagent molecule or ion. Frequently the presence of adequate concentrations of denaturants such as sodium dodecylsulfate, urea or guanidine hydrochloride will cause sufficient unfolding of the protein molecule to permit

equal access to all of the reagents for thiol groups. By varying the concentration of denaturant, the degree of unfolding can be controlled and in this way thiol groups with different degrees of reactivity may be revealed. Although up to date most of the work reported has been done with p-chloromercuribenzoate, N-ethylmaleimide and DTNB, it is likely that the other more recently developed reagents may prove equally useful. Because of  
5 their varying structures, it seems likely, in fact, that they may respond differently to changes in the steric environment of the thiol groups.

Alternatively, conditions such as low pH (preferably lower than pH 6) for preventing free SH groups from oxidizing and thus preventing the formation of large intermolecular aggregates upon recombinant expression and purification of E1 and E2 (envelope) proteins are also within the scope of the present invention.

10 A preferred SH group blocking reagent according to the present invention is N-ethylmaleimide (NEM). Said SH group blocking reagent may be administered during lysis of the recombinant host cells and after the above-mentioned partial reduction process or after any other process for cleaving disulphide bridges. Said SH group blocking reagent may also be modified with any group capable of providing a detectable label and/or any group aiding in the immobilization of said recombinant protein to a solid substrate, e.g. biotinylated NEM.

15 Methods for cleaving cysteine bridges and blocking free cysteines have also been described in Darbre (1987), Means and Feeney (1971), and by Wong (1993).

A method to purify single or specific oligomeric recombinant E1 and/or E2 and/or E1/E2 proteins according to the present invention as defined above is further characterized as comprising the following steps:

- lysing recombinant E1 and/or E2 and/or E1/E2 expressing host cells, preferably in the presence of an  
20 SH group blocking agent, such as N-ethylmaleimide (NEM), and possibly a suitable detergent, preferably Empigen-BB,
- recovering said HCV envelope protein by affinity purification for instance by means lectin-chromatography, such as lentil-lectin chromatography, or immunoaffinity chromatography using anti-E1 and/or anti-E2 specific monoclonal antibodies, followed by,
- 25 - reduction or cleavage of disulphide bonds with a disulphide bond cleaving agent, such as DTT, preferably also in the presence of an SH group blocking agent, such as NEM or Biotin-NEM, and,
- recovering the reduced HCV E1 and/or E2 and/or E1/E2 envelope proteins for instance by gel filtration (size exclusion chromatography or molecular sieving) and possibly also by an additional  $\text{Ni}^{2+}$ -IMAC chromatography and desalting step.

30 It is to be understood that the above-mentioned recovery steps may also be carried out using any other suitable technique known by the person skilled in the art.

Preferred lectin-chromatography systems include Galanthus nivalis agglutinin (GNA) - chromatography, or Lens culinaris agglutinin (LCA) (lentil) lectin chromatography as illustrated in the Examples section. Other useful lectins include those recognizing high-mannose type sugars, such as Narcissus

pseudonarcissus agglutinin (NPA), Pisum sativum agglutinin (PSA), or Allium ursinum agglutinin (AUA).

Preferably said method is usable to purify single or specific oligomeric HCV envelope protein produced intracellularly as detailed above.

For secreted E1 or E2 or E1/E2 oligomers, lectins binding complex sugars such as Ricinus communis agglutinin I (RCA I), are preferred lectins.

The present invention more particularly contemplates essentially purified recombinant HCV single or specific oligomeric envelope proteins, selected from the group consisting of E1 and/or E2 and/or E1/E2, characterized as being isolated or purified by a method as defined above.

The present invention more particularly relates to the purification or isolation of recombinant envelope proteins which are expressed from recombinant mammalian cells such as vaccinia.

The present invention also relates to the purification or isolation of recombinant envelope proteins which are expressed from recombinant yeast cells.

The present invention equally relates to the purification or isolation of recombinant envelope proteins which are expressed from recombinant bacterial (prokaryotic) cells.

The present invention also contemplates a recombinant vector comprising a vector sequence, an appropriate prokaryotic, eukaryotic or viral or synthetic promoter sequence followed by a nucleotide sequence allowing the expression of the single or specific oligomeric E1 and/or E2 and/or E1/E2 of the invention.

Particularly, the present invention contemplates a recombinant vector comprising a vector sequence, an appropriate prokaryotic, eukaryotic or viral or synthetic promoter sequence followed by a nucleotide sequence allowing the expression of the single E1 or E2 of the invention.

Particularly, the present invention contemplates a recombinant vector comprising a vector sequence, an appropriate prokaryotic, eukaryotic or viral or synthetic promoter sequence followed by a nucleotide sequence allowing the expression of the single E1 or E2 of the invention.

The segment of the HCV cDNA encoding the desired E1 and/or E2 sequence inserted into the vector sequence may be attached to a signal sequence. Said signal sequence may be that from a non-HCV source, e.g. the IgG or tissue plasminogen activator (tpa) leader sequence for expression in mammalian cells, or the  $\alpha$ -mating factor sequence for expression into yeast cells, but particularly preferred constructs according to the present invention contain signal sequences appearing in the HCV genome before the respective start points of the E1 and E2 proteins. The segment of the HCV cDNA encoding the desired E1 and/or E2 sequence inserted into the vector may also include deletions e.g. of the hydrophobic domain(s) as illustrated in the examples section, or of the E2 hypervariable region I.

More particularly, the recombinant vectors according to the present invention encompass a nucleic acid having an HCV cDNA segment encoding the polyprotein starting in the region between amino acid positions 1 and 192 and ending in the region between positions 250 and 400 of the HCV polyprotein, more preferably

ending in the region between positions 250 and 341, even more preferably ending in the region between positions 290 and 341 for expression of the HCV single E1 protein. Most preferably, the present recombinant vector encompasses a recombinant nucleic acid having a HCV cDNA segment encoding part of the HCV polyprotein starting in the region between positions 117 and 192, and ending at any position in the region between positions 263 and 326, for expression of HCV single E1 protein. Also within the scope of the present invention are forms that have the first hydrophobic domain deleted (positions 264 to 293 plus or minus 8 amino acids), or forms to which a 5'-terminal ATG codon and a 3'-terminal stop codon has been added, or forms which have a factor Xa cleavage site and/or 3 to 10, preferably 6 Histidine codons have been added.

More particularly, the recombinant vectors according to the present invention encompass a nucleic acid having an HCV cDNA segment encoding the polyprotein starting in the region between amino acid positions 290 and 406 and ending in the region between positions 600 and 820 of the HCV polyprotein, more preferably starting in the region between positions 322 and 406, even more preferably starting in the region between positions 347 and 406, even still more preferably starting in the region between positions 364 and 406 for expression of the HCV single E2 protein. Most preferably, the present recombinant vector encompasses a recombinant nucleic acid having a HCV cDNA segment encoding the polyprotein starting in the region between positions 290 and 406, and ending at any position of positions 623, 650, 661, 673, 710, 715, 720, 746 or 809, for expression of HCV single E2 protein. Also within the scope of the present invention are forms to which a 5'-terminal ATG codon and a 3'-terminal stop codon has been added, or forms which have a factor Xa cleavage site and/or 3 to 10, preferably 6 Histidine codons have been added.

A variety of vectors may be used to obtain recombinant expression of HCV single or specific oligomeric envelope proteins of the present invention. Lower eukaryotes such as yeasts and glycosylation mutant strains are typically transformed with plasmids, or are transformed with a recombinant virus. The vectors may replicate within the host independently, or may integrate into the host cell genome.

Higher eukaryotes may be transformed with vectors, or may be infected with a recombinant virus, for example a recombinant vaccinia virus. Techniques and vectors for the insertion of foreign DNA into vaccinia virus are well known in the art, and utilize, for example homologous recombination. A wide variety of viral promoter sequences, possibly terminator sequences and poly(A)-addition sequences, possibly enhancer sequences and possibly amplification sequences, all required for the mammalian expression, are available in the art. Vaccinia is particularly preferred since vaccinia halts the expression of host cell proteins. Vaccinia is also very much preferred since it allows the expression of E1 and E2 proteins of HCV in cells or individuals which are immunized with the live recombinant vaccinia virus. For vaccination of humans the avipox and Ankara Modified Virus (AMV) are particularly useful vectors.

Also known are insect expression transfer vectors derived from baculovirus *Autographa californica* nuclear polyhedrosis virus (AcNPV), which is a helper-independent viral expression vector. Expression vectors

derived from this system usually use the strong viral polyhedrin gene promoter to drive the expression of heterologous genes. Different vectors as well as methods for the introduction of heterologous DNA into the desired site of baculovirus are available to the man skilled in the art for baculovirus expression. Also different signals for posttranslational modification recognized by insect cells are known in the art.

5 Also included within the scope of the present invention is a method for producing purified recombinant single or specific oligomeric HCV E1 or E2 or E1:E2 proteins, wherein the cysteine residues involved in aggregates formation are replaced at the level of the nucleic acid sequence by other residues such that aggregate formation is prevented. The recombinant proteins expressed by recombinant vectors carrying such a mutated E1 and/or E2 protein encoding nucleic acid are also within the scope of the present invention.

10 The present invention also relates to recombinant E1 and/or E2 and/or E1/E2 proteins characterized in that at least one of their glycosylation sites has been removed and are consequently termed glycosylation mutants. As explained in the Examples section, different glycosylation mutants may be desired to diagnose (screening, confirmation, prognosis, etc.) and prevent HCV disease according to the patient in question. An E2 protein glycosylation mutant lacking the GLY4 has for instance been found to improve the reactivity of certain  
15 sera in diagnosis. These glycosylation mutants are preferably purified according to the method disclosed in the present invention. Also contemplated within the present invention are recombinant vectors carrying the nucleic acid insert encoding such a E1 and/or E2 and/or E1:E2 glycosylation mutant as well as host cells transformed with such a recombinant vector.

The present invention also relates to recombinant vectors including a polynucleotide which also forms  
20 part of the present invention. The present invention relates more particularly to the recombinant nucleic acids as represented in SEQ ID NO 3, 5, 7, 9, 11, 13, 21, 23, 25, 27, 29, 31, 35, 37, 39, 41, 43, 45, 47 and 49, or parts thereof.

The present invention also contemplates host cells transformed with a recombinant vector as defined above, wherein said vector comprises a nucleotide sequence encoding HCV E1 and/or E2 and/or E1/E2 protein  
25 as defined above in addition to a regulatory sequence operably linked to said HCV E1 and/or E2 and/or E1/E2 sequence and capable of regulating the expression of said HCV E1 and/or E2 and/or E1/E2 protein.

Eukaryotic hosts include lower and higher eukaryotic hosts as described in the definitions section. Lower eukaryotic hosts include yeast cells well known in the art. Higher eukaryotic hosts mainly include mammalian cell lines known in the art and include many immortalized cell lines available from the ATCC,  
30 including HeLa cells, Chinese hamster ovary (CHO) cells, Baby hamster kidney (BHK) cells, PK15, RK13 and a number of other cell lines.

The present invention relates particularly to a recombinant E1 and/or E2 and/or E1/E2 protein expressed by a host cell as defined above containing a recombinant vector as defined above. These recombinant proteins are particularly purified according to the method of the present invention.

A preferred method for isolating or purifying HCV envelope proteins as defined above is further characterized as comprising at least the following steps:

- growing a host cell as defined above transformed with a recombinant vector according to the present invention or with a known recombinant vector expressing E1 and/or E2 and/or E1/E2 HCV envelope proteins in a suitable culture medium,
- causing expression of said vector sequence as defined above under suitable conditions, and,
- lysing said transformed host cells, preferably in the presence of a SH group blocking agent, such as N-ethylmaleimide (NEM), and possibly a suitable detergent, preferably Empigen-BB,
- recovering said HCV envelope protein by affinity purification such as by means of lectin-chromatography or immunoaffinity chromatography using anti-E1 and/or anti-E2 specific monoclonal antibodies, with said lectin being preferably lentil-lectin or GNA, followed by,
- incubation of the eluate of the previous step with a disulphide bond cleavage means, such as DTT, preferably followed by incubation with an SH group blocking agent, such as NEM or Biotin-NEM, and,
- isolating the HCV single or specific oligomeric E1 and/or E2 and/or E1/E2 proteins such as by means of gel-filtration and possibly also by a subsequent  $\text{Ni}^{2+}$ -IMAC chromatography followed by a desalting step.

As a result of the above-mentioned process, E1 and/or E2 and/or E1/E2 proteins may be produced in a form which elute differently from the large aggregates containing vector-derived components and/or cell components in the void volume of the gel-filtration column or the IMAC column as illustrated in the Examples section. The disulphide bridge cleavage step advantageously also eliminates the false reactivity due to the presence of host and/or expression-system-derived proteins. The presence of NEM and a suitable detergent during lysis of the cells may already partly or even completely prevent the aggregation between the HCV envelope proteins and contaminants.

$\text{Ni}^{2+}$ -IMAC chromatography followed by a desalting step is preferably used for constructs bearing a (His)<sub>6</sub> as described by Janknecht et al., 1991, and Hochuli et al., 1988.

The present invention also relates to a method for producing monoclonal antibodies in small animals such as mice or rats, as well as a method for screening and isolating human B-cells that recognize anti-HCV antibodies, using the HCV single or specific oligomeric envelope proteins of the present invention.

The present invention further relates to a composition comprising at least one of the following E1 peptides as listed in Table 3:

- E1-31 (SEQ ID NO 56) spanning amino acids 181 to 200 of the Core/E1 V1 region,
- E1-33 (SEQ ID NO 57) spanning amino acids 193 to 212 of the E1 region,
- E1-35 (SEQ ID NO 58) spanning amino acids 205 to 224 of the E1 V2 region (epitope B),
- E1-35A (SEQ ID NO 59) spanning amino acids 208 to 227 of the E1 V2 region (epitope B).

1bE1 (SEQ ID NO 53) spanning amino acids 192 to 228 of E1 regions (V1, C1, and V2 regions (containing epitope B)),

E1-51 (SEQ ID NO 66) spanning amino acids 301 to 320 of the E1 region,

E1-53 (SEQ ID NO 67) spanning amino acids 313 to 332 of the E1 C4 region (epitope A),

5 E1-55 (SEQ ID NO 68) spanning amino acids 325 to 344 of the E1 region.

The present invention also relates to a composition comprising at least one of the following E2 peptides as listed in Table 3:

Env 67 or E2-67 (SEQ ID NO 72) spanning amino acid positions 397 to 416 of the E2 region (epitope A, recognized by monoclonal antibody 2F10H10, see Figure 19),

10 Env 69 or E2-69 (SEQ ID NO 73) spanning amino acid positions 409 to 428 of the E2 region (epitope A),

Env 23 or E2-23 (SEQ ID NO 86) spanning positions 583 to 602 of the E2 region (epitope E),

Env 25 or E2-25 (SEQ ID NO 87) spanning positions 595 to 614 of the E2 region (epitope E),

Env 27 or E2-27 (SEQ ID NO 88) spanning positions 607 to 626 of the E2 region (epitope E),

15 Env 17B or E2-17B (SEQ ID NO 83) spanning positions 547 to 566 of the E2 region (epitope D),

Env 13B or E2-13B (SEQ ID NO 82) spanning positions 523 to 542 of the E2 region (epitope C; recognized by monoclonal antibody 16A6E7, see Figure 19).

The present invention also relates to a composition comprising at least one of the following E2 conformational epitopes:

20 epitope F recognized by monoclonal antibodies 15C8C1, 12D11F1 and 8G10D1H9,

epitope G recognized by monoclonal antibody 9G3E3,

epitope H (or C) recognized by monoclonal antibody 10D3C4 and 4H6B2, or,

epitope I recognized by monoclonal antibody 17F2C2.

The present invention also relates to an E1 or E2 specific antibody raised upon immunization with a peptide or protein composition, with said antibody being specifically reactive with any of the polypeptides or peptides as defined above, and with said antibody being preferably a monoclonal antibody.

The present invention also relates to an E1 or E2 specific antibody screened from a variable chain library in plasmids or phages or from a population of human B-cells by means of a process known in the art, with said antibody being reactive with any of the polypeptides or peptides as defined above, and with said antibody being preferably a monoclonal antibody.

30 The E1 or E2 specific monoclonal antibodies of the invention can be produced by any hybridoma liable to be formed according to classical methods from splenic cells of an animal, particularly from a mouse or rat immunized against the HCV polypeptides or peptides according to the invention, as defined above on the one hand, and of cells of a myeloma cell line on the other hand, and to be selected by the ability of the hybridoma to



produce the monoclonal antibodies recognizing the polypeptides which has been initially used for the immunization of the animals.

The antibodies involved in the invention can be labelled by an appropriate label of the enzymatic, fluorescent, or radioactive type.

5 The monoclonal antibodies according to this preferred embodiment of the invention may be humanized versions of mouse monoclonal antibodies made by means of recombinant DNA technology, departing from parts of mouse and/or human genomic DNA sequences coding for H and L chains from cDNA or genomic clones coding for H and L chains.

10 Alternatively the monoclonal antibodies according to this preferred embodiment of the invention may be human monoclonal antibodies. These antibodies according to the present embodiment of the invention can also be derived from human peripheral blood lymphocytes of patients infected with HCV, or vaccinated against HCV. Such human monoclonal antibodies are prepared, for instance, by means of human peripheral blood lymphocytes (PBL) repopulation of severe combined immune deficiency (SCID) mice (for recent review, see Duchosal et al., 1992).

15 The invention also relates to the use of the proteins or peptides of the invention, for the selection of recombinant antibodies by the process of repertoire cloning (Persson et al., 1991).

Antibodies directed to peptides or single or specific oligomeric envelope proteins derived from a certain genotype may be used as a medicament, more particularly for incorporation into an immunoassay for the detection of HCV genotypes (for detecting the presence of HCV E1 or E2 antigen), for prognosing/monitoring of HCV disease, or as therapeutic agents.

20 Alternatively, the present invention also relates to the use of any of the above-specified E1 or E2 specific monoclonal antibodies for the preparation of an immunoassay kit for detecting the presence of E1 or E2 antigen in a biological sample, for the preparation of a kit for prognosing/monitoring of HCV disease or for the preparation of a HCV medicament.

25 The present invention also relates to the a method for *in vitro* diagnosis or detection of HCV antigen present in a biological sample, comprising at least the following steps :

- (i) contacting said biological sample with any of the E1 and/or E2 specific monoclonal antibodies as defined above, preferably in an immobilized form under appropriate conditions which allow the formation of an immune complex,
- 30 (ii) removing unbound components,
- (iii) incubating the immune complexes formed with heterologous antibodies, which specifically bind to the antibodies present in the sample to be analyzed, with said heterologous antibodies having conjugated to a detectable label under appropriate conditions,
- (iv) detecting the presence of said immune complexes visually or mechanically (e.g. by means of

densitometry, fluorimetry, colorimetry).

The present invention also relates to a kit for in vitro diagnosis of HCV antigen present in a biological sample, comprising:

- 5                   - at least one monoclonal antibody as defined above, with said antibody being preferentially immobilized on a solid substrate,
- a buffer or components necessary for producing the buffer enabling binding reaction between these antibodies and the HCV antigens present in the biological sample,
- a means for detecting the immune complexes formed in the preceding binding reaction,
- 10                  - possibly also including an automated scanning and interpretation device for inferring the HCV antigens present in the sample from the observed binding pattern.

The present invention also relates to a composition comprising E1 and/or E2 and/or E1/E2 recombinant HCV proteins purified according to the method of the present invention or a composition comprising at least one peptides as specified above for use as a medicament.

15                  The present invention more particularly relates to a composition comprising at least one of the above-specified envelope peptides or a recombinant envelope protein composition as defined above, for use as a vaccine for immunizing a mammal, preferably humans, against HCV, comprising administering a sufficient amount of the composition possibly accompanied by pharmaceutically acceptable adjuvant(s), to produce an immune response.

20                  More particularly, the present invention relates to the use of any of the compositions as described here above for the preparation of a vaccine as described above.

Also, the present invention relates to a vaccine composition for immunizing a mammal, preferably humans, against HCV, comprising HCV single or specific oligomeric proteins or peptides derived from the E1 and/or the E2 region as described above.

25                  Immunogenic compositions can be prepared according to methods known in the art. The present compositions comprise an immunogenic amount of a recombinant E1 and/or E2 and/or E1/E2 single or specific oligomeric proteins as defined above or E1 or E2 peptides as defined above, usually combined with a pharmaceutically acceptable carrier, preferably further comprising an adjuvant.

30                  The single or specific oligomeric envelope proteins of the present invention, either E1 and/or E2 and/or E1/E2, are expected to provide a particularly useful vaccine antigen, since the formation of antibodies to either E1 or E2 may be more desirable than to the other envelope protein, and since the E2 protein is cross-reactive between HCV types and the E1 protein is type-specific. Cocktails including type 1 E2 protein and E1 proteins derived from several genotypes may be particularly advantageous. Cocktails containing a molar excess of E1 versus E2 or E2 versus E1 may also be particularly useful. Immunogenic compositions may be administered to animals to induce production of antibodies, either to provide a source of antibodies or to induce protective

immunity in the animal.

Pharmaceutically acceptable carriers include any carrier that does not itself induce the production of antibodies harmful to the individual receiving the composition. Suitable carriers are typically large, slowly metabolized macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers; and inactive virus particles. Such carriers are well known to those of ordinary skill in the art.

Preferred adjuvants to enhance effectiveness of the composition include, but are not limited to : aluminum hydroxide (alum), N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP) as found in U.S. Patent No. 4,606,918, N-acetyl-normuramyl-L-alanyl-D-isoglutamine (nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (MTP-PE) and RIBI, which contains three components extracted from bacteria, monophosphoryl lipid A, trehalose dimycolate, and cell wall skeleton (MPL+TDM+CWS) in a 2% squalene:Tween 80 emulsion. Any of the 3 components MPL, TDM or CWS may also be used alone or combined 2 by 2. Additionally, adjuvants such as Stimulon (Cambridge Bioscience, Worcester, MA) or SAF-1 (Syntex) may be used. Further, Complete Freund's Adjuvant (CFA) and Incomplete Freund's Adjuvant (IFA) may be used for non-human applications and research purposes.

The immunogenic compositions typically will contain pharmaceutically acceptable vehicles, such as water, saline, glycerol, ethanol, etc. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, preservatives, and the like, may be included in such vehicles.

Typically, the immunogenic compositions are prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid vehicles prior to injection may also be prepared. The preparation also may be emulsified or encapsulated in liposomes for enhanced adjuvant effect. The E1 and E2 proteins may also be incorporated into Immune Stimulating Complexes together with saponins, for example Quil A (ISCOMS).

Immunogenic compositions used as vaccines comprise a 'sufficient amount' or 'an immunologically effective amount' of the envelope proteins of the present invention, as well as any other of the above mentioned components, as needed. 'Immunologically effective amount', means that the administration of that amount to an individual, either in a single dose or as part of a series, is effective for treatment, as defined above. This amount varies depending upon the health and physical condition of the individual to be treated, the taxonomic group of individual to be treated (e.g. nonhuman primate, primate, etc.), the capacity of the individual's immune system to synthesize antibodies, the degree of protection desired, the formulation of the vaccine, the treating doctor's assessment of the medical situation, the strain of infecting HCV, and other relevant factors. It is expected that the amount will fall in a relatively broad range that can be determined through routine trials. Usually, the amount will vary from 0.01 to 1000  $\mu\text{g}/\text{dose}$ , more particularly from 0.1 to 100  $\mu\text{g}/\text{dose}$ .

The single or specific oligomeric envelope proteins may also serve as vaccine carriers to present

homologous (e.g. T cell epitopes or B cell epitopes from the core, NS2, NS3, NS4 or NS5 regions) or heterologous (non-HCV) haptens, in the same manner as Hepatitis B surface antigen (see European Patent Application 174,444). In this use, envelope proteins provide an immunogenic carrier capable of stimulating an immune response to haptens or antigens conjugated to the aggregate. The antigen may be conjugated either by  
5 conventional chemical methods, or may be cloned into the gene encoding E1 and/or E2 at a location corresponding to a hydrophilic region of the protein. Such hydrophilic regions include the V1 region (encompassing amino acid positions 191 to 202), the V2 region (encompassing amino acid positions 213 to 223), the V3 region (encompassing amino acid positions 230 to 242), the V4 region (encompassing amino acid positions 230 to 242), the V5 region (encompassing amino acid positions 294 to 303) and the V6 region  
10 (encompassing amino acid positions 329 to 336). Another useful location for insertion of haptens is the hydrophobic region (encompassing approximately amino acid positions 264 to 293). It is shown in the present invention that this region can be deleted without affecting the reactivity of the deleted E1 protein with antisera. Therefore, haptens may be inserted at the site of the deletion.

The immunogenic compositions are conventionally administered parenterally, typically by injection, for  
15 example, subcutaneously or intramuscularly. Additional formulations suitable for other methods of administration include oral formulations and suppositories. Dosage treatment may be a single dose schedule or a multiple dose schedule. The vaccine may be administered in conjunction with other immunoregulatory agents.

The present invention also relates to a composition comprising peptides or polypeptides as described above, for *in vitro* detection of HCV antibodies present in a biological sample.

20 The present invention also relates to the use of a composition as described above for the preparation of an immunoassay kit for detecting HCV antibodies present in a biological sample.

The present invention also relates to a method for *in vitro* diagnosis of HCV antibodies present in a biological sample, comprising at least the following steps :

- 25 (i) contacting said biological sample with a composition comprising any of the envelope peptide or proteins as defined above, preferably in an immobilized form under appropriate conditions which allow the formation of an immune complex, wherein said peptide or protein can be a biotinylated peptide or protein which is covalently bound to a solid substrate by means of streptavidin or avidin complexes,
- (ii) removing unbound components,
- 30 (iii) incubating the immune complexes formed with heterologous antibodies, with said heterologous antibodies having conjugated to a detectable label under appropriate conditions,
- (iv) detecting the presence of said immune complexes visually or mechanically (e.g. by means of densitometry, fluorimetry, colorimetry).

Alternatively, the present invention also relates to competition immunoassay formats in which

recombinantly produced purified single or specific oligomeric protein E1 and/or E2 and/or E1/E2 proteins as disclosed above are used in combination with E1 and/or E2 peptides in order to compete for HCV antibodies present in a biological sample.

The present invention also relates to a kit for determining the presence of HCV antibodies, in a biological sample, comprising :

- at least one peptide or protein composition as defined above, possibly in combination with other polypeptides or peptides from HCV or other types of HCV, with said peptides or proteins being preferentially immobilized on a solid substrate, more preferably on different microwells of the same ELISA plate, and even more preferentially on one and the same membrane strip,
- a buffer or components necessary for producing the buffer enabling binding reaction between these polypeptides or peptides and the antibodies against HCV present in the biological sample,
- means for detecting the immune complexes formed in the preceding binding reaction,
- possibly also including an automated scanning and interpretation device for inferring the HCV genotypes present in the sample from the observed binding pattern.

The immunoassay methods according to the present invention utilize single or specific oligomeric antigens from the E1 and/or E2 domains that maintain linear (in case of peptides) and conformational epitopes (single or specific oligomeric proteins) recognized by antibodies in the sera from individuals infected with HCV. It is within the scope of the invention to use for instance single or specific oligomeric antigens, dimeric antigens, as well as combinations of single or specific oligomeric antigens. The HCV E1 and E2 antigens of the present invention may be employed in virtually any assay format that employs a known antigen to detect antibodies. Of course, a format that denatures the HCV conformational epitope should be avoided or adapted. A common feature of all of these assays is that the antigen is contacted with the body component suspected of containing HCV antibodies under conditions that permit the antigen to bind to any such antibody present in the component. Such conditions will typically be physiologic temperature, pH and ionic strength using an excess of antigen. The incubation of the antigen with the specimen is followed by detection of immune complexes comprised of the antigen.

Design of the immunoassays is subject to a great deal of variation, and many formats are known in the art. Protocols may, for example, use solid supports, or immunoprecipitation. Most assays involve the use of labeled antibody or polypeptide; the labels may be, for example, enzymatic, fluorescent, chemiluminescent, radioactive, or dye molecules. Assays which amplify the signals from the immune complex are also known; examples of which are assays which utilize biotin and avidin or streptavidin, and enzyme-labeled and mediated immunoassays, such as ELISA assays.

The immunoassay may be, without limitation, in a heterogeneous or in a homogeneous format, and of

a standard or competitive type. In a heterogeneous format, the polypeptide is typically bound to a solid matrix or support to facilitate separation of the sample from the polypeptide after incubation. Examples of solid supports that can be used are nitrocellulose (e.g., in membrane or microtiter well form), polyvinyl chloride (e.g., in sheets or microtiter wells), polystyrene latex (e.g., in beads or microtiter plates), polyvinylidene fluoride (known as Immunolon™), diazotized paper, nylon membranes, activated beads, and Protein A beads. For example, Dynatech Immunolon™ 1 or Immunolon™ 2 microtiter plates or 0.25 inch polystyrene beads (Precision Plastic Ball) can be used in the heterogeneous format. The solid support containing the antigenic polypeptides is typically washed after separating it from the test sample, and prior to detection of bound antibodies. Both standard and competitive formats are known in the art.

In a homogeneous format, the test sample is incubated with the combination of antigens in solution. For example, it may be under conditions that will precipitate any antigen-antibody complexes which are formed. Both standard and competitive formats for these assays are known in the art.

In a standard format, the amount of HCV antibodies in the antibody-antigen complexes is directly monitored. This may be accomplished by determining whether labeled anti-xenogeneic (e.g. anti-human) antibodies which recognize an epitope on anti-HCV antibodies will bind due to complex formation. In a competitive format, the amount of HCV antibodies in the sample is deduced by monitoring the competitive effect on the binding of a known amount of labeled antibody (or other competing ligand) in the complex.

Complexes formed comprising anti-HCV antibody (or in the case of competitive assays, the amount of competing antibody) are detected by any of a number of known techniques, depending on the format. For example, unlabeled HCV antibodies in the complex may be detected using a conjugate of anti-xenogeneic Ig complexed with a label (e.g. an enzyme label).

In an immunoprecipitation or agglutination assay format the reaction between the HCV antigens and the antibody forms a network that precipitates from the solution or suspension and forms a visible layer or film of precipitate. If no anti-HCV antibody is present in the test specimen, no visible precipitate is formed.

There currently exist three specific types of particle agglutination (PA) assays. These assays are used for the detection of antibodies to various antigens when coated to a support. One type of this assay is the hemagglutination assay using red blood cells (RBCs) that are sensitized by passively adsorbing antigen (or antibody) to the RBC. The addition of specific antigen antibodies present in the body component, if any, causes the RBCs coated with the purified antigen to agglutinate.

To eliminate potential non-specific reactions in the hemagglutination assay, two artificial carriers may be used instead of RBC in the PA. The most common of these are latex particles. However, gelatin particles may also be used. The assays utilizing either of these carriers are based on passive agglutination of the particles coated with purified antigens.

The HCV single or specific oligomeric E1 and/or E2 and/or E1/E2 antigens of the present invention

comprised of conformational epitopes will typically be packaged in the form of a kit for use in these immunoassays. The kit will normally contain in separate containers the native HCV antigen, control antibody formulations (positive and/or negative), labeled antibody when the assay format requires the same and signal generating reagents (e.g. enzyme substrate) if the label does not generate a signal directly. The native HCV antigen may be already bound to a solid matrix or separate with reagents for binding it to the matrix. Instructions (e.g. written, tape, CD-ROM, etc.) for carrying out the assay usually will be included in the kit.

Immunoassays that utilize the native HCV antigen are useful in screening blood for the preparation of a supply from which potentially infective HCV is lacking. The method for the preparation of the blood supply comprises the following steps. Reacting a body component, preferably blood or a blood component, from the individual donating blood with HCV E1 and/or E2 proteins of the present invention to allow an immunological reaction between HCV antibodies, if any, and the HCV antigen. Detecting whether anti-HCV antibody - HCV antigen complexes are formed as a result of the reacting. Blood contributed to the blood supply is from donors that do not exhibit antibodies to the native HCV antigens, E1 or E2.

In cases of a positive reactivity to the HCV antigen, it is preferable to repeat the immunoassay to lessen the possibility of false positives. For example, in the large scale screening of blood for the production of blood products (e.g. blood transfusion, plasma, Factor VIII, immunoglobulin, etc.) 'screening' tests are typically formatted to increase sensitivity (to insure no contaminated blood passes) at the expense of specificity; i.e. the false-positive rate is increased. Thus, it is typical to only defer for further testing those donors who are 'repeatedly reactive'; i.e. positive in two or more runs of the immunoassay on the donated sample. However, for confirmation of HCV-positivity, the 'confirmation' tests are typically formatted to increase specificity (to insure that no false-positive samples are confirmed) at the expense of sensitivity. Therefore the purification method described in the present invention for E1 and E2 will be very advantageous for including single or specific oligomeric envelope proteins into HCV diagnostic assays.

The solid phase selected can include polymeric or glass beads, nitrocellulose, microparticles, microwells of a reaction tray, test tubes and magnetic beads. The signal generating compound can include an enzyme, a luminescent compound, a chromogen, a radioactive element and a chemiluminescent compound. Examples of enzymes include alkaline phosphatase, horseradish peroxidase and beta-galactosidase. Examples of enhancer compounds include biotin, anti-biotin and avidin. Examples of enhancer compounds binding members include biotin, anti-biotin and avidin. In order to block the effects of rheumatoid factor-like substances, the test sample is subjected to conditions sufficient to block the effect of rheumatoid factor-like substances. These conditions comprise contacting the test sample with a quantity of anti-human IgG to form a mixture, and incubating the mixture for a time and under conditions sufficient to form a reaction mixture product substantially free of rheumatoid factor-like substance.

The present invention further contemplates the use of E1 proteins, or parts thereof, more particularly

HCV single or specific oligomeric E1 proteins as defined above, for *in vitro* monitoring HCV disease or prognosing the response to treatment (for instance with Interferon) of patients suffering from HCV infection comprising:

- incubating a biological sample from a patient with hepatitis C infection with an E1 protein or a suitable part thereof under conditions allowing the formation of an immunological complex,
- removing unbound components,
- calculating the anti-E1 titers present in said sample (for example at the start of and/or during the course of (interferon) therapy),
- monitoring the natural course of HCV disease, or prognosing the response to treatment of said patient on the basis of the amount anti-E1 titers found in said sample at the start of treatment and/or during the course of treatment.

Patients who show a decrease of 2, 3, 4, 5, 7, 10, 15, or preferably more than 20 times of the initial anti-E1 titers could be concluded to be long-term, sustained responders to HCV therapy, more particularly to interferon therapy. It is illustrated in the Examples section, that an anti-E1 assay may be very useful for prognosing long-term response to IFN treatment, or to treatment of Hepatitis C virus disease in general.

More particularly the following E1 peptides as listed in Table 3 were found to be useful for *in vitro* monitoring HCV disease or prognosing the response to interferon treatment of patients suffering from HCV infection:

- E1-31 (SEQ ID NO 56) spanning amino acids 181 to 200 of the Core/E1 V1 region,
- E1-33 (SEQ ID NO 57) spanning amino acids 193 to 212 of the E1 region,
- E1-35 (SEQ ID NO 58) spanning amino acids 205 to 224 of the E1 V2 region (epitope B),
- E1-35A (SEQ ID NO 59) spanning amino acids 208 to 227 of the E1 V2 region (epitope B),
- 1bE1 (SEQ ID NO 53) spanning amino acids 192 to 228 of E1 regions (V1, C1, and V2 regions (containing epitope B)),
- E1-51 (SEQ ID NO 66) spanning amino acids 301 to 320 of the E1 region,
- E1-53 (SEQ ID NO 67) spanning amino acids 313 to 332 of the E1 C4 region (epitope A),
- E1-55 (SEQ ID NO 68) spanning amino acids 325 to 344 of the E1 region.

It is to be understood that smaller fragments of the above-mentioned peptides also fall within the scope of the present invention. Said smaller fragments can be easily prepared by chemical synthesis and can be tested for their ability to be used in an assay as detailed above and in the Examples section.

The present invention also relates to a kit for monitoring HCV disease or prognosing the response to treatment (for instance to interferon) of patients suffering from HCV infection comprising:

- at least one E1 protein or E1 peptide, more particularly an E1 protein or E1 peptide as defined above,



- a buffer or components necessary for producing the buffer enabling the binding reaction between these proteins or peptides and the anti-E1 antibodies present in a biological sample,
- means for detecting the immune complexes formed in the preceding binding reaction,
- possibly also an automated scanning and interpretation device for inferring a decrease of anti-E1 titers during the progression of treatment.

5

It is to be understood that also E2 protein and peptides according to the present invention can be used to a certain degree to monitor/prognose HCV treatment as indicated above for the E1 proteins or peptides because also the anti-E2 levels decrease in comparison to antibodies to the other HCV antigens. It is to be understood, however, that it might be possible to determine certain epitopes in the E2 region which would also be suited for use in an test for monitoring/prognosing HCV disease.

10

The present invention also relates to a serotyping assay for detecting one or more serological types of HCV present in a biological sample, more particularly for detecting antibodies of the different types of HCV to be detected combined in one assay format, comprising at least the following steps :

15

(i) contacting the biological sample to be analyzed for the presence of HCV antibodies of one or more serological types, with at least one of the E1 and/or E2 and/or E1/E2 protein compositions or at least one of the E1 or E2 peptide compositions as defined above, preferentially in an immobilized form under appropriate conditions which allow the formation of an immune complex,

20

(ii) removing unbound components.

(iii) incubating the immune complexes formed with heterologous antibodies, with said heterologous antibodies being conjugated to a detectable label under appropriate conditions,

(iv) detecting the presence of said immune complexes visually or mechanically (e.g. by means of densitometry, fluorimetry, colorimetry) and inferring the presence of one or more HCV serological types present from the observed binding pattern.

25

It is to be understood that the compositions of proteins or peptides used in this method are recombinantly expressed type-specific envelope proteins or type-specific peptides.

The present invention further relates to a kit for serotyping one or more serological types of HCV present in a biological sample, more particularly for detecting the antibodies to these serological types of HCV comprising:

30

- at least one E1 and/or E2 and/or E1/E2 protein or E1 or E2 peptide, as defined above,
- a buffer or components necessary for producing the buffer enabling the binding reaction between these proteins or peptides and the anti-E1 antibodies present in a biological sample.
- means for detecting the immune complexes formed in the preceding binding reaction,
- possibly also an automated scanning and interpretation device for detecting the presence of

one or more serological types present from the observed binding pattern.

The present invention also relates to the use of a peptide or protein composition as defined above, for immobilization on a solid substrate and incorporation into a reversed phase hybridization assay, preferably for immobilization as parallel lines onto a solid support such as a membrane strip, for determining the presence or  
 5 the genotype of HCV according to a method as defined above. Combination with other type-specific antigens from other HCV polyprotein regions also lies within the scope of the present invention.

The present invention provides a method for purifying recombinant HCV single or specific oligomeric envelope proteins selected from E1 and/or E2 and/or E1/E2 proteins which have been produced by a recombinant process comprising contacting said proteins with a disulphide bond cleavage or reducing agent. The contacting of the method of  
 10 the invention may be carried out under partial cleavage or reducing conditions. Preferably, the disulphide bond cleavage agent is dithiothreitol (DTT), preferably in a concentration range of 0.1 to 50 mM, preferably 0.1 to 20 mM, more preferably 0.5 to 10 mM. Alternatively, the disulphide bond cleavage agent may be a detergent, such as Empigen-BB (which is a mixture containing N-Dodecyl-N,N-dimethylglycine as a major component), preferably at a concentration of 1 to 10%, more preferably at a concentration of 3.5%. Mixtures of detergents, disulphide bond cleavage agents and/or  
 15 reducing agents may also be used. In one embodiment, disulphide bond reformation is prevented with an SH group blocking agent, such as N-ethylmaleimide (NEM) or a derivative thereof. In a preferred embodiment, the disulphide bond reformation is blocked by use of low pH conditions.

The present invention further provides a method as described herein, further involving the following steps:  
 lysing recombinant E1 and/or E2 and/or E1/E2 expressing host cells, optionally in the presence of an SH blocking agent  
 20 such as N-ethylmaleimide (NEM); recovering said HCV envelope proteins by affinity purification such as by means of lectin-chromatography, such as lentil-lectin chromatography, or by means of immunoaffinity using anti-E1 and/or anti-E2 specific monoclonal antibodies; reducing or cleaving of the disulfide bonds with a disulphide bond cleaving agent, such as DTT, preferably also in the presence of an SH blocking agent, such as NEM or Biotin-NEM; and, recovering the reduced E1 and/or E2 and/or E1/E2 envelope proteins by gel filtration and optionally additionally by a subsequent Ni-  
 25 IMAC chromatography and desalting step.

The present invention provides a composition containing substantially isolated and/or purified, and/or isolated and/or purified recombinant HCV single or specific oligomeric recombinant envelope proteins selected from E1 and/or E2 and/or E1/E2, which have preferably been isolated from the methods described herein. In a preferred embodiment, the recombinant HCV envelope proteins of the invention have been expressed in recombinant mammalian cells, such as  
 30 vaccinia, recombinant yeast cells.

The present invention provides a recombinant vector containing a vector sequence, a prokaryotic, eukaryotic or viral promoter sequence and a nucleotide sequence allowing the expression of a single or specific oligomeric E1 and/or E2 and/or E1/E2 protein, in operable combination. In one embodiment, the nucleotide sequence of the vector encodes a single HCV E1 protein starting in the region between amino acid positions 1 and 192 and ending in the region  
 35 between amino acid positions 250 and 400, more particularly ending in the region between positions 250 and 341, even more preferably ending in the region between position 290 and 341. In another embodiment, the nucleotide sequence of the vector encodes a single HCV E1 protein starting in the region between amino acid positions 117 and 192 and ending

in the region between amino acid positions 263 and 400, more particularly ending in the region between positions 250 and 326. In yet another embodiment, the nucleotide sequence of the vector encodes a single HCV E1 protein bearing a deletion of the first hydrophobic domain between positions 264 to 293, plus or minus 8 amino acids. In a further embodiment, the nucleotide sequence of the vector encodes a single HCV E2 protein starting in the region between amino acid positions 290 and 406 and ending in the region between amino acid positions 600 and 820, more particularly starting in the region between positions 322 and 406, even more preferably starting in the region between position 347 and 406 and most preferably starting in the region between positions 364 and 406; and preferably ending at any of amino acid positions 623, 650, 661, 673, 710, 715, 720, 746 or 809. The vector of the present invention, in one embodiment, contains a 5'-terminal ATG codon and a 3'-terminal stop codon operably linked to the nucleotide sequence.

The vector further contains, in one embodiment, a nucleotide sequence further containing at a factor Xa cleavage site and/or 3 to 10, preferably 6, histidine codons added 3'-terminally to the coding region. The vector of the present invention optionally contains a nucleotide sequence wherein at least one of the glycosylation sites present in the E1 or E2 proteins has been removed at the nucleic acid level.

The present invention provides a nucleic acid containing any one of SEQ ID NOs: 3, 5, 7, 9, 11, 13, 21, 23, 25, 27, 29, 31, 35, 37, 39, 41, 43, 45, 47 and 49, or parts thereof. The vector of the invention may preferably contain a nucleotide sequence containing a nucleic acid containing any one of SEQ ID NOs: 3, 5, 7, 9, 11, 13, 21, 23, 25, 27, 29, 31, 35, 37, 39, 41, 43, 45, 47 and 49, or parts thereof.

The composition of the present invention further contains recombinant HCV envelope proteins which have been expressed or are the expression product of a vector described herein.

The present invention provides a host cell transformed with at least one recombinant vector as described herein, wherein the vector contains a nucleotide sequence encoding HCV E1 and/or E2 and/or E1/E2 protein as described herein in addition to a regulatory sequence operable in the host cell and capable of regulating expression of the HCV E1 and/or E2 and/or E1/E2 protein. Moreover, the present invention provides a recombinant E1 and/or E2 and/or E1/E2 protein expressed by a host cell of the invention.

The present invention further provides a method as described herein and containing the following steps: growing a host cell as described herein which has been transformed with a recombinant vector as described herein in a suitable culture medium; causing expression of the vector nucleotide sequence of the vector, as described herein under suitable conditions; lysing the transformed host cells, preferably in the presence of an SH group blocking agent, such as N-ethylmaleimide (NEM); recovering the HCV envelope protein by affinity purification by means of for instance lectin-chromatography or immunoaffinity chromatography using anti-E1 and/or anti-E2 specific monoclonal antibodies, with said lectin being preferably lentil-lectin, followed by, incubation of the eluate of the previous step with a disulphide bond cleavage agent, such as DTT, preferably also in the presence of an SH group blocking agent, such as NEM or Biotin-NEM; and, isolating the HCV single or specific oligomeric E1 and/or E2 and/or E1/E2 proteins by means of gel filtration and possibly also by means of an additional  $\text{Ni}^{2+}$ -IMAC chromatography and desalting step.

The present invention provides a composition containing at least one of the following E1 and/or E2 peptides:

E1-31 (SEQ ID NO 56) spanning amino acids 181 to 200 of the Core/E1 V1 region,

E1-33 (SEQ ID NO 57) spanning amino acids 193 to 212 of the E1 region,

E1-35 (SEQ ID NO 58) spanning amino acids 205 to 224 of the E1 V2 region (epitope B),  
 E1-35A (SEQ ID NO 59) spanning amino acids 208 to 227 of the E1 V2 region (epitope B),  
 1bE1 (SEQ ID NO 53) spanning amino acids 192 to 228 of E1 regions (V1, C1, and V2 regions  
 (containing epitope B),

5

E1-51 (SEQ ID NO 66) spanning amino acids 301 to 320 of the E1 region,  
 E1-53 (SEQ ID NO 67) spanning amino acids 313 to 332 of the E1 C4 region (epitope A),  
 E1-55 (SEQ ID NO 68) spanning amino acids 325 to 344 of the E1 region.  
 Env 67 or E2-67 (SEQ ID NO 72) spanning amino acid positions 397 to 416 of the E2 region (epitope  
 A),

10

Env 69 or E2-69 (SEQ ID NO 73) spanning amino acid positions 409 to 428 of the E2 region (epitope  
 A),

15

Env 23 or E2-23 (SEQ ID NO 86) spanning positions 583 to 602 of the E2 region (epitope E),  
 Env 25 or E2-25 (SEQ ID NO 87) spanning positions 595 to 614 of the E2 region (epitope E),  
 Env 27 or E2-27 (SEQ ID NO 88) spanning positions 607 to 626 of the E2 region (epitope E),  
 Env 17B or E2-17B (SEQ ID NO 83) spanning positions 547 to 566 of the E2 region (epitope D),  
 Env 13B or E2-13B (SEQ ID NO 82) spanning positions 523 to 542 of the E2 region (epitope C).

The present invention provides a composition containing at least one of the following E2 conformational  
 epitopes:

20

epitope F recognized by monoclonal antibodies 15C8C1, 12D11F1, and 8G10D1H9,  
 epitope G recognized by monoclonal antibody 9G3E6,  
 epitope H (or C) recognized by monoclonal antibodies 10D3C4 and 4H6B2,  
 epitope I recognized by monoclonal antibody 17F2C2.

25

The present invention provides an E1 and/or E2 specific monoclonal antibody raised upon immunization with a  
 composition as described herein. The antibodies of the present invention may be used, for example, as a medicament,  
 for incorporation into an immunoassay kit for detecting the presence of HCV E1 or E2 antigen, for prognosis/monitoring  
 of disease or for HCV therapy. The present invention provides for the use of an E1 and/or E2 specific monoclonal  
 antibody as described herein for the preparation of an immunoassay kit for detecting HCV E1 or E2 antigens, for the  
 preparation of a kit for prognosing/monitoring of HCV disease or for the preparation of a HCV medicament.

30

The present invention provides a method for in vitro diagnosis of HCV antigen present in a biological sample,  
 containing at least the following steps:

- (i) contacting said biological sample with an E1 and/or E2 specific monoclonal antibody as described herein,  
 preferably in an immobilized form under appropriate conditions which allow the formation of an immune complex,
- (ii) removing unbound components,
- (iii) incubating the immune complexes formed with heterologous antibodies, with the heterologous antibodies  
 being conjugated to a detectable label under appropriate conditions,
- (iv) detecting the presence of the immune complexes visually or mechanically.

35

The present invention provides a kit for determining the presence of HCV antigens present in a biological

sample, which includes at least the following: at least one E1 and/or E2 specific monoclonal antibody as described herein, preferably in an immobilized form on a solid substrate, a buffer or components necessary for producing the buffer enabling binding reaction between these antibodies and the HCV antigens present in a biological sample, and optionally a means for detecting the immune complexes formed in the preceding binding reaction.

5 The composition of the present invention may be provided in the form of a medicament.

The present invention provides a composition, as described herein for use as a vaccine for immunizing a mammal, preferably humans, against HCV, comprising administering an effective amount of said composition being optionally accompanied by pharmaceutically acceptable adjuvants, to produce an immune response.

10 The present invention provides a method of using the composition, as described herein, for the preparation of a vaccine for immunizing a mammal, preferably humans, against HCV, comprising administering an effective amount of said composition, optionally accompanied by pharmaceutically acceptable adjuvants, to produce an immune response.

The present invention provides a vaccine composition for immunizing a mammal, preferably humans, against HCV, which contains an effective amount of a composition containing an E1 and/or E2 containing composition as described herein, optionally also accompanied by pharmaceutically acceptable adjuvants.

15 The composition of the present invention may be provided in a form for *in vitro* detection of HCV antibodies present in a biological sample. The present invention also provides a method of preparing an immunoassay kit for detecting HCV antibodies present in a biological sample and a method of detecting HCV antibodies present in a biological sample using the kit of the invention to diagnose HCV antibodies present in a biological sample. Such a method of the present invention includes at least the following steps:

20 (i) contacting said biological sample with a composition as described herein, preferably in an immobilized form under appropriate conditions which allow the formation of an immune complex with HCV antibodies present in the biological sample,

(ii) removing unbound components,

25 (iii) incubating the immune complexes formed with heterologous antibodies, with the heterologous antibodies being conjugated to a detectable label under appropriate conditions,

(iv) detecting the presence of the immune complexes visually or mechanically.

30 The present invention provides a kit for determining the presence of HCV antibodies present in a biological sample, containing: at least one peptide or protein composition as described herein, preferably in an immobilized form on a solid substrate; a buffer or components necessary for producing the buffer enabling binding reaction between these proteins or peptides and the antibodies against HCV present in the biological sample; and, optionally, a means for detecting the immune complexes formed in the preceding binding reaction.

35 The present invention provides a method of *in vitro* monitoring HCV disease or diagnosing the response of a patient suffering from HCV infection to treatment, preferably with interferon, the method including: incubating a biological sample from the patient with HCV infection with an E1 protein or a suitable part thereof under conditions allowing the formation of an immunological complex; removing unbound components; calculating the anti-E1 titers present in the sample at the start of and during the course of treatment; monitoring the natural course of HCV disease, or diagnosing the response to treatment of the patient on the basis of the amount anti-E1 titers found in the sample at the start of

treatment and/or during the course of treatment.

The present invention provides a kit for monitoring HCV disease or prognosing the response to treatment, particularly with interferon, of patients suffering from HCV infection, wherein the kit contains: at least one E1 protein or E1 peptide, more particularly an E1 protein or E1 peptide as described herein; a buffer or components necessary for  
5 producing the buffer enabling the binding reaction between these proteins or peptides and the anti-E1 antibodies present in a biological sample; and optionally, means for detecting the immune complexes formed in the preceding binding reaction, optionally, also an automated scanning and interpretation device for inferring a decrease of anti-E1 titers during the progression of treatment.

The present invention provides a serotyping assay for detecting one or more serological types of HCV present  
10 in a biological sample, more particularly for detecting antibodies of the different types of HCV to be detected combined in one assay format, including at least the following steps : (i) contacting the biological sample to be analyzed for the presence of HCV antibodies of one or more serological types, with at least one of the E1 and/or E2 and/or E1/E2 protein compositions as described herein or at least one of the E1 or E2 peptide compositions described herein, preferentially in an immobilized form under appropriate conditions which allow the formation of an immune complex; (ii) removing  
15 unbound components; (iii) incubating the immune complexes formed with heterologous antibodies, with the heterologous antibodies being conjugated to a detectable label under appropriate conditions; and optionally, (iv) detecting the presence of said immune complexes visually or mechanically (e.g. by means of densitometry, fluorimetry, colorimetry) and inferring the presence of one or more HCV serological types present from the observed binding pattern.

The present invention provides a kit for serotyping one or more serological types of HCV present in a biological  
20 sample, more particularly for detecting the antibodies to these serological types of HCV containing: at least one E1 and/or E2 and/or E1/E2 protein as described herein or an E1 or E2 peptide as described herein; a buffer or components necessary for producing the buffer enabling the binding reaction between these proteins or peptides and the anti-E1 antibodies present in a biological sample;; optionally, means for detecting the immune complexes formed in the preceding binding reaction, optionally, also an automated scanning and interpretation device for detecting the presence  
25 of one or more serological types present from the observed binding pattern.

The present invention provides a peptide or protein composition as described herein, for immobilization on a solid substrate and incorporation into a reversed phase hybridization assay, preferably for immobilization as parallel lines onto a solid support such as a membrane strip, for determining the presence or the genotype of HCV according to a method as described herein.

The present invention provides a therapeutic vaccine composition containing a therapeutic effective amount of:  
30 a composition containing at least one purified recombinant HCV single or specific oligomeric recombinant envelope proteins selected from the group of an E1 protein and an E2 protein; and optionally a pharmaceutically acceptable adjuvant. The HCV envelope proteins of the vaccine of the present invention are optionally produced by recombinant mammalian cells or recombinant yeast cells. The invention provides a therapeutic vaccine composition  
35 containing a therapeutically effective amount of a composition containing at least one of the following E1 and E2 peptides:

E1-31 (SEQ ID NO:56) spanning amino acids 181 to 200 of the Core/E1 V1 region,

E1-33 (SEQ ID NO:57) spanning amino acids 193 to 212 of the E1 region,  
 E1-35 (SEQ ID NO:58) spanning amino acids 205 to 224 of the E1 V2 region (epitope B),  
 E1-35A (SEQ ID NO:59) spanning amino acids 208 to 227 of the E1 V2 region (epitope B),  
 1bE1 (SEQ ID NO:53) spanning amino acids 192 to 228 of E1 regions V1, C1, and V2 regions (containing  
 5 epitope B),

E1-51 (SEQ ID NO:66) spanning amino acids 301 to 320 of the E1 region,  
 E1-53 (SEQ ID NO:67) spanning amino acids 313 to 332 of the E1 C4 region (epitope A),  
 E1-55 (SEQ ID NO:68) spanning amino acids 325 to 344 of the E1 region,  
 Env 67 or E2-67 (SEQ ID NO:72) spanning amino acid positions 397 to 418 of the E2 region (epitope A),  
 10 Env 69 or E2-69 (SEQ ID NO:73) spanning amino acid positions 409 to 428 of the E2 region (epitope A),  
 Env 23 or E2-23 (SEQ ID NO:86) spanning positions 583 to 602 of the E2 region (epitope E),  
 Env 25 or E2-25 (SEQ ID NO:87) spanning positions 595 to 614 of the E2 region (epitope E),  
 Env 27 or E2-27 (SEQ ID NO:88) spanning positions 607 to 626 of the E2 region (epitope E),  
 Env 178 or E2-178 (SEQ ID NO:83) spanning positions 547 to 586 of the E2 region (epitope D), and  
 15 Env 13B or E2-13B (SEQ ID NO:82) spanning positions 523 to 542 of the E2 region (epitope C).

The present invention provides a method of treating a mammal, such as a human, infected with HCV comprising administering an effective amount of a composition as described herein, such as the above described vaccines, and optionally, a pharmaceutically acceptable adjuvant. In one embodiment, the composition of the invention is administered in combination with or at a time in conjunction with antiviral therapy, either soon prior to or subsequent to  
 20 or with administration of the composition of the invention.

The present invention provides a composition containing at least one purified recombinant HCV recombinant envelope proteins selected from the group of an E1 protein and an E2 protein, and optionally an adjuvant. In a preferred embodiment, the composition contains at least one of the following E1 and E2 peptides:

E1-31 (SEQ ID NO:56) spanning amino acids 181 to 200 of the Core/E1 V1 region,  
 25 E1-33 (SEQ ID NO:57) spanning amino acids 193 to 212 of the E1 region,  
 E1-35 (SEQ ID NO:58) spanning amino acids 205 to 224 of the E1 V2 region (epitope B),  
 E1-35A (SEQ ID NO:59) spanning amino acids 208 to 227 of the E1 V2 region (epitope B),  
 1bE1 (SEQ ID NO:53) spanning amino acids 192 to 228 of E1 regions V1, C1, and V2 regions (containing  
 epitope B),  
 30 E1-51 (SEQ ID NO:66) spanning amino acids 301 to 320 of the E1 region,  
 E1-53 (SEQ ID NO:67) spanning amino acids 313 to 332 of the E1 C4 region (epitope A),  
 E1-55 (SEQ ID NO:68) spanning amino acids 325 to 344 of the E1 region,  
 Env 67 or E2-67 (SEQ ID NO:72) spanning amino acid positions 397 to 418 of the E2 region (epitope A),  
 Env 69 or E2-69 (SEQ ID NO:73) spanning amino acid positions 409 to 428 of the E2 region (epitope A),  
 35 Env 23 or E2-23 (SEQ ID NO:86) spanning positions 583 to 602 of the E2 region (epitope E),  
 Env 25 or E2-25 (SEQ ID NO:87) spanning positions 595 to 614 of the E2 region (epitope E),  
 Env 27 or E2-27 (SEQ ID NO:88) spanning positions 607 to 626 of the E2 region (epitope E).

Env 178 or E2-178 (SEQ ID NO:83) spanning positions 547 to 586 of the E2 region (epitope D), and  
Env 13B or E2-13B (SEQ ID NO:82) spanning positions 523 to 542 of the E2 region (epitope C).

The present invention provides a therapeutic composition for inducing HCV-specific antibodies containing a  
therapeutic effective amount of a composition containing an E1/E2 complex formed from purified recombinant HCV  
5 single or specific oligomeric recombinant E1 or E2 proteins; and optionally a pharmaceutically acceptable adjuvant. The  
recombinant HCV envelope proteins of the invention may be produced by recombinant mammalian cells or recombinant  
HCV envelope proteins are produced by recombinant yeast cells. The present invention provides a method of treating a  
mammal, such as a human, infected with HCV including administering an effective amount of a composition as described  
herein and, optionally, a pharmaceutically acceptable adjuvant. The present invention provides a therapeutic  
10 composition for inducing HCV-specific antibodies containing a therapeutic effective amount of a composition containing  
at least one purified recombinant HCV single or specific oligomeric recombinant envelope protein selected from the  
group of an E1 protein and an E2 protein; and optionally a pharmaceutically acceptable adjuvant.



Figure and Table legends

- Figure 1 : Restriction map of plasmid pgpt ATA 18
- Figure 2 : Restriction map of plasmid pgs ATA 18
- 5 Figure 3 : Restriction map of plasmid pMS 66
- Figure 4 : Restriction map of plasmid pv HCV-11A
- Figure 5 : Anti-E1 levels in non-responders to IFN treatment
- Figure 6 : Anti-E1 levels in responders to IFN treatment
- Figure 7 : Anti-E1 levels in patients with complete response to IFN treatment
- 10 Figure 8 : Anti-E1 levels in incomplete responders to IFN treatment
- Figure 9 : Anti-E2 levels in non-responders to IFN treatment
- Figure 10 : Anti-E2 levels in responders to IFN treatment
- Figure 11 : Anti-E2 levels in incomplete responders to IFN treatment
- Figure 12 : Anti-E2 levels in complete responders to IFN treatment
- 15 Figure 13 : Human anti-E1 reactivity competed with peptides
- Figure 14 : Competition of reactivity of anti-E1 monoclonal antibodies with peptides
- Figure 15 : Anti-E1 (epitope 1) levels in non-responders to IFN treatment
- Figure 16 : Anti-E1 (epitope 1) levels in responders to IFN treatment
- Figure 17 : Anti-E1 (epitope 2) levels in non-responders to IFN treatment
- 20 Figure 18 : Anti-E1 (epitope 2) levels in responders to IFN treatment
- Figure 19 : Competition of reactivity of anti-E2 monoclonal antibodies with peptides
- Figure 20 : Human anti-E2 reactivity competed with peptides
- Figure 21 : Nucleic acid sequences of the present invention. The nucleic acid sequences encoding an E1 or E2 protein according to the present invention may be translated (SEQ ID NO 3 to 13, 21-31, 35 and 41-49 are translated in a reading frame starting from residue number 1, SEQ ID NO 37-39 are translated in a reading frame starting from residue number 2), into the amino acid sequences of the respective E1 or E2 proteins as shown in the sequence listing.
- 25
- Figure 22 : ELISA results obtained from lentil lectin chromatography eluate fractions of 4 different E1 purifications of cell lysates infected with vvHCV39 (type 1b), vvHCV40 (type 1b), vvHCV62 (type 3a), and vvHCV63 (type 5a).
- 30
- Figure 23 : Elution profiles obtained from the lentil lectin chromatography of the 4 different E1 constructs on the basis of the values as shown in Figure 22.
- Figure 24 : ELISA results obtained from fractions obtained after gel filtration chromatography of 4 different E1 purifications of cell lysates infected with vvHCV39 (type 1b), vvHCV40 (type 1b), vvHCV62

(type 3a), and vvHCV63 (type 5a).

- Figure 25: Profiles obtained from purifications of E1 proteins of type 1b (1), type 3a (2), and type 5a (3) (from RK13 cells infected with vvHCV39, vvHCV62, and vvHCV63, respectively; purified on lentil lectin and reduced as in example 5.2 - 5.3) and a standard (4). The peaks indicated with '1', '2', and '3', represent pure E1 protein peaks (see Figure 24, E1 reactivity mainly in fractions 26 to 30).
- Figure 26: Silver staining of an SDS-PAGE as described in example 4 of a raw lysate of E1 vvHCV40 (type 1b) (lane 1), pool 1 of the gelfiltration of vvHCV40 representing fractions 10 to 17 as shown in Figure 25 (lane 2), pool 2 of the gelfiltration of vvHCV40 representing fractions 18 to 25 as shown in Figure 25 (lane 3), and E1 pool (fractions 26 to 30) (lane 4).
- Figure 27: Streptavidine-alkaline phosphatase blot of the fractions of the gelfiltration of E1 constructs 39 (type 1b) and 62 (type 3a). The proteins were labelled with NEM-biotin. Lane 1: start gelfiltration construct 39, lane 2: fraction 26 construct 39, lane 3: fraction 27 construct 39, lane 4: fraction 28 construct 39, lane 5: fraction 29 construct 39, lane 6: fraction 30 construct 39, lane 7 fraction 31 construct 39, lane 8: molecular weight marker, lane 9: start gelfiltration construct 62, lane 10: fraction 26 construct 62, lane 11: fraction 27 construct 62, lane 12: fraction 28 construct 62, lane 13: fraction 29 construct 62, lane 14: fraction 30 construct 62, lane 15: fraction 31 construct 62.
- Figure 28: Silver staining of an SDS-PAGE gel of the gelfiltration fractions of vvHCV-39 (E1s, type 1b) and vvHCV-62 (E1s, type 3a) run under identical conditions as Figure 26. Lane 1: start gelfiltration construct 39, lane 2: fraction 26 construct 39, lane 3: fraction 27 construct 39, lane 4: fraction 28 construct 39, lane 5: fraction 29 construct 39, lane 6: fraction 30 construct 39, lane 7 fraction 31 construct 39, lane 8: molecular weight marker, lane 9: start gelfiltration construct 62, lane 10: fraction 26 construct 62, lane 11: fraction 27 construct 62, lane 12: fraction 28 construct 62, lane 13: fraction 29 construct 62, lane 14: fraction 30 construct 62, lane 15: fraction 31 construct 62.
- Figure 29: Western Blot analysis with anti-E1 mouse monoclonal antibody 5E1A10 giving a complete overview of the purification procedure. Lane 1: crude lysate, Lane 2: flow through of lentil chromatography, Lane 3: wash with Empigen BB after lentil chromatography, Lane 4: Eluate of lentil chromatography, Lane 5: Flow through during concentration of the lentil eluate, Lane 6: Pool of E1 after Size Exclusion Chromatography (gelfiltration).
- Figure 30: OD<sub>280</sub> profile (continuous line) of the lentil lectin chromatography of E2 protein from RK13 cells infected with vvHCV44. The dotted line represents the E2 reactivity as detected by ELISA (as in example 6).

- Figure 31A: OD<sub>280</sub> profile (continuous line) of the lentil-lectin gelfiltration chromatography E2 protein pool from RK13 cells infected with vvhCV44 in which the E2 pool is applied immediately on the gelfiltration column (non-reduced conditions). The dotted line represents the E2 reactivity as detected by ELISA (as in example 6).
- 5     Figure 31B: OD<sub>280</sub> profile (continuous line) of the lentil-lectin gelfiltration chromatography E2 protein pool from RK13 cells infected with vvhCV44 in which the E2 pool was reduced and blocked according to Example 5.3 (reduced conditions). The dotted line represents the E2 reactivity as detected by ELISA (as in example 6).
- 10     Figure 32: Ni<sup>2+</sup>-IMAC chromatography and ELISA reactivity of the E2 protein as expressed from vvhCV44 after gelfiltration under reducing conditions as shown in Figure 31B.
- Figure 33: Silver staining of an SDS-PAGE of 0.5 µg of purified E2 protein recovered by a 200 mM imidazole elution step (lane 2) and a 30mM imidazole wash (lane 1) of the Ni<sup>2+</sup>-IMAC chromatography as shown in Figure 32.
- 15     Figure 34: OD profiles of a desalting step of the purified E2 protein recovered by 200 mM imidazole as shown in Figure 33, intended to remove imidazole.
- Figure 35A: Antibody levels to the different HCV antigens (Core 1, Core 2, E2HCVR, NS3) for NR and LTR followed during treatment and over a period of 6 to 12 months after treatment determined by means of the LIAscan method. The average values are indicated by the curves with the open squares.
- 20     Figure 35B: Antibody levels to the different HCV antigens (NS4, NS5, E1 and E2) for NR and LTR followed during treatment and over a period of 6 to 12 months after treatment determined by means of the LIAscan method. The average values are indicated by the curve with the open squares.
- Figure 36: Average E1 antibody (E1Ab) and E2 antibody (E2Ab) levels in the LTR and NR groups.
- 25     Figure 37: Averages E1 antibody (E1Ab) levels for non-responders (NR) and long term responders (LTR) for type 1b and type 3a.
- Figure 38: Relative map positions of the anti-E2 monoclonal antibodies.
- Figure 39: Partial deglycosylation of HCV E1 envelope protein. The lysate of vvhCV10A-infected RK13 cells were incubated with different concentrations of glycosidases according to the manufacturer's instructions. Right panel: Glycopeptidase F (PNGase F). Left panel: Endoglycosidase H (Endo H).
- 30     Figure 40: Partial deglycosylation of HCV E2 envelope proteins. The lysate of vvhCV64-infected (E2) and vvhCV41-infected (E2s) RK13 cells were incubated with different concentrations of Glycopeptidase F (PNGase F) according to the manufacturer's instructions.

- Figure 41: In vitro mutagenesis of HCV E1 glycoproteins. Map of the mutated sequences and the creation of new restriction sites.
- Figure 42A: In vitro mutagenesis of HCV E1 glycoprotein (part 1). First step of PCR amplification.
- Figure 42B: In vitro mutagenesis of HCV E1 glycoprotein (part 2). Overlap extension and nested PCR.
- 5 Figure 43: In vitro mutagenesis of HCV E1 glycoproteins. Map of the PCR mutated fragments (GLY-# and OVR-#) synthesized during the first step of amplification.
- Figure 44A: Analysis of E1 glycoprotein mutants by Western blot expressed in HeLa (left) and RK13 (right) cells. Lane 1: wild type VV (vaccinia virus), Lane 2: original E1 protein (vHCV-10A), Lane 3: E1 mutant Gly-1 (vHCV-81), Lane 4: E1 mutant Gly-2 (vHCV-82), Lane 5: E1 mutant Gly-3 (vHCV-83), Lane 6: E1 mutant Gly-4 (vHCV-84), Lane 7: E1 mutant Gly-5 (vHCV-85), Lane 8: E1 mutant Gly-6 (vHCV-86).
- 10 Figure 44B: Analysis of E1 glycosylation mutant vaccinia viruses by PCR amplification/restriction. Lane 1: E1 (vHCV-10A), *BspE I*, Lane 2: E1.GLY-1 (vHCV-81), *BspE I*, Lane 4: E1 (vHCV-10A), *Sac I*, Lane 5: E1.GLY-2 (vHCV-82), *Sac I*, Lane 7: E1 (vHCV-10A), *Sac I*, Lane 8: E1.GLY-3 (vHCV-83), *Sac I*, Lane 10: E1 (vHCV-10A), *Stu I*, Lane 11: E1.GLY-4 (vHCV-84), *Stu I*, Lane 13: E1 (vHCV-10A), *Sma I*, Lane 14: E1.GLY-5 (vHCV-85), *Sma I*, Lane 16: E1 (vHCV-10A), *Stu I*, Lane 17: E1.GLY-6 (vHCV-86), *Stu I*, Lane 3 - 6 - 9 - 12 - 15 : Low Molecular Weight Marker, pBluescript SK+, *Msp I*.
- 15 Figure 45: SDS polyacrylamide gel electrophoresis of recombinant E2 expressed in *S. cerevisiae*. Inoculates were grown in leucine selective medium for 72 hrs. and diluted 1/15 in complete medium. After 10 days of culture at 28°C, medium samples were taken. The equivalent of 200 µl of culture supernatant concentrated by speedvac was loaded on the gel. Two independent transformants were analysed.
- 20 Figure 46: SDS polyacrylamide gel electrophoresis of recombinant E2 expressed in a glycosylation deficient *S. cerevisiae* mutant. Inoculae were grown in leucine selective medium for 72 hrs. and diluted 1/15 in complete medium. After 10 days of culture at 28°C, medium samples were taken. The equivalent of 350 µl of culture supernatant, concentrated by ion exchange chromatography, was loaded on the gel.
- 25 Figure 47: Profile of chimpanzees and immunization schedule.
- 30 Figure 48: Cellular response after 3 immunizations.
- Figure 49: Evolution of cellular response upon repeated E1 immunizations.
- Figure 50: Cellular response upon NS3 immunizations.
- Table 1 : Features of the respective clones and primers used for amplification for constructing the different forms of the E1 protein as despected in Example 1.

	Table 2 :	Summary of Anti-E1 tests
	Table 3 :	Synthetic peptides for competition studies
	Table 4:	Changes of envelope antibody levels over time.
	Table 5:	Difference between LTR and NR
5	Table 6:	Competition experiments between murine E2 monoclonal antibodies
	Table 7:	Primers for construction of E1 glycosylation mutants
	Table 8:	Analysis of E1 glycosylation mutants by ELISA
	Table 9:	Profile of adjuvanted E1 Balb/c mice.
	Table 10:	Humoral responses: No. of immunizations required for different E1-antibodies levels.

Example 1: Cloning and expression of the hepatitis C virus E1 protein1. Construction of vaccinia virus recombination vectors

5 The pgpATA18 vaccinia recombination plasmid is a modified version of pATA18 (Stunnenberg et al, 1988) with an additional insertion containing the *E. coli* xanthine guanine phosphoribosyl transferase gene under the control of the vaccinia virus I3 intermediate promoter (Figure 1). The plasmid pgsATA18 was constructed by inserting an oligonucleotide linker with SEQ ID NO 1/94, containing stop codons in the three reading frames, into the Pst I and HindIII-cut pATA18 vector. This created an extra Pac I restriction site (Figure 2). The original HindIII

10 site was not restored.

Oligonucleotide linker with SEQ ID NO 1/94:

5' G GCATGC AAGCTT AATTAAAT 3'  
 3' ACGTC CGTACG TTCGAA TTAATTAA TCGA 5'

15 PstI SphI HindIII Pac I (HindIII)

In order to facilitate rapid and efficient purification by means of  $\text{Ni}^{2+}$  chelation of engineered histidine stretches fused to the recombinant proteins, the vaccinia recombination vector pMS66 was designed to express secreted proteins with an additional carboxy-terminal histidine tag. An oligonucleotide linker, with SEQ ID NO 2/95, containing unique sites for 3 restriction enzymes generating blunt ends (Sma I, Stu I and Pml I/Bbr PI) was synthesized in such a way that the carboxy-terminal end of any cDNA could be inserted in frame with a sequence encoding the protease factor Xa cleavage site followed by a nucleotide sequence encoding 6 histidines and 2 stop codons (a new Pac I restriction site was also created downstream the 3' end). This

20 oligonucleotide with SEQ ID NO 2/95 was introduced between the Xma I and Pst I sites of pgpATA18 (Figure 3).

Oligonucleotide linker with SEQ ID NO 2/95:

5' CCGGG GAGGCCGTGCACGTGATCGAGGGCAGACACCATCACCACCATCACTAATAGTTAATTAA CTGCA  
 3'

30 3' C CTCCGGACGTGCACTAGCTCCCGTCTGTGGTAGTGGTGGTAGTGATTATCAATTAATT G

XmaI PstI

Plasmid pgpATA-18 contained within *Escherichia coli* MC1061(lambda) has been deposited under the terms of the Budapest Treaty at BCCM/LMBP (Belgian Coordinated Collections of microorganisms/Laboratorium voor Moleculaire Biologie - Plasmidencollectie, Universiteit Gent, K.L. Ledeganckstraat 35, B-9000 Ghent, Belgium), and bears accession number LMBP4486. Said deposit was made on January 9, 2002.

35

Example 2. Construction of HCV recombinant plasmids

### 2.1. Constructs encoding different forms of the E1 protein

Polymerase Chain Reaction (PCR) products were derived from the serum samples by RNA preparation and subsequent reverse-transcription and PCR as described previously (Stuyver et al., 1993b). Table 1 shows the features of the respective clones and the primers used for amplification. The PCR fragments were cloned into the Sma I-cut pSP72 (Promega) plasmids. The following clones were selected for insertion into vaccinia recombination vectors: HCCI9A (SEQ ID NO 3), HCCI10A (SEQ ID NO 5), HCCI11A (SEQ ID NO 7), HCCI12A (SEQ ID NO 9), HCCI13A (SEQ ID NO 11), and HCCI17A (SEQ ID NO 13) as depicted in Figure 21. cDNA fragments containing the E1-coding regions were cleaved by EcoRI and HindIII restriction from the respective pSP72 plasmids and inserted into the EcoRI/HindIII-cut pgptATA-18 vaccinia recombination vector (described in example 1), downstream of the 11K vaccinia virus late promoter. The respective plasmids were designated pvHCV-9A, pvHCV-10A, pvHCV-11A, pvHCV-12A, pvHCV-13A and pvHCV-17A, of which pvHCV-11A is shown in Figure 4.

### 2.2. Hydrophobic region E1 deletion mutants

Clone HCCI37, containing a deletion of codons Asp264 to Val287 (nucleotides 790 to 861, region encoding hydrophobic domain I) was generated as follows: 2 PCR fragments were generated from clone HCCI10A with primer sets HCP52 (SEQ ID NO 16)/HCP107 (SEQ ID NO 19) and HCP108 (SEQ ID NO 20)/HCP54 (SEQ ID NO 18). These primers are shown in Figure 21. The two PCR fragments were purified from agarose gel after electrophoresis and 1 ng of each fragment was used together as template for PCR by means of primers HCP52 (SEQ ID NO 16) and HCP54 (SEQ ID NO 18). The resulting fragment was cloned into the Sma I-cut pSP72 vector and clones containing the deletion were readily identified because of the deletion of 24 codons (72 base pairs). Plasmid pSP72HCCI37 containing clone HCCI37 (SEQ ID NO 15) was selected. A recombinant vaccinia plasmid containing the full-length E1 cDNA lacking hydrophobic domain I was constructed by inserting the HCV sequence surrounding the deletion (fragment cleaved by Xma I and BamH I from the vector pSP72-HCCI37) into the Xma I-Bam H I sites of the vaccinia plasmid pvHCV-10A. The resulting plasmid was named pvHCV-37. After confirmatory sequencing, the amino-terminal region containing the internal deletion was isolated from this vector pvHCV-37 (cleavage by EcoR I and BstE II) and reinserted into the Eco RI and Bst EII-cut pvHCV-11A plasmid. This construct was expected to express an E1 protein with both hydrophobic domains deleted and was named pvHCV-38. The E1-coding region of clone HCCI38 is represented by SEQ ID NO 23.

As the hydrophilic region at the E1 carboxyterminus (theoretically extending to around amino acids 337-340) was not completely included in construct pvHCV-38, a larger E1 region lacking hydrophobic domain I

was isolated from the pvHCV-37 plasmid by EcoR I/Bam HI cleavage and cloned into an EcoRI/BamHI-cut pgsATA-18 vector. The resulting plasmid was named pvHCV-39 and contained clone HCCI39 (SEQ ID NO 25). The same fragment was cleaved from the pvHCV-37 vector by BamH I (of which the sticky ends were filled with Klenow DNA Polymerase I (Boehringer)) and subsequently by EcoR I (5' cohesive end). This sequence was inserted into the EcoRI and Bbr PI-cut vector pMS-66. This resulted in clone HCCI40 (SEQ ID NO 27) in plasmid pvHCV-40, containing a 6 histidine tail at its carboxy-terminal end.

### 2.3. E1 of other genotypes

Clone HCCI62 (SEQ ID NO 29) was derived from a type 3a-infected patient with chronic hepatitis C (serum BR36, clone BR36-9-13, SEQ ID NO 19 in WO 94/25601, and see also Stuyver et al. 1993a) and HCCI63 (SEQ ID NO 31) was derived from a type 5a-infected child with post-transfusion hepatitis (serum BE95, clone PC-4-1, SEQ ID NO 45 in WO 94/25601).

### 2.4. E2 constructs

The HCV E2 PCR fragment 22 was obtained from serum BE11 (genotype 1b) by means of primers HCPri09 (SEQ ID NO 33) and HCPri72 (SEQ ID NO 34) using techniques of RNA preparation, reverse-transcription and PCR, as described in Stuyver et al., 1993b, and the fragment was cloned into the Sma I-cut pSP72 vector. Clone HCCI22A (SEQ ID NO 35) was cut with NcoI/AlwNI or by BamHI/AlwNI and the sticky ends of the fragments were blunted (NcoI and BamHI sites with Klenow DNA Polymerase I (Boehringer), and AlwNI with T4 DNA polymerase (Boehringer)). The BamHI/AlwNI cDNA fragment was then inserted into the vaccinia pgsATA-18 vector that had been linearized by EcoR I and Hind III cleavage and of which the cohesive ends had been filled with Klenow DNA Polymerase (Boehringer). The resulting plasmid was named pvHCV-41 and encoded the E2 region from amino acids Met347 to Gln673, including 37 amino acids (from Met347 to Gly383) of the E1 protein that can serve as signal sequence. The same HCV cDNA was inserted into the EcoR I and Bbr PI-cut vector pMS66, that had subsequently been blunt ended with Klenow DNA Polymerase. The resulting plasmid was named pvHCV-42 and also encoded amino acids 347 to 683. The NcoI/AlwNI fragment was inserted in a similar way into the same sites of pgsATA-18 (pvHCV-43) or pMS-66 vaccinia vectors (pvHCV-44). pvHCV-43 and pvHCV-44 encoded amino acids 364 to 673 of the HCV polyprotein, of which amino acids 364 to 383 were derived from the natural carboxyterminal region of the E1 protein encoding the signal sequence for E2, and amino acids 384 to 673 of the mature E2 protein.

### 2.5. Generation of recombinant HCV-vaccinia viruses



Rabbit kidney RK13 cells (ATCC CCL 37), human osteosarcoma 143B thymidine kinase deficient (TK<sup>-</sup>) (ATCC CRL 8303), HeLa (ATCC CCL 2), and Hep G2 (ATCC HB 8065) cell lines were obtained from the American Type Culture Collection (ATCC, Rockville, Md, USA). The cells were grown in Dulbecco's modified Eagle medium (DMEM) supplemented with 10 % foetal calf serum, and with Earle's salts (EMEM) for RK13 and 143 B (TK<sup>-</sup>), and with glucose (4 g/l) for Hep G2. The vaccinia virus WR strain (Western Reserve, ATCC VR119) was routinely propagated in either 143B or RK13 cells, as described previously (Panicali & Paoletti, 1982; Piccini et al., 1987; Mackett et al., 1982, 1984, and 1986). A confluent monolayer of 143B cells was infected with wild type vaccinia virus at a multiplicity of infection (m.o.i.) of 0.1 (= 0.1 plaque forming unit (PFU) per cell). Two hours later, the vaccinia recombination plasmid was transfected into the infected cells in the form of a calcium phosphate coprecipitate containing 500 ng of the plasmid DNA to allow homologous recombination (Graham & van der Eb, 1973; Mackett et al., 1985). Recombinant viruses expressing the *Escherichia coli* xanthine-guanine phosphoribosyl transferase (gpt) protein were selected on rabbit kidney RK13 cells incubated in selection medium (EMEM containing 25 µg/ml mycophenolic acid (MPA), 250 µg/ml xanthine, and 15 µg/ml hypoxanthine; Falkner and Moss, 1988; Janknecht et al, 1991). Single recombinant viruses were purified on fresh monolayers of RK13 cells under a 0.9% agarose overlay in selection medium. Thymidine kinase deficient (TK<sup>-</sup>) recombinant viruses were selected and then plaque purified on fresh monolayers of human 143B cells (TK<sup>-</sup>) in the presence of 25 µg/ml 5-bromo-2'-deoxyuridine. Stocks of purified recombinant HCV-vaccinia viruses were prepared by infecting either human 143 B or rabbit RK13 cells at an m.o.i. of 0.05 (Mackett et al, 1988). The insertion of the HCV cDNA fragment in the recombinant vaccinia viruses was confirmed on an aliquot (50 µl) of the cell lysate after the MPA selection by means of PCR with the primers used to clone the respective HCV fragments (see Table 1). The recombinant vaccinia-HCV viruses were named according to the vaccinia recombination plasmid number, e.g. the recombinant vaccinia virus vvHCV-10A was derived from recombining the wild type WR strain with the pvHCV-10A plasmid.

25

### Example 3: infection of cells with recombinant vaccinia viruses

A confluent monolayer of RK13 cells was infected at a m.o.i. of 3 with the recombinant HCV-vaccinia viruses as described in example 2. For infection, the cell monolayer was washed twice with phosphate-buffered saline pH 7.4 (PBS) and the recombinant vaccinia virus stock was diluted in MEM medium. Two hundred µl of the virus solution was added per 10<sup>6</sup> cells such that the m.o.i. was 3, and incubated for 45 min at 24°C. The virus solution was aspirated and 2 ml of complete growth medium (see example 2) was added per 10<sup>6</sup> cells. The cells were incubated for 24 hr at 37°C during which expression of the HCV proteins took place.

30

Example 4: Analysis of recombinant proteins by means of western blotting

The infected cells were washed two times with PBS, directly lysed with lysis buffer (50 mM Tris.HCl pH 7.5, 150 mM NaCl, 1% Triton X-100, 5 mM MgCl<sub>2</sub>, 1 µg/ml aprotinin (Sigma, Bornem, Belgium)) or detached  
5 from the flasks by incubation in 50 mM Tris.HCl pH 7.5/ 10 mM EDTA/ 150 mM NaCl for 5 min, and collected by centrifugation (5 min at 1000g). The cell pellet was then resuspended in 200 µl lysis buffer (50 mM Tris.HCl pH 8.0, 2 mM EDTA, 150 mM NaCl, 5 mM MgCl<sub>2</sub>, aprotinin, 1% Triton X-100) per 10<sup>6</sup> cells. The cell lysates were cleared for 5 min at 14,000 rpm in an Eppendorf centrifuge to remove the insoluble debris. Proteins of 20 µl lysate were separated by means of sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-  
10 PAGE). The proteins were then electro-transferred from the gel to a nitrocellulose sheet (Amersham) using a Hoefer HSI transfer unit cooled to 4°C for 2 hr at 100 V constant voltage, in transfer buffer (25 mM Tris.HCl pH 8.0, 192 mM glycine, 20% (v/v) methanol). Nitrocellulose filters were blocked with Blotto (5 % (w/v) fat-free instant milk powder in PBS; Johnson et al., 1981) and incubated with primary antibodies diluted in Blotto/0.1 % Tween 20. Usually, a human negative control serum or serum of a patient infected with HCV were 200 times  
15 diluted and preincubated for 1 hour at room temperature with 200 times diluted wild type vaccinia virus-infected cell lysate in order to decrease the non-specific binding. After washing with Blotto/0.1% Tween 20, the nitrocellulose filters were incubated with alkaline phosphatase substrate solution diluted in Blotto/0.1 % Tween 20. After washing with 0.1% Tween 20 in PBS, the filters were incubated with alkaline phosphatase substrate solution (100 mM Tris.HCl pH 9.5, 100 mM NaCl, 5 mM MgCl<sub>2</sub>, 0.38 µg/ml nitroblue tetrazolium, 0.165 µg/ml 5-  
20 bromo-4-chloro-3-indolylphosphate). All steps, except the electrotransfer, were performed at room temperature.

Example 5: Purification of recombinant E1 or E2 protein25 5.1. Lysis

Infected RK13 cells (carrying E1 or E2 constructs) were washed 2 times with phosphate-buffered saline (PBS) and detached from the culture recipients by incubation in PBS containing 10 mM EDTA. The detached cells were washed twice with PBS and 1 ml of lysis buffer (50 mM Tris.HCl pH 7.5, 150 mM NaCl, 1% Triton X-  
30 100, 5 mM MgCl<sub>2</sub>, 1 µg/ml aprotinin (Sigma, Bornem, Belgium) containing 2 mM biotinylated N-ethylmaleimide (biotin-NEM) (Sigma) was added per 10<sup>5</sup> cells at 4°C. This lysate was homogenized with a type B douncer and left at room temperature for 0.5 hours. Another 5 volumes of lysis buffer containing 10 mM N-ethylmaleimide (NEM, Aldrich, Bornem, Belgium) was added to the primary lysate and the mixture was left at room temperature for 15 min. Insoluble cell debris was cleared from the solution by centrifugation in a Beckman JA-14 rotor at

14,000 rpm (30100 g at  $r_{max}$ ) for 1 hour at 4°C.

## 5.2. Lectin Chromatography

5

The cleared cell lysate was loaded at a rate of 1ml/min on a 0.8 by 10 cm Lentil-lectin Sepharose 4B column (Pharmacia) that had been equilibrated with 5 column volumes of lysis buffer at a rate of 1ml/min. The lentil-lectin column was washed with 5 to 10 column volumes of buffer 1 (0.1M potassium phosphate pH 7.3, 500 mM KCl, 5% glycerol, 1 mM 6-NH<sub>2</sub>-hexanoic acid, 1 mM MgCl<sub>2</sub>, and 1% DecylPEG (KWANT, Bedum, The Netherlands). In some experiments, the column was subsequently washed with 10 column volumes of buffer 1 containing 0.5% Empigen-BB (Calbiochem, San Diego, CA, USA) instead of 1% DecylPEG. The bound material was eluted by applying elution buffer (10 mM potassium phosphate pH 7.3, 5% glycerol, 1 mM hexanoic acid, 1mM MgCl<sub>2</sub>, 0.5% Empigen-BB, and 0.5 M  $\alpha$ -methyl-mannopyranoside). The eluted material was fractionated and fractions were screened for the presence of E1 or E2 protein by means of ELISA as described in example 6.

10

15 Figure 22 shows ELISA results obtained from lentil lectin eluate fractions of 4 different E1 purifications of cell lysates infected with wHCV39 (type 1b), wHCV40 (type 1b), wHCV62 (type 3a), and wHCV63 (type 5a). Figure 23 shows the profiles obtained from the values shown in Figure 22. These results show that the lectin affinity column can be employed for envelope proteins of the different types of HCV.

20

## 5.3. Concentration and partial reduction

The E1- or E2-positive fractions were pooled and concentrated on a Centricon 30 kDa (Amicon) by centrifugation for 3 hours at 5,000 rpm in a Beckman JA-20 rotor at 4°C. In some experiments the E1- or E2-positive fractions were pooled and concentrated by nitrogen evaporation. An equivalent of  $3.10^8$  cells was concentrated to approximately 200  $\mu$ l. For partial reduction, 30% Empigen-BB (Calbiochem, San Diego, CA, USA) was added to this 200  $\mu$ l to a final concentration of 3.5 %, and 1M DTT in H<sub>2</sub>O was subsequently added to a final concentration of 1.5 to 7.5 mM and incubated for 30 min at 37 °C. NEM (1M in dimethylsulphoxide) was subsequently added to a final concentration of 50 mM and left to react for another 30 min at 37°C to block the free sulphydryl groups.

25

30

## 5.4. Gel filtration chromatography

A Superdex-200 HR 10/20 column (Pharmacia) was equilibrated with 3 column volumes PBS/3%

Empigen-BB. The reduced mixture was injected in a 500  $\mu$ l sample loop of the Smart System (Pharmacia) and PBS/3% Empigen-BB buffer was added for gelfiltration. Fractions of 250  $\mu$ l were collected from  $V_0$  to  $V_6$ . The fractions were screened for the presence of E1 or E2 protein as described in example 6.

Figure 24 shows ELISA results obtained from fractions obtained after gelfiltration chromatography of 4 different E1 purifications of cell lysates infected with vvHCV39 (type 1b), vvHCV40 (type 1b), vvHCV62 (type 3a), and vvHCV63 (type 5a). Figure 25 shows the profiles obtained from purifications of E1 proteins of types 1b, 3a, and 5a (from RK13 cells infected with vvHCV39, vvHCV62, and vvHCV63, respectively; purified on lentil lectin and reduced as in the previous examples). The peaks indicated with '1', '2', and '3', represent pure E1 protein peaks (E1 reactivity mainly in fractions 26 to 30). These peaks show very similar molecular weights of approximately 70 kDa, corresponding to dimeric E1 protein. Other peaks in the three profiles represent vaccinia virus and/or cellular proteins which could be separated from E1 only because of the reduction step as outlined in example 5.3. and because of the subsequent gelfiltration step in the presence of the proper detergent. As shown in Figure 26 pool 1 (representing fractions 10 to 17) and pool 2 (representing fractions 18 to 25) contain contaminating proteins not present in the E1 pool (fractions 26 to 30). The E1 peak fractions were ran on SDS/PAGE and blotted as described in example 4. Proteins labelled with NEM-biotin were detected by streptavidin-alkaline phosphatase as shown in Figure 27. It can be readily observed that, amongst others, the 29 kDa and 45kDa contaminating proteins present before the gelfiltration chromatography (lane 1) are only present at very low levels in the fractions 26 to 30. The band at approximately 65kDa represents the E1 dimeric form that could not be entirely disrupted into the monomeric E1 form. Similar results were obtained for the type 3a E1 protein (lanes 10 to 15), which shows a faster mobility on SDS/PAGE because of the presence of only 5 carbohydrates instead of 6. Figure 28 shows a silver stain of an SDS/PAGE gel run in identical conditions as in Figure 26. A complete overview of the purification procedure is given in Figure 29.

The presence of purified E1 protein was further confirmed by means of western blotting as described in example 4. The dimeric E1 protein appeared to be non-aggregated and free of contaminants. The subtype 1b E1 protein purified from vvHCV40-infected cells according to the above scheme was aminoterminally sequenced on an 477 Perkins-Elmer sequencer and appeared to contain a tyrosine as first residue. This confirmed that the E1 protein had been cleaved by the signal peptidase at the correct position (between A191 and Y192) from its signal sequence. This confirms the finding of Hijikata et al. (1991) that the aminotermminus of the mature E1 protein starts at amino acid position 192.

#### 5.5. Purification of the E2 protein

The E2 protein (amino acids 384 to 673) was purified from RK13 cells infected with vvHCV44 as indicated in Examples 5.1 to 5.4. Figure 30 shows the OD<sub>230</sub> profile (continuous line) of the lentil lectin

chromatography. The dotted line represents the E2 reactivity as detected by ELISA (see example 6). Figure 31 shows the same profiles obtained from gelfiltration chromatography of the lentil-lectin E2 pool (see Figure 30), part of which was reduced and blocked according to the methods as set out in example 5.3., and part of which was immediately applied to the column. Both parts of the E2 pool were run on separate gelfiltration columns. It could be demonstrated that E2 forms covalently-linked aggregates with contaminating proteins if no reduction has been performed. After reduction and blocking, the majority of contaminating proteins segregated into the  $V_0$  fraction. Other contaminating proteins copurified with the E2 protein, were not covalently linked to the E2 protein any more because these contaminants could be removed in a subsequent step. Figure 32 shows an additional  $\text{Ni}^{2+}$ -IMAC purification step carried out for the E2 protein purification. This affinity purification step employs the 6 histidine residues added to the E2 protein as expressed from vHCV44. Contaminating proteins either run through the column or can be removed by a 30 mM imidazole wash. Figure 33 shows a silver-stained SDS/PAGE of 0.5  $\mu\text{g}$  of purified E2 protein and a 30 mM imidazole wash. The pure E2 protein could be easily recovered by a 200 mM imidazole elution step. Figure 34 shows an additional desalting step intended to remove imidazole and to be able to switch to the desired buffer, e.g. PBS, carbonate buffer, saline.

Starting from about 50,000  $\text{cm}^2$  of RK13 cells infected with vHCV11A (or vHCV40) for the production of E1 or vHCV41, vHCV42, vHCV43, or vHCV44 for production of E2 protein, the procedures described in examples 5.1 to 5.3 allow the purification of approximately 1.3 mg of E1 protein and 0.6 mg of E2 protein.

It should also be remarked that secreted E2 protein (constituting approximately 30-40%, 60-70% being in the intracellular form) is characterized by aggregate formation (contrary to expectations). The same problem is thus posed to purify secreted E2. The secreted E2 can be purified as disclosed above.

#### Example 6: ELISA for the detection of anti-E1 or anti-E2 antibodies or for the detection of E1 or E2 proteins

Maxisorb microwell plates (Nunc, Roskilde, Denmark) were coated with 1 volume (e.g. 50  $\mu\text{l}$  or 100  $\mu\text{l}$  or 200  $\mu\text{l}$ ) per well of a 5  $\mu\text{g}/\text{ml}$  solution of Streptavidin (Boehringer Mannheim) in PBS for 16 hours at 4°C or for 1 hour at 37°C. Alternatively, the wells were coated with 1 volume of 5  $\mu\text{g}/\text{ml}$  of Galanthus nivalis agglutinin (GNA) in 50 mM sodium carbonate buffer pH 9.6 for 16 hours at 4°C or for 1 hour at 37°C. In the case of coating with GNA, the plates were washed 2 times with 400  $\mu\text{l}$  of Washing Solution of the Innatest HCV Ab III kit (Innogenetics, Zwijndrecht, Belgium). Unbound coating surfaces were blocked with 1.5 to 2 volumes of blocking solution (0.1% casein and 0.1%  $\text{NaN}_3$  in PBS) for 1 hour at 37°C or for 16 hours at 4°C. Blocking solution was aspirated. Purified E1 or E2 was diluted to 100-1000 ng/ml (concentration measured at  $A = 280 \text{ nm}$ ) or column fractions to be screened for E1 or E2 (see example 5), or E1 or E2 in non-purified cell lysates (example 5.1.)

were diluted 20 times in blocking solution, and 1 volume of the E1 or E2 solution was added to each well and incubated for 1 hour at 37°C on the Streptavidin- or GNA-coated plates. The microwells were washed 3 times with 1 volume of Washing Solution of the Innatest HCV Ab III kit (Innogenetics, Zwijndrecht, Belgium). Serum samples were diluted 20 times or monoclonal anti-E1 or anti-E2 antibodies were diluted to a concentration of 20 ng/ml in Sample Diluent of the Innatest HCV Ab III kit and 1 volume of the solution was left to react with the E1 or E2 protein for 1 hour at 37°C. The microwells were washed 5 times with 400 µl of Washing Solution of the Innatest HCV Ab III kit (Innogenetics, Zwijndrecht, Belgium). The bound antibodies were detected by incubating each well for 1 hour at 37°C with a goat anti-human or anti-mouse IgG, peroxidase-conjugated secondary antibody (DAKO, Glostrup, Denmark) diluted 1/80,000 in 1 volume of Conjugate Diluent of the Innatest HCV Ab III kit (Innogenetics, Zwijndrecht, Belgium), and color development was obtained by addition of substrate of the Innatest HCV Ab III kit (Innogenetics, Zwijndrecht, Belgium) diluted 100 times in 1 volume of Substrate Solution of the Innatest HCV Ab III kit (Innogenetics, Zwijndrecht, Belgium) for 30 min at 24°C after washing of the plates 3 times with 400 µl of Washing Solution of the Innatest HCV Ab III kit (Innogenetics, Zwijndrecht, Belgium).

15

#### Example 7: Follow up of patient groups with different clinical profiles

##### 7.1. Monitoring of anti-E1 and anti-E2 antibodies

The current hepatitis C virus (HCV) diagnostic assays have been developed for screening and confirmation of the presence of HCV antibodies. Such assays do not seem to provide information useful for monitoring of treatment or for prognosis of the outcome of disease. However, as is the case for hepatitis B, detection and quantification of anti-envelope antibodies may prove more useful in a clinical setting. To investigate the possibility of the use of anti-E1 antibody titer and anti-E2 antibody titer as prognostic markers for outcome of hepatitis C disease, a series of IFN-α treated patients with long-term sustained response (defined as patients with normal transaminase levels and negative HCV-RNA test (PCR in the 5' non-coding region) in the blood for a period of at least 1 year after treatment) was compared with patients showing no response or showing biochemical response with relapse at the end of treatment.

A group of 8 IFN-α treated patients with long-term sustained response (LTR, follow up 1 to 3.5 years, 3 type 3a and 5 type 1b) was compared with 9 patients showing non-complete responses to treatment (NR, follow up 1 to 4 years, 6 type 1b and 3 type 3a). Type 1b (vHCV-39, see example 2.5.) and 3a E1 (vHCV-62, see example 2.5.) proteins were expressed by the vaccinia virus system (see examples 3 and 4) and purified to homogeneity (example 5). The samples derived from patients infected with a type 1b hepatitis C virus were tested for reactivity with purified type 1b E1 protein, while samples of a type 3a infection were tested for reactivity

of anti-type 3a E1 antibodies in an ELISA as described in example 6. The genotypes of hepatitis C viruses infecting the different patients were determined by means of the Inno-LiPA genotyping assay (Innogenetics, Zwijndrecht, Belgium). Figure 5 shows the anti-E1 signal-to-noise ratios of these patients followed during the course of interferon treatment and during the follow-up period after treatment. LTR cases consistently showed rapidly declining anti-E1 levels (with complete negatvation in 3 cases), while anti-E1 levels of NR cases remained approximately constant. Some of the obtained anti-E1 data are shown in Table 2 as average S/N ratios  $\pm$  SD (mean anti-E1 titer). The anti-E1 titer could be deduced from the signal to noise ratio as show in Figures 5, 6, 7, and 8.

Already at the end of treatment, marked differences could be observed between the 2 groups. Anti-E1 antibody titers had decreased 6.9 times in LTR but only 1.5 times in NR. At the end of follow up, the anti-E1 titers had declined by a factor of 22.5 in the patients with sustained response and even slightly increased in NR. Therefore, based on these data, decrease of anti-E1 antibody levels during monitoring of IFN- $\alpha$  therapy correlates with long-term, sustained response to treatment. The anti-E1 assay may be very useful for prognosis of long-term response to IFN treatment, or to treatment of the hepatitis C disease in general.

This finding was not expected. On the contrary, the inventors had expected the anti-E1 antibody levels to increase during the course of IFN treatment in patients with long term response. As is the case for hepatitis B, the virus is cleared as a consequence of the seroconversion for anti-HBsAg antibodies. Also in many other virus infections, the virus is eliminated when anti-envelope antibodies are raised. However, in the experiments of the present invention, anti-E1 antibodies clearly decreased in patients with a long-term response to treatment, while the antibody-level remained approximately at the same level in non-responding patients. Although the outcome of these experiments was not expected, this non-obvious finding may be very important and useful for clinical diagnosis of HCV infections. As shown in Figures 9, 10, 11, and 12, anti-E2 levels behaved very differently in the same patients studied and no obvious decline in titers was observed as for anti-E1 antibodies. Figure 35 gives a complete overview of the pilot study.

As can be deduced from Table 2, the anti-E1 titers were on average at least 2 times higher at the start of treatment in long term responders compared with incomplete responders to treatment. Therefore, measuring the titer of anti-E1 antibodies at the start of treatment, or monitoring the patient during the course of infection and measuring the anti-E1 titer, may become a useful marker for clinical diagnosis of hepatitis C. Furthermore, the use of more defined regions of the E1 or E2 proteins may become desirable, as shown in example 7.3.

#### 7.2. Analysis of E1 and E2 antibodies in a larger patient cohort

The pilot study lead the inventors to conclude that, in case infection was completely cleared, antibodies to the HCV envelope proteins changed more rapidly than antibodies to the more conventionally studied HCV

antigens, with E1 antibodies changing most vigorously. We therefore included more type 1b and 3a-infected LTR and further supplemented the cohort with a matched series of NR, such that both groups included 14 patients each. Some partial responders (PR) and responders with relapse (RR) were also analyzed.

Figure 36 depicts average E1 antibody (E1Ab) and E2 antibody (E2Ab) levels in the LTR and NR groups and Tables 4 and 5 show the statistical analyses. In this larger cohort, higher E1 antibody levels before IFN- $\alpha$  therapy were associated with LTR ( $P < 0.03$ ). Since much higher E1 antibody levels were observed in type 3a-infected patients compared with type 1b-infected patients (Figure 37), the genotype was taken into account (Table 4). Within the type 1b-infected group, LTR also had higher E1 antibody levels than NR at the initiation of treatment ( $P < 0.05$ ); the limited number of type 3a-infected NR did not allow statistical analysis.

Of antibody levels monitored in LTR during the 1.5-year follow up period, only E1 antibodies cleared rapidly compared with levels measured at initiation of treatment [ $P = 0.0058$ , end of therapy;  $P = 0.0047$  and  $P = 0.0051$  at 6 and 12 months after therapy, respectively]. This clearance remained significant within type 1- or type 3-infected LTR (average  $P$  values  $< 0.05$ ). These data confirmed the initial finding that E1Ab levels decrease rapidly in the early phase of resolution. This feature seems to be independent of viral genotype. In NR, PR, or RR, no changes in any of the antibodies measured were observed throughout the follow up period. In patients who responded favourably to treatment with normalization of ALT levels and HCV-RNA negative during treatment, there was a marked difference between sustained responders (LTR) and responders with a relapse (RR). In contrast to LTR, RR did not show any decreasing E1 antibody levels, indicating the presence of occult HCV infection that could neither be demonstrated by PCR or other classical techniques for detection of HCV-RNA, nor by raised ALT levels. The minute quantities of viral RNA, still present in the RR group during treatment, seemed to be capable of anti-E1 B cell stimulation. Anti-E1 monitoring may therefore not only be able to discriminate LTR from NR, but also from RR.

### 7.3. Monitoring of antibodies of defined regions of the E1 protein

Although the molecular biological approach of identifying HCV antigens resulted in unprecedented breakthrough in the development of viral diagnostics, the method of immune screening of  $\lambda$ gt11 libraries predominantly yielded linear epitopes dispersed throughout the core and non-structural regions, and analysis of the envelope regions had to await cloning and expression of the E1/E2 region in mammalian cells. This approach sharply contrasts with many other viral infections of which epitopes to the envelope regions had already been mapped long before the deciphering of the genomic structure. Such epitopes and corresponding antibodies often had neutralizing activity useful for vaccine development and/or allowed the development of diagnostic assays with clinical or prognostic significance (e.g. antibodies to hepatitis B surface antigen).



As no HCV vaccines or tests allowing clinical diagnosis and prognosis of hepatitis C disease are available today, the characterization of viral envelope regions exposed to immune surveillance may significantly contribute to new directions in HCV diagnosis and prophylaxis.

Several 20-mer peptides (Table 3) that overlapped each other by 8 amino acids, were synthesized according to a previously described method (EP-A-0 489 968) based on the HC-J1 sequence (Okamoto et al., 1990). None of these, except peptide env35 (also referred to as E1-35), was able to detect antibodies in sera of approximately 200 HCV cases. Only 2 sera reacted slightly with the env35 peptide. However, by means of the anti-E1 ELISA as described in example 6, it was possible to discover additional epitopes as follows: The anti-E1 ELISA as described in example 6 was modified by mixing 50 µg/ml of E1 peptide with the 1/20 diluted human serum in sample diluent. Figure 13 shows the results of reactivity of human sera to the recombinant E1 (expressed from vvHCV-40) protein, in the presence of single or of a mixture of E1 peptides. While only 2% of the sera could be detected by means of E1 peptides coated on strips in a Line Immunoassay format, over half of the sera contained anti-E1 antibodies which could be competed by means of the same peptides, when tested on the recombinant E1 protein. Some of the murine monoclonal antibodies obtained from Balb/C mice after injection with purified E1 protein were subsequently competed for reactivity to E1 with the single peptides (Figure 14). Clearly, the region of env53 contained the predominant epitope, as the addition of env53 could substantially compete reactivity of several sera with E1, and antibodies to the env31 region were also detected. This finding was surprising, since the env53 and env31 peptides had not shown any reactivity when coated directly to the solid phase.

Therefore peptides were synthesized using technology described by applicant previously (in WO 93/18054). The following peptides were synthesized:

peptide env35A-biotin

NH<sub>2</sub>-SNSSEAADMIMHTPGCV-GKbiotin (SEQ ID NO 51)

spanning amino acids 208 to 227 of the HCV polyprotein in the E1 region

peptide biotin-env53 ('epitope A')

biotin-GG-ITGHRMAWDMMMNWSPTTAL-COOH (SEQ ID NO 52)

spanning amino acids 313 to 332 of the HCV polyprotein in the E1 region

peptide 1bE1 ('epitope B')

H<sub>2</sub>N-YEVRNVSGIYHVTNDCSNSSIVYEAADMIMHTPGCGK-biotin (SEQ ID NO 53)

spanning amino acids 192 to 228 of the HCV polyprotein in the E1 region

and compared with the reactivities of peptides E1a-8B (biotin-GG-TPTVATRDGKLPATQLRRHIDL, SEQ ID NO 54) and E1b-8B (biotin-GG-TPTLAARDASVPTTTIRRHVDLL, SEQ ID NO 55) which are derived from the same region of sequences of genotype 1a and 1b respectively and which have been described at the IXth international virology meeting in Glasgow, 1993 ('epitope C'). Reactivity of a panel of HCV sera was tested on

epitopes A, B and C and epitope B was also compared with env35A (of 47 HCV-positive sera, 8 were positive on epitope B and none reacted with env35A). Reactivity towards epitopes A, B, and C was tested directly to the biotinylated peptides (50 µg/ml) bound to streptavidin-coated plates as described in example 6. Clearly, epitopes A and B were most reactive while epitopes C and env35A-biotin were much less reactive. The same series of patients that had been monitored for their reactivity towards the complete E1 protein (example 7.1.) was tested for reactivity towards epitopes A, B, and C. Little reactivity was seen to epitope C, while as shown in Figures 15, 16, 17, and 18, epitopes A and B reacted with the majority of sera. However, antibodies to the most reactive epitope (epitope A) did not seem to predict remission of disease, while the anti-1bE1 antibodies (epitope B) were present almost exclusively in long term responders at the start of IFN treatment. Therefore, anti-1bE1 (epitope B) antibodies and anti-env53 (epitope A) antibodies could be shown to be useful markers for prognosis of hepatitis C disease. The env53 epitope may be advantageously used for the detection of cross-reactive antibodies (antibodies that cross-react between major genotypes) and antibodies to the env53 region may be very useful for universal E1 antigen detection in serum or liver tissue. Monoclonal antibodies that recognized the env53 region were reacted with a random epitope library. In 4 clones that reacted upon immunoscreening with the monoclonal antibody 5E1A10, the sequence -GWD- was present. Because of its analogy with the universal HCV sequence present in all HCV variants in the env53 region, the sequence AWD is thought to contain the essential sequence of the env53 cross-reactive murine epitope. The env31 clearly also contains a variable region which may contain an epitope in the amino terminal sequence -YQVRNSTGL- (SEQ ID NO 93) and may be useful for diagnosis. Env31 or E1-31 as shown in Table 3, is a part of the peptide 1bE1. Peptides E1-33 and E1-51 also reacted to some extent with the murine antibodies, and peptide E1-55 (containing the variable region 6 (V6); spanning amino acid positions 329-336) also reacted with some of the patient sera.

Anti-E2 antibodies clearly followed a different pattern than the anti-E1 antibodies, especially in patients with a long-term response to treatment. Therefore, it is clear that the decrease in anti-envelope antibodies could not be measured as efficiently with an assay employing a recombinant E1/E2 protein as with a single anti-E1 or anti-E2 protein. The anti-E2 response would clearly blur the anti-E1 response in an assay measuring both kinds of antibodies at the same time. Therefore, the ability to test anti-envelope antibodies to the single E1 and E2 proteins, was shown to be useful.

#### 7.4. Mapping of anti-E2 antibodies

Of the 24 anti-E2 Mabs only three could be competed for reactivity to recombinant E2 by peptides, two of which reacted with the HVRI region (peptides E2-67 and E2-69, designated as epitope A) and one which recognized an epitope competed by peptide E2-13B (epitope C). The majority of murine antibodies recognized conformational anti-E2 epitopes (Figure 19). A human response to HVRI (epitope A), and to a lesser extent

HVRII (epitope B) and a third linear epitope region (competed by peptides E2-23, E2-25 or E2-27, designated epitope E) and a fourth linear epitope region (competed by peptide E2-17B, epitope D) could also frequently be observed, but the majority of sera reacted with conformational epitopes (Figure 20). These conformational epitopes could be grouped according to their relative positions as follows: the IgG antibodies in the supernatant of hybridomas 15C8C1, 12D11F1, 9G3E6, 8G10D1H9, 10D3C4, 4H6B2, 17F2C2, 5H6A7, 15B7A2 recognizing conformational epitopes were purified by means of protein A affinity chromatography and 1 mg/ml of the resulting IgG's were biotinylated in borate buffer in the presence of biotin. Biotinylated antibodies were separated from free biotin by means of gelfiltration chromatography. Pooled biotinylated antibody fractions were diluted 100 to 10,000 times. E2 protein bound to the solid phase was detected by the biotinylated IgG in the presence of 100 times the amount of non-biotinylated competing antibody and subsequently detected by alkaline phosphatase labeled streptavidin.

Percentages of competition are given in Table 6. Based on these results, 4 conformational anti-E2 epitope regions (epitopes F, G, H and I) could be delineated (Figure 38). Alternatively, these Mabs may recognize mutant linear epitopes not represented by the peptides used in this study. Mabs 4H6B2 and 10D3C4 competed reactivity of 16A6E7, but unlike 16A6E7, they did not recognize peptide E2-13B. These Mabs may recognize variants of the same linear epitope (epitope C) or recognize a conformational epitope which is sterically hindered or changes conformation after binding of 16A6E7 to the E2-13B region (epitope H).

### Example 8: E1 glycosylation mutants

#### 8.1. Introduction

The E1 protein encoded by vvhCV10A, and the E2 protein encoded by vvhCV41 to 44 expressed from mammalian cells contain 6 and 11 carbohydrate moieties, respectively. This could be shown by incubating the lysate of vvhCV10A-infected or vvhCV44-infected RK13 cells with decreasing concentrations of glycosidases (PNGase F or Endoglycosidase H, (Boehringer Mannheim Biochemica) according to the manufacturer's instructions), such that the proteins in the lysate (including E1) are partially deglycosylated (Fig. 39 and 40, respectively).

Mutants devoid of some of their glycosylation sites could allow the selection of envelope proteins with

improved immunological reactivity. For HIV for example, gp120 proteins lacking certain selected sugar-addition motifs, have been found to be particularly useful for diagnostic or vaccine purpose. The addition of a new oligosaccharide side chain in the hemagglutinin protein of an escape mutant of the A/Hong Kong/3/68 (H3N2) influenza virus prevents reactivity with a neutralizing monoclonal antibody (Skehel et al, 1984). When novel glycosylation sites were introduced into the influenza hemagglutinin protein by site-specific mutagenesis, dramatic antigenic changes were observed, suggesting that the carbohydrates serve as a modulator of antigenicity (Gallagher et al., 1988). In another analysis, the 8 carbohydrate-addition motifs of the surface protein gp70 of the Friend Murine Leukemia Virus were deleted. Although seven of the mutations did not affect virus infectivity, mutation of the fourth glycosylation signal with respect to the amino terminus resulted in a non-infectious phenotype (Kayman et al., 1991). Furthermore, it is known in the art that addition of N-linked carbohydrate chains is important for stabilization of folding intermediates and thus for efficient folding, prevention of misfolding and degradation in the endoplasmic reticulum, oligomerization, biological activity, and transport of glycoproteins (see reviews by Rose et al., 1988; Doms et al., 1993; Helenius, 1994).

After alignment of the different envelope protein sequences of HCV genotypes, it may be inferred that not all 6 glycosylation sites on the HCV subtype 1b E1 protein are required for proper folding and reactivity, since some are absent in certain (sub)types. The fourth carbohydrate motif (on Asn251), present in types 1b, 6a, 7, 8, and 9, is absent in all other types known today. This sugar-addition motif may be mutated to yield a type 1b E1 protein with improved reactivity. Also the type 2b sequences show an extra glycosylation site in the V5 region (on Asn299). The isolate S83, belonging to genotype 2c, even lacks the first carbohydrate motif in the V1 region (on Asn), while it is present on all other isolates (Stuyver et al., 1994). However, even among the completely conserved sugar-addition motifs, the presence of the carbohydrate may not be required for folding, but may have a role in evasion of immune surveillance. Therefore, identification of the carbohydrate addition motifs which are not required for proper folding (and reactivity) is not obvious, and each mutant has to be analyzed and tested for reactivity. Mutagenesis of a glycosylation motif (NXS or NXT sequences) can be achieved by either mutating the codons for N, S, or T, in such a way that these codons encode amino acids different from N in the case of N, and/or amino acids different from S or T in the case of S and in the case of T. Alternatively, the X position may be mutated into P, since it is known that NPS or NPT are not frequently modified with carbohydrates. After establishing which carbohydrate-addition motifs are required for folding and/or reactivity and which are not, combinations of such mutations may be made.

30

## 8.2. Mutagenesis of the E1 protein

All mutations were performed on the E1 sequence of clone HCC110A (SEQ ID NO. 5). The first round of PCR was performed using sense primer 'GPT' (see Table 7) targeting the GPT sequence located upstream of

the vaccinia 11K late promoter, and an antisense primer (designated GLY#, with # representing the number of the glycosylation site, see Fig. 41) containing the desired base change to obtain the mutagenesis. The six GLY# primers (each specific for a given glycosylation site) were designed such that:

- Modification of the codon encoding for the N-glycosylated Asn (AAC or AAT) to a Gln codon (CAA or CAG).
- 5 Glutamine was chosen because it is very similar to asparagine (both amino acids are neutral and contain non-polar residues, glutamine has a longer side chain (one more -CH<sub>2</sub>- group).
- The introduction of silent mutations in one or several of the codons downstream of the glycosylation site, in order to create a new unique or rare (e.g. a second SmaI site for E1Gly5) restriction enzyme site. Without modifying the amino acid sequence, this mutation will provide a way to distinguish the mutated sequences from the original E1 sequence (pvHCV-10A) or from each other (Figure 41). This additional restriction site may also
- 10 be useful for the construction of new hybrid (double, triple, etc.) glycosylation mutants.
- 18 nucleotides extend 5' of the first mismatched nucleotide and 12 to 16 nucleotides extend to the 3' end. Table 7 depicts the sequences of the six GLY# primers overlapping the sequence of N-linked glycosylation sites.

For site-directed mutagenesis, the 'mispriming' or 'overlap extension' (Horton, 1993) was used. The concept is illustrated in Figures 42 and 43. First, two separate fragments were amplified from the target gene for each mutated site. The PCR product obtained from the 5' end (product GLY#) was amplified with the 5' sense GPT primer (see Table 7) and with the respective 3' antisense GLY# primers. The second fragment (product OVR#) was amplified with the 3' antisense TK<sub>A</sub> primer and the respective 5' sense primers (OVR# primers, see Table 7, Figure 43).

20 The OVR# primers target part of the GLY# primer sequence. Therefore, the two groups of PCR products share an overlap region of identical sequence. When these intermediate products are mixed (GLY-1 with OVR-1, GLY-2 with OVR-2, etc.), melted at high temperature, and reannealed, the top sense strand of product GLY# can anneal to the antisense strand of product OVR# (and vice versa) in such a way that the two strands act as primers for one another (see Fig. 42.B.). Extension of the annealed overlap by Taq polymerase

25 during two PCR cycles created the full-length mutant molecule E1GLY#, which carries the mutation destroying the glycosylation site number #. Sufficient quantities of the E1GLY# products for cloning were generated in a third PCR by means of a common set of two internal nested primers. These two new primers are respectively overlapping the 3' end of the vaccinia 11K promoter (sense GPT-2 primer) and the 5' end of the vaccinia thymidine kinase locus (antisense TK<sub>A</sub>-2 primer, see Table 7). All PCR conditions were performed as described

30 in Stuyver et al. (1993).

Each of these PCR products was cloned by EcoRI/BamHI cleavage into the EcoRI/BamHI-cut vaccinia vector containing the original E1 sequence (pvHCV-10A).

The selected clones were analyzed for length of insert by EcoRI/BamHI cleavage and for the presence of each new restriction site. The sequences overlapping the mutated sites were confirmed by double-stranded

sequencing.

### 8.3. Analysis of E1 glycosylation mutants

5 Starting from the 6 plasmids containing the mutant E1 sequences as described in example 8.2, recombinant vaccinia viruses were generated by recombination with wt vaccinia virus as described in example 2.5. Briefly, 175 cm<sup>2</sup>-flasks of subconfluent RK13 cells were infected with the 6 recombinant vaccinia viruses carrying the mutant E1 sequences, as well as with the vvHCV-10A (carrying the non-mutated E1 sequence) and wt vaccinia viruses. Cells were lysed after 24 hours of infection and analyzed on western blot as described in  
10 example 4 (see Figure 44A). All mutants showed a faster mobility (corresponding to a smaller molecular weight of approximately 2 to 3 kDa) on SDS-PAGE than the original E1 protein; confirming that one carbohydrate moiety was not added. Recombinant viruses were also analyzed by PCR and restriction enzyme analysis to confirm the identity of the different mutants. Figure 44B shows that all mutants (as shown in Figure 41) contained the expected additional restriction sites. Another part of the cell lysate was used to test the reactivity of  
15 the different mutant by ELISA. The lysates were diluted 20 times and added to microwell plates coated with the lectin GNA as described in example 6. Captured (mutant) E1 glycoproteins were left to react with 20-times diluted sera of 24 HCV-infected patients as described in example 6. Signal to noise (S/N) values (OD of GLY#/OD of wt) for the six mutants and E1 are shown in Table 8. The table also shows the ratios between S/N values of GLY# and E1 proteins. It should be understood that the approach to use cell lysates of the different  
20 mutants for comparison of reactivity with patient sera may result in observations that are the consequence of different expression levels rather than reactivity levels. Such difficulties can be overcome by purification of the different mutants as described in example 5, and by testing identical quantities of all the different E1 proteins. However, the results shown in table 5 already indicate that removal of the 1st (GLY1), 3rd (GLY3), and 6th (GLY6) glycosylation motifs reduces reactivity of some sera, while removal of the 2nd and 5th site does not.  
25 Removal of GLY4 seems to improve the reactivity of certain sera. These data indicate that different patients react differently to the glycosylation mutants of the present invention. Thus, such mutant E1 proteins may be useful for the diagnosis (screening, confirmation, prognosis, etc.) and prevention of HCV disease.

### 30 Example 9: Expression of HCV E2 protein in glycosylation-deficient yeasts

The E2 sequence corresponding to clone HCCL41 was provided with the  $\alpha$ -mating factor pre/pro signal sequence, inserted in a yeast expression vector and *S. cerevisiae* cells transformed with this construct secreted E2 protein into the growth medium. It was observed that most glycosylation sites were modified with

high-mannose type glycosylations upon expression of such a construct in *S. cerevisiae* strains (Figure 45). This resulted in a too high level of heterogeneity and in shielding of reactivity, which is not desirable for either vaccine or diagnostic purposes. To overcome this problem, *S. cerevisiae* mutants with modified glycosylation pathways were generated by means of selection of vanadate-resistant clones. Such clones were analyzed for modified glycosylation pathways by analysis of the molecular weight and heterogeneity of the glycoprotein invertase. This allowed us to identify different glycosylation deficient *S. cerevisiae* mutants. The E2 protein was subsequently expressed in some of the selected mutants and left to react with a monoclonal antibody as described in example 7, on western blot as described in example 4 (Figure 46).

#### 10 Example 10. General utility

The present results show that not only a good expression system but also a good purification protocol are required to reach a high reactivity of the HCV envelope proteins with human patient sera. This can be obtained using the proper HCV envelope protein expression system and/or purification protocols of the present invention which guarantee the conservation of the natural folding of the protein and the purification protocols of the present invention which guarantee the elimination of contaminating proteins and which preserve the conformation, and thus the reactivity of the HCV envelope proteins. The amounts of purified HCV envelope protein needed for diagnostic screening assays are in the range of grams per year. For vaccine purposes, even higher amounts of envelope protein would be needed. Therefore, the vaccinia virus system may be used for selecting the best expression constructs and for limited upscaling, and large-scale expression and purification of single or specific oligomeric envelope proteins containing high-mannose carbohydrates may be achieved when expressed from several yeast strains. In the case of hepatitis B for example, manufacturing of HBsAg from mammalian cells was much more costly compared with yeast-derived hepatitis B vaccines.

The purification method disclosed in the present invention may also be used for 'viral envelope proteins' in general. Examples are those derived from Flaviviruses, the newly discovered GB-A, GB-B and GB-C Hepatitis viruses, Pestiviruses (such as Bovine viral Diarrhoea Virus (BVDV), Hog Cholera Virus (HCV), Border Disease Virus (BDV)), but also less related viruses such as Hepatitis B Virus (mainly for the purification of HBsAg).

The envelope protein purification method of the present invention may be used for intra- as well as extracellularly expressed proteins in lower or higher eukaryotic cells or in prokaryotes as set out in the detailed description section.

#### Example 11. Demonstration of Prophylactic and Therapeutic Utility

Liver disease in chimpanzees chronically infected with HCV can be reduced by immunization with E1. Multiple immunizations, however, were required in order to reach a significant immune response. One of ordinary skill will appreciate that viral persistence is produced with immune modulation which is either orchestrated by the virus itself or by the host. In order to analyze if such an immune modulation does exist in HCV, the immune responses against E1 and NS3 in naive and chronically infected chimpanzees were compared. Since a lower response in the chronically infected animals was anticipated, this group of animals was selected for a more rigorous immunization schedule including the following: use of an adjuvant proven in mice to be more potent for inducing cellular responses (Table 9) compared to alum, which was the adjuvant used for naive animals; and the immunization schedule for chronically infected animals consisted of 12 immunizations compared to 6 for naive animals (Fig. 47).

Although the number of immunized animals does not allow statistical analysis, the following clear tendency can be detected in the humoral responses (Table 10): the number of immunizations for seroconversion is lower in naive animals; and the magnitude of the immune response is substantially greater in the naive animals, 2/3 infected animals do not reach the level of 10 internal units, even after 12 immunizations.

The analysis of the cellular responses, after three immunizations, reveals an even larger difference (Fig. 48a-d), including the following: E1-specific T-cell proliferation is almost absent in the chronically infected animals, while a clear stimulation can be seen in the naive setting; IL-2 measurements confirmed that the low stimulation of the T-cell compartment in chronic carriers; and, a clear Th2 (IL-4) response in naive animals is induced, as expected for an alum-adjuvant containing vaccine.

This confirms that at least E1 immunization provides a prophylactic effect in naive animals and suggest that E2 and/or combinations of E1 and E2 proteins and/or peptides may provide useful therapeutic and/or prophylactic benefits in naive animals.

The 'impairment' to induce both cellular and humoral responses against an HCV E1 antigen can be only partially overcome by multiple immunizations, as demonstrated by the following results: an increase in antibody titer after each injection was noted but the levels as in naive animals were not reached in 2/3 animals; and the T-cell proliferative responses remain very low (Fig. 49). The ELISPOT results show, however, a minor increase in IL-2 (not shown), no change in IFN- $\gamma$  (not shown) and an increase in IL-4 (Fig. 49) which indicates that Th2 type responses are more readily induced. IL-4 was noted to remain at a low level compared to the level reached after three immunizations in naive animals.

A quite similar observation was made for NS3 immunizations where an even stronger adjuvant (RIBI) was used in the chronic chimpanzee. As compared with an alum formulation in naive animals the following has been noted: the induced antibody titers are comparable in both groups (not shown); and both cytokine secretion and T-cell proliferation are almost absent in the chronic animals compared to the responses in naive animals (Fig. 49a-b).

Currently there have been some indications that immune responses against HCV in chronic carriers are low or at least insufficient to allow clearance of infection. The above results support the hypothesis that the immune system of HCV chronic carriers may be impaired and that they do not respond to HCV antigens as efficiently as in a naive situation.

In a study by Wiedmann et al., (Hepatology 2000; 31: 230-234), vaccination for HBV was less effective in HCV chronic carriers, which indicates that such an immune impairment is not limited to HCV antigens. De Maria et al.



(Hepatology 2000; 32: 444-445) confirmed these data and have proposed adapted vaccine dosing regimens for HCV patients. The data presented herein indicates that increasing the number of immunizations may indeed augment humoral responses but that cellular (especially Th1) responses are difficult to induce, even when powerful adjuvants are used. It may be advantageous to begin immunization at the time of antiviral therapy, when the immune system is more prone to respond.

Table 1: Recombinant vaccinia plasmids and viruses

Plasmid name	Name	cDNA subclone construction	Length (nt/aa)	Vector used for insertion
pvHCV-13A	E1s	EcoR I - Hind III	472/157	pgptATA-18
pvHCV-12A	E1s	EcoR I - Hind III	472/158	pgptATA-18
pvHCV-9A	E1	EcoR I - Hind III	631/211	pgptATA-18
pvHCV-11A	E1s	EcoR I - Hind III	625/207	pgptATA-18
pvHCV-17A	E1s	EcoR I - Hind III	625/208	pgptATA-18
pvHCV-10A	E1	EcoR I - Hind III	783/262	pgptATA-18
pvHCV-18A	COREs	Acc I (KI) - EcoR I (KI)	403/130	pgptATA-18
pvHCV-34	CORE	Acc I (KI) - Fsp I	595/197	pgptATA-18
pvHCV-33	CORE-E1	Acc I (KI)	1150/380	pgptATA-18
pvHCV-35	CORE-E1b.his	EcoR I - BamH I (KI)	1032/352	pMS-66
pvHCV-36	CORE-E1n.his	EcoR I - Nco I (KI)	1106/376	pMS-66
pvHCV-37	E1 $\Delta$	Xma I - BamH I	711/239	pvHCV-10A
pvHCV-38	E1 $\Delta$ s	EcoR I - EstE II	553/183	pvHCV-11A
pvHCV-39	E1 $\Delta$ b	EcoR I - BamH I	960/313	pgsATA-18
pvHCV-40	E1 $\Delta$ b.his	EcoR I - BamH I (KI)	960/323	pMS-66
pvHCV-41	E2bs	BamH I (KI)-AlwN I (T4)	1005/331	pgsATA-18
pvHCV-42	E2bs.his	BamH I (KI)-AlwN I (T4)	1005/341	pMS-66
pvHCV-43	E2ns	Nco I (KI) - AlwN I (T4)	932/314	pgsATA-18
pvHCV-44	E2ns.his	Nco I (KI) - AlwN I (T4)	932/321	pMS-66
pvHCV-62	E1s (type 3a)	EcoR I - Hind III	625/207	pgsATA-18
pvHCV-63	E1s (type 5)	EcoR I - Hind III	625/207	pgsATA-18
pvHCV-64	E2	BamH I - Hind III	1410/463	pgsATA-18
pvHCV-65	E1-E2	BamH I - Hind III	2072/691	pvHCV-10A
pvHCV-66	CORE-E1-E2	BamH I - Hind III	2427/809	pvHCV-33

nt: nucleotide aa: aminoacid KI: Klenow DNA Pol filling T4: T4 DNA Pol filling  
Position: aminoacid position in the HCV polypeptide sequence

Table 1 - continued: Recombinant vaccinia plasmids and viruses

Plasmid Name	HCV cDNA subclone			Vector used for insertion
	Name	Construction	Length (nt/aa)	
pvHCV-81	E1*-GLY 1	EcoRI - BamH I	783/262	pvHCV-10A
pvHCV-82	E1*-GLY 2	EcoRI - BamH I	783/262	pvHCV-10A
pvHCV-83	E1*-GLY 3	EcoRI - BamH I	783/262	pvHCV-10A
pvHCV-84	E1*-GLY 4	EcoRI - BamH I	783/262	pvHCV-10A
pvHCV-85	E1*-GLY 5	EcoRI - BamH I	783/262	pvHCV-10A
pvHCV-86	E1*-GLY 6	EcoRI - BamH I	783/262	pvHCV-10A

nt: nucleotide    aa: aminoacid    KI: Klenow DNA Pol filling    T4: T4 DNA Pol filling  
Position: aminoacid position in the HCV polyprotein sequence

Table 2 : Summary of anti-E1 tests

S/N  $\pm$  SD (mean anti-E1 titer)

	Start of treatment	End of treatment	Follow-up
LTR	6.94 $\pm$ 2.29 (1:3946)	4.48 $\pm$ 2.69 (1:568)	2.99 $\pm$ 2.69 (1:175)
NR	5.77 $\pm$ 3.77 (1:1607)	5.29 $\pm$ 3.99 (1:1060)	6.08 $\pm$ 3.73 (1:1978)

LTR : Long-term, sustained response for more than 1 year

NR : No response, response with relapse, or partial response

**Table 3****Synthetic peptides for competition studies**

PROTEIN	PEPTIDE	AMINO ACID SEQUENCE	POSITION	SEQ ID NO
E1	E1-31	LLSCLTPASAYQVRNSTGL	181-200	56
	E1-33	QVRNSTGLYHVTNDCPNSSI	193-212	57
	E1-35	NDCPNSSIVYEAHDAILHTP	205-224	58
	E1-35A	SNSSIVYEAADMIMHTPGCV	208-227	59
	E1-37	HDAILHTPGCVPCVREGNVS	217-236	60
	E1-39	CVREGNVSRCWVAMTPTVAT	229-248	61
	E1-41	AMTPTVATRDGKLPATQLRR	241-260	62
	E1-43	LPATQLRRHIDLLVGSATLC	253-272	63
	E1-45	LVGSATLCSALYVGDLGGSV	265-284	64
	E1-49	QLFTFSPRRHWTTOGCNCSI	289-308	65
	E1-51	TQGCNCSIYPGHITGHRMAW	301-320	66
	E1-53	ITGHRMAWDMMNWSPTAAL	313-332	67
	E1-55	NWSPTAALVMAQLLRIPQAI	325-344	68
	E1-57	LLRIPQAILDMIAGAHWGV	337-356	69
	E1-59	AGAHWGVLAGIAYFSMVGNM	349-368	70
	E1-63	VVLLFAGVDAETIVSGGQA	373-392	71
E2	E2-67	SGLVSLFTPGAKQNIQLINT	397-416	72
	E2-69	QNIQLINTNGSWHINSTALN	409-428	73
	E2-33B	LNCNESLNTGWWLAGLIYQHK	427-446	74
	E2-11B	AGLIYQHKFNSSGCPERLAS	439-458	75
	E2-1B	GCPERLASCRPLTDFQGGWG	451-470	76
	E2-3B	TDFQGGWGPISYANGSGPOQ	463-482	77
	E2-5B	ANGSGEDQRPYCWHYPPKPC	475-494	78
	E2-7B	WHYPPKPCGIVPAKSVCGPV	487-506	79
	E2-9B	AKSVCGPVYCFTPSPVVVGT	499-518	80
	E2-11B	PSPVVVGTDRSGAPTYSWG	511-530	81
	E2-13B	GAPTYSWGENDTDFVFLNNT	523-542	82

E2-17B	GNWFGCTWMNSTGFTKVC GA	547-566	83
E2-19B	GFTKVC GAPPVCIGGAGNNT	559-578	84
E2-21	IGGAGNNTLHCPTDCFRKHP	571-590	85
E2-23	TDCFRKHPDATYSRCGSGPW	583-602	86
E2-25	SRCGSGPWITPRCLVDYPYR	595-614	87
E2-27	CLVDYPYRLWHYPCTINYTI	607-626	88
E2-29	PCTINYTIFKIRMYVGGVEH	619-638	89
E2-31	MYVGGVEHRLEAACNWT PGE	631-650	90
E2-33	ACNWT PGERCDLED RDRSEL	643-662	91
E2-35	EDRDRSELSP LLLTTTQWQV	655-674	92



Table 5. Difference between LTR and NR (complete study)

Mann-Whitney U test (P values)	E1Ab S/N All	E1Ab titers All	E1Ab S/N type 1b	E1Ab S/N type 3a	E1Ab S/N All	E2Ab S/N
Initiation of therapy	0.0257	0.05	0.68		0.1078	
End of therapy	0.1742				0.1295	
6 months follow up	1	0.6099	0.425		0.3081	
12 months follow up	0.67	0.23	0.4386		0.6629	

\* P values &lt; 0.05



**Table 6. Competition experiments between murine E2 monoclonal antibodies**

Decrease (%) of anti-E2 reactivity of biotinylated anti-E2 mabs										
compellor	17H10F4D10	2F10H10	16A6E7	10D3C4	4H6B2	17C2F2	9G3E6	12D11F1	15C8C1	8G10D1H9
17H10F4D10	-	62	10	ND	11	ND	5	6	30	ND
2F10H10	90	-	1	ND	30	ND	0	4	12	ND
16A6E7	ND	ND	-	ND	ND	ND	ND	ND	ND	ND
10D3C4	11	50	92	-	94	28	28	43	53	30
4H6B2	ND	ND	82	ND	-	ND	ND	ND	ND	ND
17C2F2	2	ND	75	ND	56	-	11	10	0	0
9G3E6	ND	ND	88	ND	11	ND	-	60	76	ND
12D11F1	ND	ND	26	ND	13	ND	ND	-	88	ND
15C8C1	ND	ND	18	ND	10	ND	ND	ND	-	ND
8G10D1H9	2	2	11	ND	15	ND	67	082	81	-
compellor controls										
15B7A2	0	0	9	15	10	9	0	0	0	5
5H6A7	0	2	0	12	8	0	0	4	0	0
23C12H9	ND	ND	2	12	ND	4	ND	ND	ND	2
ND = not done										

Table 7. Primers

5	SEQ ID NO. 96	GPT	5'-GTTTAACCACTGCATGATG-3'
	SEQ ID NO. 97	TK <sub>n</sub>	5'-GTCCATCGAGTGGGGTAC-3'
	SEQ ID NO. 98	GLY1	5'-CGTGACATGGTACATTCGGGACACTTGGGCACITTCATAAGCGGA-3'
	SEQ ID NO. 99	GLY2	5'-TGCCTCATACACAATGGAGCTCTGGGACGAGTCGTTCTGTGAC-3'
	SEQ ID NO. 100	GLY3	5'-TACCCAGCAGCGGGAGCCTCTGCTCCCGAACGCGAGGGGAC-3'
10	SEQ ID NO. 101	GLY4	5'-TGTCGTGTGGGACGGAGGGCTGCCCTAGCTGGAGCGGTGGG-3'
	SEQ ID NO. 102	GLY5	5'-CGTTAAGTGCCCGGGTAGATTGAGCACTGGCAGTCCCTGCACCGTCTC-3'
	SEQ ID NO. 103	GLY6	5'-CAGGGCCGTTGTAGGCCCTCCACTGCATCATCATATCCCAAGC-3'
	SEQ ID NO. 104	OVR1	5'-CCGGAATGTACCATGTCACGAACGAC-3'
	SEQ ID NO. 105	OVR2	5'-GCTCCATTGTGTATGAGGCAGCGG-3'
15	SEQ ID NO. 106	OVR3	5'-GAGCTCCCGCTGCTGGGTAGCGC-3'
	SEQ ID NO. 107	OVR4	5'-CCTCCGTCCCACCACGACAATACG-3'
	SEQ ID NO. 108	OVR5	5'-CTACCCGGGCCACATAACGGGTACCG-3'
	SEQ ID NO. 109	OVR6	5'-GGAGGGCTACAACGGCCCTGGTGG-3'
	SEQ ID NO. 110	GPT-2	5'-TTCTATCGATTAAATAGAATTC-3'
20	SEQ ID NO. 111	TK <sub>n</sub> -2	5'-GCCATACGCTCACAGCCGATCCC-3'

nucleotides underlined represent additional restriction site

20 nucleotides in bold represent mutations with respect to the original HCC10A sequence

Table B. Analysis of E1 glycosylation mutants by ELISA

SERUM													Sum	Average
1	2	3	4	5	6	7	8	9	10	11	12	24		
SN GLY1	1.802402	2.120971	1.403871	1.205597	2.120191	2.806913	1.950345	1.860103	1.730103	2.400162	1.220054	1.629403		
SN GLY2	2.400786	1.76818	2.328486	2.839308	2.458019	6.043893	2.148302	1.580477	1.608873	2.402212	1.487502	2.070524		
SN GLY3	1.042716	1.716477	2.261646	2.354748	1.661816	4.033742	1.90092	1.402099	1.602222	2.101858	1.404210	1.721104		
SN GLY4	2.570154	3.824038	3.874805	1.489387	3.15	4.71302	4.198761	3.958542	3.710507	5.170841	4.250704	3.955153		
SN GLY5	2.482051	1.783781	2.409344	2.627358	1.716311	4.894765	2.13912	1.570336	1.708937	3.021807	1.562092	2.07278		
SN GLY6	2.031487	1.485737	2.131613	2.627925	2.494833	4.704027	2.02089	1.498480	1.704876	2.877757	1.529808	1.744221		
SN E1	2.828205	2.227038	2.512792	2.780881	3.131579	4.889128	2.287763	1.954198	1.805556	2.616822	1.55718	2.593886		
Sum														
Average														
SN GLY1	5.885581	3.233604	3.763488	1.885105	2.317721	8.875178	1.93470	2.47171	4.378033	1.188748	2.186880	1.708992	59.88534	2.495223
SN GLY2	7.558892	2.567813	3.821928	3.055849	2.933792	7.05433	2.127712	2.921268	4.680101	1.160781	1.861914	1.032705	68.65243	2.902185
SN GLY3	7.930538	3.018099	2.945828	2.515305	5.775357	1.880186	2.557384	4.260033	0.97707	1.306776	1.368776	1.703378	82.09872	2.607447
SN GLY4	8.178816	6.581122	6.707688	5.684498	5.804813	6.4125	3.813321	3.002535	4.293038	2.393011	3.08713	2.481585	102.0970	4.279076
SN GLY5	8.883408	2.940334	3.125501	3.330912	2.654224	5.424107	2.442004	3.128761	1.153858	1.817901	1.038211	69.26511	2.888048	
SN GLY6	8.005581	2.499952	2.621704	2.672385	2.363301	6.194107	1.508718	2.685433	2.781003	1.280743	1.475082	1.716423	61.32181	2.655076
SN E1	8.826112	3.183771	3.087285	3.280335	2.880354	7.181884	2.771218	3.878008	6.36443	1.167286	2.083333	1.78252	76.54088	3.109195

SERUM													Sum	Average
1	2	3	4	5	6	7	8	9	10	11	12	24		
GLY1/E1	0.837316	0.952374	0.65869	0.431977	0.877038	0.688794	0.852516	0.954801	0.958261	0.94319	0.783892	0.628171		
GLY2/E1	0.848876	0.783901	0.825483	0.94569	0.785233	1.035913	0.93817	0.816436	0.935431	0.94856	0.842456	0.798232		
GLY3/E1	0.580834	0.770288	0.900053	0.84373	0.508312	0.882733	0.859781	0.768418	0.807385	0.037488	0.940294	0.803547		
GLY4/E1	0.811587	1.717087	1.541862	0.837245	1.005882	0.987939	1.835317	2.028172	2.05505	1.978	2.72978	1.524798		
GLY5/E1	0.877007	0.805447	0.958831	0.941408	0.547746	1.018842	0.935031	0.806641	0.948488	1.154782	1.003148	0.799102		
GLY6/E1	0.718296	0.871828	0.848305	0.905578	0.798688	0.882622	0.883284	0.785781	0.944284	1.023200	0.982280	0.872435		
Sum														
Average														
GLY1/E1	0.644248	1.015852	1.226988	0.605153	0.777688	0.828144	0.898182	0.072013	0.817759	1.018300	1.030267	0.957628	19.36524	0.806005
GLY2/E1	0.85827	0.808489	1.180833	0.931505	0.944377	1.054289	0.78779	0.794245	0.874081	0.98586	0.787719	0.815888	21.67304	0.903077
GLY3/E1	0.998033	0.807858	0.903319	0.887968	0.843962	0.803029	0.714554	0.695308	0.757215	0.837558	0.641052	0.675314	19.19921	0.709167
GLY4/E1	0.92654	2.005802	1.060833	1.732902	1.880587	0.88102	1.376045	0.810335	0.801773	2.050084	1.787422	1.392178	36.38592	1.51600
GLY5/E1	1.006608	0.923538	1.019008	1.017857	0.890574	0.75418	0.861491	0.850109	0.867612	0.988323	0.872593	0.918042	21.78879	0.907703
GLY6/E1	0.907134	0.786217	0.854737	0.784184	0.792266	0.72221	0.643702	0.724603	0.518335	1.097197	0.70003	0.962918	18.59681	0.816330

Table 9. Profile of adjuvanted E1 in Balb/c mice

	alum	T-cell adjuvant	RBI
antibody titre (mean $\pm$ SD, n=6)	96000 $\pm$ 101000	62000 $\pm$ 60000	176000 $\pm$ 149000
antibody isotypes	IgG1	IgG1/2b	IgG1/2a
T-cell proliferation in spleen <sup>1</sup> (n=3)	11750 (2/3)	48300 (3/3)	26000 (3/3)
T-cell proliferation in lymph node <sup>2</sup>	no specific stimulation	4000	8000
cytokine profile (spleen)	IL-4	IFN- $\gamma$ /IL-4	IFN- $\gamma$ /IL-4

5 <sup>1</sup> after three s.c./i.m. immunizations, 3 randomly selected mice were analyzed individually, the result is expressed as the mean specific cpm obtained after 4 days of E1 stimulation (1  $\mu$ g/ml), the number in brackets refers to the number of mice with specific stimulation above background

10 <sup>2</sup> after one single intra footpath immunization (n=2), the result is expressed as the mean specific cpm obtained after 5 days of E1 stimulation (1  $\mu$ g/ml)

Table 10. Humoral Responses: No. of immunizations required for different E-1 antibodies levels

Animal	status	seroconversion <sup>1</sup>	> 1 U/ml <sup>2</sup>	> 10 <sup>3</sup> U/ml
Marcel	chronic	3	4	5
Peggy	chronic	3	5	>12
Femma	chronic	4	5	>12
Yoran	naive	3	4	5
Marti	naive	2	3	5

5

<sup>1</sup> defined as ELISA signal higher than cut-off level if no E1-antibodies were present prior to immunization, in the other cases the observation of a titer higher than the 3 individual time points of pre-immunization titers was considered as the point of seroconversion.

10

<sup>2</sup> the unit is defined as follows: the level of E1 antibodies in human chronic carriers prior to interferon therapy and infected with genotype 1b is < 0.1 U/ml for 50% of the patients; between 0.1 to 1 U/ml for 25% of the patients and > 1 U/ml in the remaining 25% of patients, n=58

**Example 12: Immunization of a chimpanzee chronically infected with HCV subtype 1b**

5 A chimpanzee (Phil) already infected for over 13 years (5015 days before immunization) with an HCV subtype 1b strain was vaccinated with E1 (aa 192-326) which was derived from a different strain of genotype 1b, with a 95.1% identity on the amino acid level (see also Table 2 of WO 99/67285 the whole of which is incorporated herein by reference), and which was prepared as described in examples 1-3 of WO 99/97285. The chimpanzee received in total 6 intramuscular  
10 immunizations of each 50 µg E1 in PBS/0.05% CHAPS mixed with RIBI R-730 (MPLA+TDM+CWS) according to the manufacturer's protocol (Ribi Inc. Hamilton, MT). The 6 immunizations were given in two series of three shots with a three week interval and with a lag period of 6 weeks between the two series. Starting 150 days prior to immunization, during the immunization period and until 1 year post immunization (but see below and WO 99/67285) the  
15 chimpanzee was continuously monitored for various parameters indicative for the activity of the HCV induced disease. These parameters included blood chemistry, ALT, AST, gammaGT, blood chemistry, viral load in the serum, viral load in the liver and liver histology. In addition, the immune answer to the immunization was monitored both on the humoral and cellular level. During this period the animal was also monitored for any adverse effects of the immunization, such as change in  
20 behaviour, clinical symptoms, body weight, temperature and local reactions (redness, swelling, indurations). Such effects were not detected.

Clearly, ALT (and especially gammaGT, data not shown) levels decreased as soon as the antibody level against E1 reached its maximum (see, Figure 8 of WO 99/67285). ALT rebounded rather  
25 rapidly as soon as the antibody levels started to decline, but gammaGT remained at a lower level as long as anti-E1 remained detectable.

E2 antigen in the liver decreased to almost undetectable levels during the period in which anti-E1 was detectable and the E2 antigen rebounded shortly after the disappearance of these antibodies.  
30 Together with the Core and E2 antigen becoming undetectable in the liver, the inflammation of the liver markedly decreased (see also Table 3 of WO 99/67285). This is a major proof that the vaccine induces a reduction of the liver damage, probably by clearing, at least partially, the viral antigens from its major target organ, the liver.

The viraemia level, as measured by Amplicor HCV Monitor (Roche, Basel, Switzerland), remained approximately unchanged in the serum during the whole study period.

More detailed analyses of the humoral response revealed that the maximum end-point titer reached  
5  $14.5 \times 10^3$  (after the sixth immunization) and that this titer dropped to undetectable 1 year post  
immunization (Figure 8 of WO 99/67285). Figure 9 of WO 99/67285 shows that the main epitopes,  
which can be mimicked by peptides, recognized by the B-cells are located at the N-terminal region of  
E2 (peptides V1V2 and V2V3, for details on the peptides used see Table 4 of WO 99/67285). Since  
the reactivity against the recombinant E1 is higher and longer lasting, it can also be deduced from  
10 this figure, that the antibodies recognizing these peptides represent only part of the total antibody  
population against E1. The remaining part is directed against epitopes which cannot be mimicked by  
peptides, i.e. discontinuous epitopes. Such epitopes are only present on the complete E1 molecule or  
even only on the particle-like structure. Such an immune response against E1 is unique, at least  
compared to what is normally observed in human chronic HCV carriers (WO 96/13590 to Maertens  
15 et al.) and in chimpanzees (van Doorn et al., 1996), who raise anti-E1 antibodies in their natural  
course of infection. In those patients, anti-E1 is in part also directed to discontinuous epitopes but a  
large proportion is directed against the C4 epitope ( $\pm 50\%$  of the patient sera), a minor proportion  
against V1V2 (ranging from 2-70% depending on the genotype), and reactivity against V2V3 was  
only exceptionally recorded (Maertens et al., 1997).

20 Analysis of the T-cell reactivity indicated that also this compartment of the immune system is  
stimulated by the vaccine in a specific way, as the stimulation index of these T-cells rises from 1 to  
2.5, and remains somewhat elevated during the follow up period (Figure 10 of WO 99/67285). It is  
this T cell reactivity that is only seen in Long term responders to interferon therapy (see: PCT/EP  
25 94/03555 to Leroux-Roels et al.; Leroux-Roels et al., 1996).

#### Example 13: Immunization of a chronic HCV carrier with different subtype

30 A chimpanzee (Ton) already infected for over 10 years (3809 days before immunization) with HCV  
from genotype 1a was vaccinated with E1 from genotype 1b, with only a 79.3 % identity on the  
amino acid level (see also Table 2 of WO 99/67285), and prepared as described in the previous  
examples. The chimpanzee received a total of 6 intramuscular immunizations of 50  $\mu$ g E1 in  
PBS/0.05% CHAPS each mixed with RIBI R-730 according to the manufacturer's protocol (Ribi

Inc. Hamilton, MT). The 6 immunizations were given in two series of three shots with a three week interval and with a lag period of 4 weeks between the two series. Starting 250 days prior to immunization, during the immunization period and until 9 months (but see below and WO 99/67285) post immunization the chimpanzee was continuously monitored for various parameters indicative for the activity of the HCV induced disease. These parameters included blood chemistry, ALT, AST, gammaGT, viral load in the serum, viral load in the liver and liver histology. In addition, the immune answer to the immunization was monitored both on the humoral and cellular level. During this period the animal was also monitored for any adverse effects of the immunization, such as change in behaviour, clinical symptoms, body weight, temperature and local reactions (redness, swelling, indurations). Such effects were not detected.

Clearly, ALT levels (and gammaGT levels, data not shown) decreased as soon as the antibody level against E1 reached its maximum (Figure 11 of WO 99/67285). ALT and gammaGT rebounded as soon as the antibody levels started to decline, but ALT and gammaGT remained at a lower level during the complete follow up period. ALT levels were even significantly reduced after vaccination ( $62 \pm 6$  U/l) as compared to the period before vaccination ( $85 \pm 11$  U/l). Since less markers of tissue damage were recovered in the serum, these findings were a first indication that the vaccination induced an improvement of the liver disease.

E2 antigen levels became undetectable in the period in which anti-E1 remained above a titer of  $1.0 \times 10^3$ , but became detectable again at the time of lower E1 antibody levels. Together with the disappearance of HCV antigens, the inflammation of the liver markedly decreased from moderate chronic active hepatitis to minimal forms of chronic persistent hepatitis (Table 3 of WO 99/67285). This is another major proof that the vaccine induces a reduction of the liver damage, probably by clearing, at least partially, the virus from its major target organ, the liver.

The viraemia level, as measured by Amplicor HCV Monitor (Roche, Basel, Switzerland), in the serum remained at approximately similar levels during the whole study period. More detailed analysis of the humoral response revealed that the maximum end-point titer reached was  $30 \times 10^3$  (after the sixth immunization) and that this titer dropped to  $0.5 \times 10^3$  nine months after immunization (Figure 11 of WO 99/67285). Figure 12 of WO 99/67285 shows that the main epitopes, which can be mimicked by peptides and are recognized by the B-cells, are located at the N-terminal region (peptides V1V2 and V2V3, for details on the peptides used see Table 4 of WO 99/67285). Since the reactivity against the recombinant E1 is higher and longer lasting, it can also be deduced from this figure, that the antibodies recognizing these peptides represent only part of the total antibody



population against E1. The remaining part is most likely directed against epitopes which cannot be mimicked by peptides, i.e. discontinuous epitopes. Such epitopes are probably only present on the complete E1 molecule or even only on the particle-like structure. Such an immune response against E1 is unique, at least compared to what is normally observed in human chronic HCV carriers, which have detectable anti-E1. In those patients, anti-E1 is in part also discontinuous, but a large proportion is directed against the C4 epitope (50% of the patient sera), a minor proportion against V1V2 (ranging from 2-70% depending on the genotype) and exceptionally reactivity against V2V3 was recorded (Maertens et al., 1997). As this chimpanzee is infected with an 1a isolate the antibody response was also evaluated for cross-reactivity towards a E1-1a antigen. As can be seen in Figure 13 of WO 99/67285, such cross-reactive antibodies are indeed generated, although, they form only part of the total antibody population. Remarkable is the correlation between the reappearance of viral antigen in the liver and the disappearance of detectable anti-1a E1 antibodies in the serum.

Analysis of the T-cell reactivity indicated that also this compartment of the immune system is stimulated by the vaccine in a specific way, as the stimulation index of these T-cells rises from 0.5 to 5, and remains elevated during the follow up period (Figure 14 of WO 99/67285).

#### Example 14: Reboosting of HCV chronic carriers with E1

As the E1 antibody titers as observed in examples 12 and 13 were not stable and declined over time, even to undetectable levels for the 1b infected chimp, it was investigated if this antibody response could be increased again by additional boosting. Both chimpanzees were immunized again with three consecutive intramuscular immunization with a three week interval (50 µg E1 mixed with RIBI adjuvant). As can be judged from Figures 8 and 11 of WO 99/67285, the anti-E1 response could indeed be boosted, once again the viral antigen in the liver decreased below detection limit. The viral load in the serum remained constant although in Ton (Figure 11 of WO 99/67285). A viremia level of  $< 10^5$  genome equivalents per ml was measured for the first time during the follow up period.

Notable is the finding that, as was already the case for the first series of immunizations, the chimpanzee infected with the subtype 1b HCV strain (Phil) responds with lower anti-E1 titers, than the chimpanzee infected with subtype 1a HCV strain (maximum titer in the first round  $14.5 \times 10^3$  versus  $30 \times 10^3$  for Ton and after additional boosting only  $1.2 \times 10^3$  for Phil versus  $40 \times 10^3$  for Ton). Although for both animals the beneficial effect seems to be similar, it could be concluded from

this experiment that immunization of a chronic carrier with an E1 protein derived from another subtype or genotype may be especially beneficial to reach higher titers, maybe circumventing a preexisting and specific immune suppression existing in the host and induced by the infecting subtype or genotype. Alternatively, the lower titers observed in the homologous setting (1b vaccine + 1b infection) may indicate binding of the bulk of the antibodies to virus. Therefore, the induced antibodies may possess neutralizing capacity.

**Example 15: Demonstration of prophylactic utility of E1-vaccination in chimpanzee**

10 The HCV E1s protein (amino acids 192-326) was expressed in Vero cells using recombinant vaccinia virus HCV11B. This vaccinia virus is essentially identical to vvHCV11A (as described in U.S. Patent No. 6,150,134, the entire contents of which is hereby incorporated by reference) but has been passaged from RK13 to Vero cells. The protein was purified (by means of lentil chromatography, reduction-alkylation and size exclusion chromatography) essentially as described in  
15 example 9 of PCT/E99/04342 (WO 99/67285), making use of iodoacetamide as alkylating agent for the cysteines. After purification the 3% empigen-BB was exchanged to 3% betain by size exclusion chromatography as described in example 1 of PCT/E99/04342 this process allows to recover E1s as a particle. Finally the material was desalted to PBS containing 0.5% betain and an E1s concentration of 500 µg/ml. This E1 was mixed with an equal volume of Alhydrogel 1.3% (Superfos, Denmark)  
20 and finally further diluted with 8 volumes of 0.9% NaCl to yield alum-adjuvanted E1 at a concentration of 50 µg E1/ml and 0.13% of Alhydrogel.

The HCV E2deltaHVRI (amino acids 412-715) was expressed in and purified from Vero essentially as described for E1 using recombinant vaccinia virus HCV101 which has been recombined from  
25 pvHCV-101 described in Example 8 of PCT/E99/04342 and wild type vaccinia virus. Also E2deltaHVRI behaves as a particle (measured by dynamic light scattering) after exchange of empigen to betain.

Five chimpanzees were selected which tested negative for HCV-RNA and HCV-antibodies. One of  
30 the animals (Huub) was not immunized, 2 animals received 6 immunizations with 50 µg E1 adjuvanted with alum (Marti and Yoran) while the remaining 2 animals received 6 immunizations with 50 µg E2deltaHVRI adjuvanted with alum (Joost and Karlien). All immunizations were administered intra-muscularly with a 3 week interval. Humoral and cellular immune responses were assessed in each animal against the antigen with which they where immunized and in each animal

both type of responses was detected as shown in Table 11.

Table 11: antibody titers were determined by ELISA two weeks after the 6<sup>th</sup> immunization. A serial dilution of the sample was compared to an in house standard (this in house standard defined as having 1000 mU/ml of E1 or anti-E2deltaHVR I antibody is a mixture of three sera from HCV chronic carriers selected based on a high anti-envelope titer). The stimulation index, which reflects the cellular immune response, was obtained by culturing PBMC, drawn from the animals two weeks after the third immunization, in the presence or absence of envelope antigen and determining the amount of tritiated thymidine incorporated in these cells during a pulse of 18 hours after 5 days of culture. The stimulation index is the ratio of thymidine incorporated in the cells cultured with envelope antigen versus the ones cultured without antigen. A stimulation index of >3 is considered a positive signal.

	Anti-E1 response		Anti-E2deltaHVRI response	
	Antibody titer	Stimulation index	Antibody titer	Stimulation index
Yoran	14110	10.9		
Marti	5630	14.2		
Joost			3210	8.5
Karlien			1770	11.2

Three weeks after the last of 6 immunizations all animals including the control were challenged with 100 CID (chimpanzee infectious doses) of a genotype 1b inoculum (J4.91, kindly provided by Dr. J. Bukh, NIH, Bethesda, Maryland). The amino acid sequence divergence between the vaccine proteins and the J4.91 isolate (of which the sequence information is available under accession number BAA01583) is 7% (9 out of 135 amino acids) for E1s and 11% (32 out of 304 amino acids) for E2delta HVRI; Consequently this challenge is considered heterologous and reflects a real life challenge.

All chimpanzees became HCV-RNA positive (determined with Monitor HCV, Roche, Basel, Switzerland) on day 7 post challenge and a first ALT and gammaGT peak was measured between days 35 and 63. This evidences that all chimps developed acute hepatitis. Remarkably, both E1 immunized animals resolved their infection while the E2deltaHVRI and the control animal did not. This is evidenced by the fact that the E1 immunized animals lost HCV-RNA (determined with

Monitor HCV, Roche, Basel, Switzerland) at day 98 (Yoran) and 133 (Marti) and remained negative so far until day 273 with monthly testing. All the other animals stayed RNA-positive during the entire follow up period of 273 days so far with ALT and gammaGT values not returning to normal as for the E1 immunized chimpanzees but gradually increasing.

5

In conclusion we have shown that E1-immunization changes the natural history of HCV infection by preventing evolution to a chronic infection, which is the major health problem related with HCV.

10 **Example 16: Similar E1 responses which allowed clearing of infection in chimpanzee can be induced in man**

In order to obtain a prophylactic effect of E1 immunization in man it is required that similar immune responses can be induced in man compared to chimpanzee. Therefore we vaccinated 20 male human  
15 volunteers, in which no anti-E1 responses (humoral or cellular) could be detected, with 3 doses of 20 µg E1s formulated on 0.13% Alhydrogel in 0.5 ml. All immunizations were given intramuscularly with a 3 week interval. As evidenced in Table 12, 17 out of 20 volunteers indeed mounted a significant humoral and cellular immune response against E1 and this without serious adverse events. Only 1 volunteer (subject 021) should be considered as a non-responder since neither  
20 humoral nor cellular responses were above the cut-off level after 3 E1 immunizations. The observation that the humoral response is lower compared to chimpanzee relates to the fact that only 3 immunizations with 20 µg were given and not 6 with 50 µg.

Table 12: antibody titers were determined by ELISA two weeks after the third immunization. A  
25 serial dilution of the sample was compared to an in house standards (this in house standard defined as having 1000 mU/ml of E1 or anti-E2deltaHVR I antibody is a mixture of three sera from HCV chronic carriers selected based on a high anti-envelope titer). The stimulation index (cellular immune response) was obtained by culturing PBMC, drawn from the individuals two weeks after the third immunization, in the presence or absence of 1 µg of E1s and determining the amount of tritiated  
30 thymidine incorporated in these cells during a pulse of 18 hours after 5 days of culture. The stimulation index is the ratio of thymidine incorporated in the cells cultured with envelope antigen versus the ones cultured without antigen. A stimulation index of >3 is considered a positive signal.

Subject no	Antibody titer	Stimulation index
------------	----------------	-------------------

002	1370	30.9
003	717	13.2
004	800	9.1
007	680	3.8
008	1026	3.9
009	325	4.6
010	898	7.7
011	284	4.1
012	181	3.6
013	<20	3.5
014	49	4.6
015	228	3.8
016	324	4.1
017	<20*	6.2
018	<20	6.7
019	624	3.1
020	84	5.5
021	<20	2.1
022	226	2.7
023	163	7.6

\* this individual is considered anti-E1 positive after immunization since a significant increase in ELISA signal was seen between the preimmune sample and the sample after three immunization, the titer however is very low and does not allow accurate determination.

Example 17: Boosting of E1 responses in vaccinated healthy volunteers

- 19 out of the 20 human volunteers of example 16 were boosted once more with 20 µg E1s formulated on 0.13% Alhydrogel in 0.5 ml at week 26 (i.e. 20 weeks after the third immunization). Again antibody titers and cellular immune responses were determined 2 weeks after this additional immunization. In all individuals the antibody titer had decreased during the 20 week interval but could easily be boosted by this additional immunization to a level equal or higher of that observed at week 8. On average the antibody titer was double as high after this boost compared to the week 8 titer, and 7 times as high compared to the week 26 titer (Table 13).
- Table 13: antibody titers were determined by ELISA two weeks (= week 8) and 20 weeks (= week 26) after the third immunization and finally also 2 weeks after the boost (= Week 28). A serial dilution of the sample was compared to an in house standards (this in house standard defined as having 1000 mU/ml of E1 antibody, is a mixture of three sera from HCV chronic carriers selected based on a high anti-envelope titer). For accurate comparison the determination of the titer at week 8 was repeated within the same assay as for the week 26 and 28 samples, which explains the differences with table 12 of example 16.

Subject no	Antibody titer		
	Week 8	Week 26	Week 28
002	1471	443	3119
003	963	95	2355
004	1006	409	2043
007	630	65	541
008	926	81	819
009	704	77	269
010	1296	657	3773
011	253	65	368
012	254	148	760
013	36	<20	166
014	53	40	123
016	159	45	231
017	109	39	568
018	43	23	50

019	425	157	1894
020	73	33	113
021	25	<20	26
022	280	150	357
024	177	81	184
average	467	138	936

Remarkably the T-cell responses were for the majority of individuals still high after the 20 week interval. Taking in account a normalization to the tetanos response, which is present in most individuals as a consequence of previous vaccinations, there is no change in the geometric mean of the stimulation index. After the additional boost, taking in account a normalization to the tetanos response, no change is noted (figure 51). This confirms that a strong T-help response was induced after 3 E1 immunizations and indicates that these immunizations induced already a very good T-help memory which requires, at least for a period of 6 months, no further boosting.

- 10 Legend to figure 51: The stimulation index (cellular immune response) was obtained by culturing PBMC ( $10^5$  cells), drawn from the individuals before immunization (week 0), two weeks after the third immunization (week 8), before the booster immunization (week 26) and two weeks after the booster immunization (week 28), in the presence or absence of 3  $\mu$ g of recombinant E1s or 2  $\mu$ g tetanos toxoid and determining the amount of tritiated thymidine incorporated in these cells during a pulse of 18 hours after 5 days of culture. The stimulation index is
- 15 the ratio of thymidine incorporated in the cells cultured with envelope antigen versus the ones cultured without antigen. Samples of week 0 and 8 were determined in a first assay (A), while the samples of week 26 and 28 were determined in a second assay (B) in which the samples of week 0 were reanalyzed. Results are expressed as the geometric mean stimulation index of all 20 (A, experiment) or 19 (B, experiment) volunteers.
- 20 In addition the Th1 cytokine interferon-gamma and Th2 cytokine interleukin-5 were measured in the supernatants of the PBMC cultures of samples taken at week 26 and 28 and restimulated with E1. As can be judged from figure 52 the predominant cytokine secreted by the E1 stimulated PBMC is interferon-gamma. It is highly surprising to see that a strong Th1 biased response is observed with an alum adjuvanted E1, since alum is known to be a Th2 inducer. Once more the results confirm that a good T-cell memory response is induced, as prior to
- 25 the final boost (week 26) already a very strong response is observed. The interferon-gamma secretion was found to be specific as in an additional experiment we saw no difference in interferon-gamma secretion between E1 stimulated cell cultures and non-stimulated cell cultures of these volunteers using samples drawn at week 0.

Legend to figure 52: PBMC ( $10^6$  cells), drawn from the individuals before the booster immunization (week 26) and two weeks after the booster immunization (week 28), were cultured in the presence of 3  $\mu$ g of recombinant E1s (E1) or 2  $\mu$ g of tetanus toxoid (TT) or no antigen (BI). Cytokines were measured in the supernatant taken after 24 hours (interleukin-5) or after 120 hours (interferon- $\gamma$ ) by means of ELISA. The stimulation index is the ratio of cytokine measured in the supernatants of cells cultured with envelope antigen versus the ones cultured without antigen. Results are expressed as the geometric mean of pg cytokine/ml secreted of all 19 volunteers. Samples with a cytokine amount below detection limit were assigned the value of the detection limit. Similarly samples with extremely high concentrations of cytokine out of the linear range of the assay were assigned the value of the limit of the linear range of the assay.

Example 18: Fine mapping of cellular response against E1 in vaccinated healthy volunteers.

In order to map the E1 specific responses a series of 20-mer peptides was synthesized, using standard Fmoc chemistry, with 8 amino acids overlap and covering the entire sequence of E1s. All peptides were C-terminally amidated and N-terminally acetylated, with the exception of IGP 1626 which has a free amino-terminus.

IGP 1626	YEVARNVSGIYHVTNDCSNSS (amino acid 192-211)
IGP 1627	TNDCSNSSIVYEAADMIMHT (amino acid 204-223)
IGP 1628	AADMIMHTPGCVPCVRENNS (amino acid 216-235)
IGP 1629	PCVRENNSRCWVALTPTLA (amino acid 228-247)
IGP 1630	VALTPTLAARNASVPTTIR (amino acid 240-259)
IGP 1631	SVPTTIRRHVDLLVGAAAF (amino acid 252-271)
IGP 1632	LLVGAAAFCSAMYVGDLGGS (amino acid 264-283)
IGP 1633	YVGDLGGSVFLVSQLFTISP (amino acid 276-295)
IGP 1634	SQLFTISPRRHETVQDCNCS (amino acid 288-307)
IGP 1635	TVQDCNCSIYPGHITGHRMA (amino acid 300-319)
IGP 1636	HITGHRMAWDMMNWSPTTA (amino acid 312-331)

PBMC from 14 different healthy donors not vaccinated with E1s or 10 donors vaccinated with E1s were cultured in the presence of 25  $\mu$ g/ml (non vaccinated persons) or 10  $\mu$ g/ml (vaccinated persons, samples taken after the third or booster injection) of each peptide separately. As can be judged from figure 53 the peptides IGP 1627, 1629, 1630, 1631, 1633, 1635 and 1636 all induced significantly higher responses in vaccinated persons



compared to non-vaccinated persons. Using a stimulation index of 3 as cut-off the peptides IGP 1627, 1629, 1631 and 1635 were the most frequently recognized (i.e. recognized by at least half of the vaccinated persons tested). This experiment proves that the T-cell responses induced by E1s derived from mammalian cell culture are specific against E1 since these responses can not only be recalled by the same E1s derived from mammalian cell culture but also by synthetic peptides. In addition this experiment delineates the most immunogenic T-cell domains in E1 are located between amino acids 204-223, 228-271, 276-295, 300-331 and more particularly even between amino acids 204-223, 228-247, 252-271 and 300-319.

Legend to figure 53: The stimulation index (cellular immune response) was obtained by culturing PBMC ( $3 \times 10^5$  cells), in the presence or absence of peptides and determining the amount of tritiated thymidine incorporated in these cells during a pulse after 5-6 days of culture. The stimulation index is the ratio of thymidine incorporated in the cells cultured with peptide versus the ones cultured without peptide. Results are expressed as individual values for vaccinated persons (top panel) or non vaccinated or controls (lower panel).

The present invention also provides therefor, the following E1 peptides, proteins, compositions and kits containing the same, nucleic acid sequences coding for these peptides and proteins containing the same, and methods of their manufacture and use, as are generally described herein for other E1 and related peptides of the present invention.

	IGP 1626	spanning positions 192-211 of the E1 region (SEQ ID NO:112),
20	IGP 1627	spanning positions 204-223 of the E1 region (SEQ ID NO:113),
	IGP 1628	spanning positions 216-235 of the E1 region (SEQ ID NO:114),
	IGP 1629	spanning positions 228-247 of the E1 region (SEQ ID NO:115),
	IGP 1630	spanning positions 240-259 of the E1 region (SEQ ID NO:116),
	IGP 1631	spanning positions 252-271 of the E1 region (SEQ ID NO:117),
25	IGP 1632	spanning positions 264-283 of the E1 region (SEQ ID NO:118),
	IGP 1633	spanning positions 276-295 of the E1 region (SEQ ID NO:119),
	IGP 1634	spanning positions 288-307 of the E1 region (SEQ ID NO:120),
	IGP 1635	spanning positions 300-319 of the E1 region (SEQ ID NO:121),
30	IGP 1636	spanning positions 312-331 of the E1 region (SEQ ID NO:122).

## REFERENCES

- Bailey, J. and Cole, R. (1959) *J. Biol. Chem.* 234, 1733-1739.
- Ballou, L., Hitzeman, R., Lewis, M. & Ballou, C. (1991) *PNAS* 88, 3209-3212.
- Benesch, R., Benesch, R.E., Gutcho, M. & Lanfer, L. (1956) *Science* 123, 981.
- Cavins, J. & Friedman. (1970) *Anal. Biochem.* 35, 489.
- Cleland, W. (1964) *Biochemistry* 3, 480
- Creighton, E. (1988) *BioEssays* 8, 57
- Darbre, A., John Wiley & Sons Ltd. (1987) *Practical Protein Chemistry - A Handbook*.
- Darbre, A., John Wiley & Sons Ltd. (1987) *Practical Proteinchemistry* p.69-79.
- Doms et al, (1993), *Virology* 193, 545-562.
- Ellman, G. (1959) *Arch. Biochem. Biohys.* 82, 70.
- Falkner, F. & Moss, B. (1988) *J. Virol.* 62, 1849-1854.
- Friedman, M. & Krull. (1969) *Biochem. Biophys. Res. Commun.* 37, 630.
- Gallagher J. (1988) *J. Cell Biol.* 107, 2059-2073.
- Glazer, A., Delange, R., Sigman, D. (1975) North Holland publishing company, Elsevier, Biomedical. Part :  
Modification of protein (p. 116).
- Graham, F. & van der Eb, A. (1973) *Virology* 52, 456-467.
- Grakoui et al. (1993) *Journal of Virology* 67:1385-1395.

Grassetti, D. & Murray, J. (1969) *Analyt. Chim. Acta.* 46, 139.

Grassetti, D. & Murray, J. (1967) *Arch. Biochem Biophys.* 119, 41.

Helenius, *Mol. Biol. Cell* (1994), 5: 253-265.

Hijikata, M., Kato, N., Ootsuyama, Y., Nakagawa, M. & Shimotohno, K. (1991) *Proc. Natl. Acad. Sci. U.S.A.* 88(13):5547-51.

Hochuli, E., Bannwarth, W., Döbeli, H., Gentz, R., Stüber, D. (1988) *Biochemistry* 88, 8976.

Hsu, H., Donets, M., Greenberg, H. & Feinstone, S. (1993) *Hepatology* 17:763-771.

Inoue, Y., Suzuki, R., Matsuura, Y., Harada, S., Chiba, J., Watanabe, Y., Saito, I. & Miyamura, T. (1992) *J. Gen. Virol.* 73:2151-2154.

Janknecht, R., de Martynoff, G. *et al.*, (1991) *Proc. Natl. Acad. Sci. USA* 88, 8972-8976.

Kayman (1991) *J. Virology* 65, 5323-5332.

Kato, N., Oostuyama, Y., Tanaka, T., Nakagawa, M., Muraiso, K., Ohkoshi, S., Hijikata, M., Shimotohno, K. (1992) *Virus Res.* 22:107-123.

Kniskern, P., Hagopian, A., Burke, P., Schultz, L., Montgomery, D., Humi, W., Yu Ip, C., Schulman, C., Malgetter, R., Wampler, D., Kubek, D., Sitrin, R., West, D., Ellis, R., Miller, W. (1994) *Vaccine* 12:1021-1025.

Kohara, M., Tsukiyama-Kohara, K., Maki, N., Asano, K., Yoshizawa, K., Miki, K., Tanaka, S., Hattori, N., Matsuura, Y., Saito, I., Miyamura, T. & Nomoto, A. (1992) *J. Gen. Virol.* 73:2313-2318.

Mackett, M., Smith, G. & Moss, B. (1985) In: 'DNA cloning: a practical approach' (Ed. Glover, D.) IRL Press, Oxford.

Mackett, M., & Smith, G. (1986) J. Gen. Virol. 67, 2067-2082.

Mackett, M., Smith, G. & Moss, B. (1984) J. Virol. 49, 857-864.

Mackett, M., Smith, G. & Moss, B. (1984) Proc. Natl. Acad. Sci. USA 79, 7415-7419.

Means, G. (1971) Holden Day, Inc.

Means, G. & Feeney, R. (1971) Holden Day p.105 & p. 217.

Mita, E., Hayashi, N., Ueda, K., Kasahara, A., Fusamoto, H., Takamizawa, A., Matsubara, K., Okayama, H. & Kamada T. (1992) Biochem. Biophys. Res. Comm. 183:925-930.

Moore, S. (1963) J. Biol. Chem. 238, 235-237.

Okamoto, H., Okada, S., Sugiyama, Y., Yotsumoto, S., Tanaka, T., Yoshizawa, H., Tsuda, F., Miyakawa, Y. & Mayumi, M. (1990) Jpn. J. Exp. Med. 60:167-177.

Panicali & Paoletti (1982) Proc. Natl. Acad. Sci. USA 79, 4927-4931.

Piccini, A., Perkus, M. & Paoletti, E. (1987) Meth. Enzymol. 153, 545-563.

Rose (1988) Annu. Rev. Cell Biol. 1988, 4: 257-288;

Ruegg, V. and Rudinger, J. (1977) Methods Enzymol. 47, 111-116.

Shan, S. & Wong (1993) CRC-press p. 30-33.

Spaete, R., Alexander, D., Rugroden, M., Choo, Q., Berger, K., Crawford, K., Kuo, C., Leng, S., Lee, C., Ralston, R., et al. (1992) Virology 188(2):819-30.

Skehel, J., (1984) Proc. Natl. Acad. Sci. USA 81, 1179-1783.

Stunnenberg, H., Lange, H., Philipson, L., Miltenburg, R. & van der Vliet, R. (1988) Nucl. Acids Res. 16, 2431-2444.

Stuyver, L., Van Arnhem, W., Wyseur, A., DeLeys, R. & Maertens, G. (1993a) Biochem. Biophys. Res. Commun. 192, 635-641.

Stuyver, L., Rossau, R., Wyseur, A., Duhamel, M., Vanderborght, B., Van Heuverswyn, H., & Maertens, G. (1993b) J. Gen. Virol. 74, 1093-1102.

Stuyver, L., Van Arnhem, W., Wyseur, A., Hernandez, F., Delaporte, E., Maertens, G. (1994), Proc. Natl. Acad. Sci. USA 91:10134-10138.

Weil, L. & Seibler, S. (1961) Arch. Biochem. Biophys. 95, 470.

Yokosuka, O., Ito, Y., Imazeki, F., Ohto, M. & Omata, M. (1992) Biochem. Biophys. Res. Commun. 189:565-571.

Miller P, Yano J, Yano E, Carroll C, Jayaram K, Ts'o P (1979) Biochemistry 18:5134-43.

Nielsen P, Egholm M, Berg R, Buchardt O (1991) Science 254:1497-500.

Nielsen P, Egholm M, Berg R, Buchardt O (1993) Nucleic-Acids-Res. 21:197-200.

Asseline U, Delarue M, Lancelot G, Toulme F, Thuong N (1984) Proc. Natl. Acad. Sci. USA 81:3297-301.

Matsukura M, Shinozuka K, Zon G, Mitsuya H, Reitz M, Cohen J, Broder S (1987) Proc. Natl. Acad. Sci. USA 84:7706-10.

WO 96/04385 (PCT/EP95/03031) - Purified Hepatitis C Virus Envelope Proteins for Diagnostic and Therapeutic Use.

All references cited herein are incorporated in their entirety by reference.

**We claim:**

1. A therapeutic vaccine composition comprising a therapeutic effective amount of:  
a composition comprising at least one purified recombinant HCV single or specific oligomeric recombinant envelope proteins selected from the group consisting of an E1 protein and an E2 protein; and optionally a pharmaceutically acceptable adjuvant.
2. A composition according to claim 1 wherein said recombinant HCV envelope proteins are produced by recombinant mammalian cells.
3. A composition according to claim 1 wherein said recombinant HCV envelope proteins are produced by recombinant yeast cells.
4. A therapeutic vaccine composition comprising a therapeutically effective amount of a composition comprising at least one of the following E1 and E2 peptides:
  - E1-31 (SEQ ID NO:56) spanning amino acids 181 to 200 of the Core/E1 V1 region,
  - E1-33 (SEQ ID NO:57) spanning amino acids 193 to 212 of the E1 region,
  - E1-35 (SEQ ID NO:58) spanning amino acids 205 to 224 of the E1 V2 region (epitope B),
  - E1-35A (SEQ ID NO:59) spanning amino acids 208 to 227 of the E1 V2 region (epitope B),
  - 1bE1 (SEQ ID NO:53) spanning amino acids 192 to 228 of E1 regions V1, C1, and V2 regions (containing epitope B),
  - E1-51 (SEQ ID NO:66) spanning amino acids 301 to 320 of the E1 region,
  - E1-53 (SEQ ID NO:67) spanning amino acids 313 to 332 of the E1 C4 region (epitope A),
  - E1-55 (SEQ ID NO:68) spanning amino acids 325 to 344 of the E1 region,
  - Env 67 or E2-67 (SEQ ID NO:72) spanning amino acid positions 397 to 418 of the E2 region (epitope A),
  - Env 69 or E2-69 (SEQ ID NO:73) spanning amino acid positions 409 to 428 of the E2 region (epitope A),
  - Env 23 or E2-23 (SEQ ID NO:86) spanning positions 583 to 602 of the E2 region (epitope E),
  - Env 25 or E2-25 (SEQ ID NO:87) spanning positions 595 to 614 of the E2 region (epitope E),
  - Env 27 or E2-27 (SEQ ID NO:88) spanning positions 607 to 626 of the E2 region (epitope E),
  - Env 178 or E2-178 (SEQ ID NO:83) spanning positions 547 to 586 of the E2 region (epitope D),
  - Env 13B or E2-13B (SEQ ID NO:82) spanning positions 523 to 542 of the E2 region (epitope C),
  - IGP 1626 spanning positions 192-211 of the E1 region (SEQ ID NO:112),
  - IGP 1627 spanning positions 204-223 of the E1 region (SEQ ID NO:113),
  - IGP 1628 spanning positions 216-235 of the E1 region (SEQ ID NO:114),
  - IGP 1629 spanning positions 228-247 of the E1 region (SEQ ID NO:115),
  - IGP 1630 spanning positions 240-259 of the E1 region (SEQ ID NO:116),

IGP 1631 spanning positions 252-271 of the E1 region (SEQ ID NO:117),  
 IGP 1632 spanning positions 264-283 of the E1 region (SEQ ID NO:118),  
 IGP 1633 spanning positions 276-295 of the E1 region (SEQ ID NO:119),  
 IGP 1634 spanning positions 288-307 of the E1 region (SEQ ID NO:120),  
 IGP 1635 spanning positions 300-319 of the E1 region (SEQ ID NO:121) and  
 IGP 1636 spanning positions 312-331 of the E1 region (SEQ ID NO:122).

5. A method of treating a mammal infected with HCV comprising administering an effective amount of a composition according to any one of claims 1-4 and, optionally, a pharmaceutically acceptable adjuvant.

6. The method of claim 5 wherein said mammal is a human.

7. A composition comprising at least one purified recombinant HCV recombinant envelope proteins selected from the group consisting of an E1 protein and an E2 protein, and optionally an adjuvant.

8. A composition comprising at least one of the following E1 and E2 peptides:

E1-31 (SEQ ID NO:56) spanning amino acids 181 to 200 of the Core/E1 V1 region,  
 E1-33 (SEQ ID NO:57) spanning amino acids 193 to 212 of the E1 region,  
 E1-35 (SEQ ID NO:58) spanning amino acids 205 to 224 of the E1 V2 region (epitope B),  
 E1-35A (SEQ ID NO:59) spanning amino acids 208 to 227 of the E1 V2 region (epitope B),  
 1bE1 (SEQ ID NO:53) spanning amino acids 192 to 228 of E1 regions V1, C1, and V2 regions (containing epitope B),  
 E1-51 (SEQ ID NO:66) spanning amino acids 301 to 320 of the E1 region,  
 E1-53 (SEQ ID NO:67) spanning amino acids 313 to 332 of the E1 C4 region (epitope A),  
 E1-55 (SEQ ID NO:68) spanning amino acids 325 to 344 of the E1 region,  
 Env 67 or E2-67 (SEQ ID NO:72) spanning amino acid positions 397 to 418 of the E2 region (epitope A),  
 Env 69 or E2-69 (SEQ ID NO:73) spanning amino acid positions 409 to 428 of the E2 region (epitope A),  
 Env 23 or E2-23 (SEQ ID NO:86) spanning positions 583 to 602 of the E2 region (epitope E),  
 Env 25 or E2-25 (SEQ ID NO:87) spanning positions 595 to 614 of the E2 region (epitope E),  
 Env 27 or E2-27 (SEQ ID NO:88) spanning positions 607 to 626 of the E2 region (epitope E),  
 Env 178 or E2-178 (SEQ ID NO:83) spanning positions 547 to 586 of the E2 region (epitope D),  
 Env 13B or E2-13B (SEQ ID NO:82) spanning positions 523 to 542 of the E2 region (epitope C),  
 IGP 1626 spanning positions 192-211 of the E1 region (SEQ ID NO:112),  
 IGP 1627 spanning positions 204-223 of the E1 region (SEQ ID NO:113),  
 IGP 1628 spanning positions 216-235 of the E1 region (SEQ ID NO:114),  
 IGP 1629 spanning positions 228-247 of the E1 region (SEQ ID NO:115),  
 IGP 1630 spanning positions 240-259 of the E1 region (SEQ ID NO:116),

IGP 1631 spanning positions 252-271 of the E1 region (SEQ ID NO:117),  
IGP 1632 spanning positions 264-283 of the E1 region (SEQ ID NO:118),  
IGP 1633 spanning positions 276-295 of the E1 region (SEQ ID NO:119),  
IGP 1634 spanning positions 288-307 of the E1 region (SEQ ID NO:120),  
IGP 1635 spanning positions 300-319 of the E1 region (SEQ ID NO:121) and  
IGP 1636 spanning positions 312-331 of the E1 region (SEQ ID NO:122).

9. A therapeutic composition for inducing HCV-specific antibodies comprising a therapeutic effective amount of a composition comprising an E1/E2 complex formed from purified recombinant HCV single or specific oligomeric recombinant E1 or E2 proteins; and optionally a pharmaceutically acceptable adjuvant.

10. A composition according to claim 9 wherein said recombinant HCV envelope proteins are produced by recombinant mammalian cells.

11. A composition according to claim 9 wherein said recombinant HCV envelope proteins are produced by recombinant yeast cells.

12. A method of treating a mammal infected with HCV comprising administering an effective amount of a composition according to any one of claims 9-11 and, optionally, a pharmaceutically acceptable adjuvant.

13. The method of claim 12 wherein said mammal is a human.

14. A therapeutic composition for inducing HCV-specific antibodies comprising a therapeutic effective amount of a composition comprising at least one purified recombinant HCV single or specific oligomeric recombinant envelope protein selected from the group consisting of an E1 protein and an E2 protein; and optionally a pharmaceutically acceptable adjuvant.



1/65




Fig. 1

2/65




Fig. 2

3/65




Fig. 3

4/65




Fig. 4

Anti-E1 levels in NON-responders to IFN treatment




Fig. 5

6/65

# Anti-E1 levels in RESPONDERS to IFN treatment




Fig. 6

7/65

Anti-E1 levels in patients with COMPLETE response to IFN

SERIES 2





Fig. 7

8/65

Anti-E1 levels in INCOMPLETE responders to IFN treatment

SERIES 2



months after start of treatment


Fig. 8



9/65

# Anti-E2 levels in NON-RESPONDERS to IFN treatment

SERIES 1




weeks after start of treatment

Fig. 9

10/65


# Anti-E2 levels in RESPONDERS to IFN treatment



11/65

Anti-E2 levels in INCOMPLETE responders to IFN treatment

SERIES 2



months after start of treatment

Fig.11

12/65




Fig.12

Human anti-E1 reactivity competed with peptides




Fig.13

14/65

# Competition of reactivity of anti-E1 Mabs with peptides




Fig.14

Anti-E1 (epitope 1) levels in NON-RESPONDERS to IFN treatment

SERIES 1





Fig.15





17/65

# Anti-E1 (epitope 2) levels in NON-RESPONDERS to IFN treatment

SERIES 1




Fig.17

18/65

Anti-E1 (epitope 2) levels in RESPONDERS to IFN treatment




Fig.18

Competition of reactivity of anti-E2 Mabs with peptides




Fig.19

# Human anti-E2 reactivity competed with peptides




Fig. 20

21/65

## Fig. 21A

5' GGCATGCAAGCTTAATTAATT3' (SEQ ID NO 1)  
3'ACGTCCGTACGTTCTGAATTAATTAATCGA5' (SEQ ID NO 94)

5'CCGGGGAGGCCTGCACGTGATCGAGGGCAGACACCATCACCACCATCACTAATAGT  
TAATTAAGTCA 3' (SEQ ID NO 2)  
3'CCTCCGGACGTGCACTAGCTCCCGTCTGTGGTAGTGGTGGTAGTGATTATCAATTAATTG  
5' (SEQ ID NO 95)

SEQ ID NO 3 (HCC19A)

ATGCCCGGTTGCTCTTTCTCTATCTTCTCTTGGCTTTACTGTCCTGTCTGACCATTC  
GCTTCCGCTTATGAGGTGCGCAACGTGTCCGGGATGTACCATGTCACGAACGACTGCT  
CCAACCTCAAGCATTGTGTATGAGGCAGCGGACATGATCATGCACACCCCCGGGTGCGT  
GCCCTGCGTTCGGGAGAACAACTCTTCCCGCTGCTGGGTAGCGCTACCCCCACGCTC  
GCAGCTAGGAACGCCAGCGTCCCCACCGACAATACGACGCCACGTGATTTGCTCG  
TTGGGGCGGCTGCTCTCTGTTCCGCTATGTACGTGGGGGATCTCTGCGGATCTGTCTTC  
CTCGTCTCCAGCTGTTACCATCTCGCCTCGCCGGCATGAGACGGTGCAGGACTGCA  
ATTGCTCAATCTATCCCGGCCACATAACAGGTCACCGTATGGCTTGGGATATGATGAT  
GAACTGGTCGCCTACAACGGCCCTGGTGGTATCGCAGCTGCTCCGGATCCCACAAGCT  
GTCGTGGACATGGTGGCGGGGGCCCATTTGGGGAGTCCTGGCGGGCCTCGCCTACTATT  
CCATGGTGGGGAACCTGGGCTAAGGTTTTGATTGTGATGCTACTCTTGCTCTCTAATAG

SEQ ID NO 5 (HCC10A)

ATGTTGGGTAAGGTCATCGATACCTTACATGCGGCTTCGCCGACCTCGTGGGGTACA  
TTCCGCTCGTCGGCGCCCCCTAGGGGGCGCTGCCAGGGCCCTGGCGCATGGCGTCCG  
GGTCTGGAGGACGGCGTGAACATGCAACAGGGAATTTGCCCGGTTGCTCTTTCTCT  
ATCTTCTCTTGGCTTTGCTGTCCTGTCTGACCGTTCCAGCTTCCGCTTATGAAGTGCG  
CAACGTGTCCGGGATGTACCATGTCACGAACGACTGCTCCAACCTCAAGCATTGTGTAT  
GAGGCAGCGGACATGATCATGCACACCCCCGGGTGCGTGCCCTGCGTTCGGGAGAAC  
AACTCTTCCCGCTGCTGGGTAGCGCTACCCCCACGCTCGCAGCTAGGAACGCCAGCG  
TCCCCACCGACAATACGACGCCACGTGATTTGCTCGTTGGGGCGGCTGCTTTCTG

22/65

## Fig. 21B

TTCCGCTATGTACGTGGGGACCTCTGCGGATCTGTCTTCCTCGTCTCCCAGCTGTTCA  
CCATCTCGCCTCGCCGGCATGAGACGGTGCAGGACTGCAATTGCTCAATCTATCCCGG  
CCACATAACGGGTCACCGTATGGCTTGGGATATGATGATGAACTGGTCGCCTACAACG  
GCCCTGGTGGTATCGCAGCTGCTCCGGATCCCACAAGCTGTCGTGGACATGGTGGCGG  
GGGCCATTGGGGAGTCTGCGGGTCTCGCCTACTATTCCATGGTGGGGAAGTGGGC  
TAAGGTTTTGATTGTGATGCTACTCTTTGCTCCCTAATAG

SEQ ID NO 7 (HCC111A)

ATGTTGGGTAAGGTCATCGATACCTTACGTGCGGCTTCGCCGACCTCATGGGGTACA  
TTCCGCTCGTGGCGCCCCCTAGGGGGTGTGCCAGAGCCCTGGCGCATGGCGTCCG  
GGTCTGGAAGACGGCGTGAACATGCAACAGGGAATTTGCCTGGTTGCTCTTTCTCTA  
TCTTCTCTTGGCTTTACTGTCTGTCTGACCATTCAGCTTCGGCTTATGAGGTGCGC  
AACGTGTCGGGGATGTACCATGTCACGAACGACTGCTCCAACCAAGCATTGTGTATG  
AGGCAGCGGACATGATCATGCACACCCCCGGGTGCGTGCCCTGCGTTCGGGAGAACA  
ACTCTTCCCGCTGCTGGGTAGCGCTCACCCCCACGCTCGCAGCTAGGAACGCCAGCGT  
CCCCACTACGACAATACGACGCCACGTCGATTTGCTCGTTGGGGCGGCTGCTTTCTGTT  
CCGCTATGTACGTGGGGGATCTCTGCGGATCTGTCTTCCTCGTCTCCCAGCTGTTACC  
ATCTCGCCTCGCCGGCATGAGACGGTGCAGGACTGCAATTGCTCAATCTATCCCGGCC  
ACATAACAGGTCACCGTATGGCTTGGGATATGATGATGAACTGGTAATAG

SEQ ID NO 9 (HCC112A)

ATGCCCGGTTGCTCTTTCTCTATCTTCTCTTGGCCCTGCTGTCTGTCTGACCATACCA  
GCTTCCGCTTATGAAGTGCACAACGTGTCCGGGGTGTACCATGTCACGAACGACTGCT  
CCAACCTCAAGCATAGTGTATGAGGCAGCGGACATGATCATGCACACCCCCGGGTGCGT  
GCCCTGCGTTCGGGAGGGCAACTCCTCCCGTTGCTGGGTGGCGCTCACTCCACGCTC  
GCGGCCAGGAACGCCAGCGTCCCCACAACGACAATACGACGCCACGTCGATTTGCTC  
GTTGGGGCTGCTGCTTTCTGTTCCGCTATGTACGTGGGGGATCTCTGCGGATCTGTTTT  
CCTTGTTTCCAGCTGTTACCTTCTACCTCGCCGGCATCAAACAGTACAGGACTGCA  
ACTGCTCAATCTATCCCGGCCATGTATCAGGTCACCGCATGGCTTGGGATATGATGAT  
GAACTGGTCCTAATAG

SEQ ID NO 11 (HCC113A)

ATGTCCGGTTGCTCTTTCTCTATCTTCTCTTGGCCCTGCTGTCTGTCTGACCATACCA  
GCTTCCGCTTATGAAGTGCACAACGTGTCCGGGGTGTACCATGTCACGAACGACTGCT  
CCAACCTCAAGCATAGTGTATGAGGCAGCGGACATGATCATGCACACCCCCGGGTGCGT

23/65

## Fig. 21C

GCCCTGCGTTTCGGGAGGGCAACTCCTCCCGTTGCTGGGTGGCGCTCACTCCACGCTC  
GCGGCCAGGAACGCCAGCGTCCCCACAACGACAATACGACGCCACGTGCGATTTGCTC  
GTTGGGGCTGCTGCTTTCTGTTCCGCTATGTACGTGGGGGATCTCTGCGGATCTGTTTT  
CCTTGTTTCCCAGCTGTTACCTTCTCACCTCGCCGGCATCAAACAGTACAGGACTGCA  
ACTGCTCAATCTATCCCGGCCATGTATCAGGTCACCGCATGGCTTGGGATATGATGAT  
GAACTGGTAATAG

SEQ ID NO 13 (HCCI17A)

ATGCTGGGTAAGGCCATCGATACCCTTACGTGCGGCTTCGCCGACCTCGTGGGGTACA  
TTCGCTCGTCGGCGCCCCCTAGGGGGCGCTGCCAGGGCCCTGGCGCATGCGCTCCG  
GGTTCTGGAAGACGGCGTGAACATGCAACAGGGAATTTGCCTGGTTGCTCTTCTCTA  
TCTTCCTCTTGGCTTTACTGTCTGTCTAACCATTCCAGCTTCCGCTTACGAGGTGCGC  
AACGTGTCCGGGATGTACCATGTACGAACGACTGCTCCAACCAAGCATTGTGTATG  
AGGCAGCGGACATGATCATGCACACCCCGGGTGCGTGCCCTGCGTTGGGAGAACA  
ACTCTTCCCGCTGCTGGGTAGCGCTACCCCCACGCTCGCGGCTAGGAACGCCAGCAT  
CCCCACTACAACAATACGACGCCACGTGCGATTTGCTCGTTGGGGCGGCTGCTTTCTGTT  
CCGCTATGTACGTGGGGGATCTCTGCGGATCTGTCTTCTCGTCTCCAGCTGTTACCC  
ATCTCGCTCGCCGGCATGAGACGGTGACGACTGCAATTGCTCAATCTATCCCGGCC  
ACATAACGGGTCACCGTATGGCTTGGGATATGATGATGAACTGGTACTAATAG

SEQ ID NO 15 (HCP#51)

ATGCCCGGTTGCTCTTCTCTATCTT

SEQ ID NO 16 (HCP#52)

ATGTTGGGTAAGGTCATCGATACCCT

SEQ ID NO 17 (HCP#53)

CTATTAGGACCAGTTCATCATCATATCCCA

SEQ ID NO 18 (HCP#54)

CTATTACCAGTTCATCATCATATCCCA

SEQ ID NO 19 (HCP#107)

ATACGACGCCACGTGCGATTCAGCTGTTACCATC

24/65

## Fig. 21D

SEQ ID NO 20 (HCP108)

GATGGTGAACAGCTGGGAATCGACGTGGCGTCGTAT

SEQ ID NO 21 (HCC137)

ATGTTGGGTAAGGTCATCGATACCCCTTACATGCGGCTTCGCCGACCTCGTGGGGTACA  
TTCCGCTCGTCGGCGCCCCCTAGGGGGCGCTGCCAGGGCCCTGGCGCATGGCGTCCG  
GGTTCTGGAGGACGGCGTGAACATATGCAACAGGGAATTTGCCCGGTTGCTCTTTCTCT  
ATCTTCCTCTTGCTTTGCTGTCTGTCTGACCGTTCCAGCTTCCGCTTATGAAGTGCG  
CAACGTGTCCGGGATGTACCATGTACGAACGACTGCTCCAACCTCAAGCATTGTGTAT  
GAGGCAGCGGACATGATCATGCACACCCCCGGGTGCGTGCCCTGCGTTCCGGGAGAAC  
AACTCTTCCCGCTGCTGGGTAGCGCTCACCCCCACGCTCGCAGCTAGGAACGCCAGCG  
TCCCCACCACGACAATACGACGCCACGTCGATTCCAGCTGTTCAACATCTCGCCTCG  
CCGGCATGAGACGGTGCCAGGACTGCAATTGCTCAATCTATCCCGGCCACATAACGGGT  
CACCGTATGGCTTGGGATATGATGATGAACTGGTCGCCTACAACGGCCCTGGTGTAT  
CGCAGCTGCTCCGATCCCAAGCTGTCGTGGACATGGTGGCGGGGGCCCAATTGGGG  
AGTCCTGGCGGGTCTCGCCTACTATTCCATGGTGGGGAACGGGCTAAGGTTTGTATTG  
TGATGCTACTCTTTGCTCCCTAATAG

SEQ ID NO 23 (HCC138)

ATGTTGGGTAAGGTCATCGATACCCCTTACATGCGGCTTCGCCGACCTCGTGGGGTACA  
TTCCGCTCGTCGGCGCCCCCTAGGGGGCGCTGCCAGGGCCCTGGCGCATGGCGTCCG  
GGTTCTGGAGGACGGCGTGAACATATGCAACAGGGAATTTGCCCGGTTGCTCTTTCTCT  
ATCTTCCTCTTGCTTTGCTGTCTGTCTGACCGTTCCAGCTTCCGCTTATGAAGTGCG  
CAACGTGTCCGGGATGTACCATGTACGAACGACTGCTCCAACCTCAAGCATTGTGTAT  
GAGGCAGCGGACATGATCATGCACACCCCCGGGTGCGTGCCCTGCGTTCCGGGAGAAC  
AACTCTTCCCGCTGCTGGGTAGCGCTCACCCCCACGCTCGCAGCTAGGAACGCCAGCG  
TCCCCACCACGACAATACGACGCCACGTCGATTCCAGCTGTTCAACATCTCGCCTCG  
CCGGCATGAGACGGTGCCAGGACTGCAATTGCTCAATCTATCCCGGCCACATAACGGGT  
CACCGTATGGCTTGGGATATGATGATGAACTGGTAA  
TAG

SEQ ID NO 25 (HCC139)

ATGTTGGGTAAGGTCATCGATACCCCTTACATGCGGCTTCGCCGACCTCGTGGGGTACA  
TTCCGCTCGTCGGCGCCCCCTAGGGGGCGCTGCCAGGGCCCTGGCGCATGGCGTCCG  
GGTTCTGGAGGACGGCGTGAACATATGCAACAGGGAATTTGCCCGGTTGCTCTTTCTCT



25/65

## Fig. 21E

ATCTTCCTCTTGGCTTTGCTGTCCTGTCTGACCGTTCCAGCTTCCGCTTATGAAGTGCG  
CAACGTGTCCGGGATGTACCATGTACGAACGACTGCTCCAACCAAGCATTGTGTAT  
GAGGCAGCGGACATGATCATGCACACCCCGGGTGCGTGCCCTGCGTTCCGGGAGAAC  
AACTCTTCCCGCTGCTGGGTAGCGCTCACCCCCACGCTCGCAGCTAGGAACGCCAGCG  
TCCCCACCACGACAATACGACGCCACGTGATTCCCAGCTGTTACCATCTCGCCTCG  
CCGGCATGAGACGGTGCAGGACTGCAATTGCTCAATCTATCCCGGCCACATAACGGGT  
CACCGTATGGCTTGGGATATGATGATGAACTGGTCGCCTACAACGGCCCTGGTGGTAT  
CGCAGCTGCTCCGGATCCTCTAATAG

SEQ ID NO 27 (HCC140)

ATGTTGGGTAAGGTCATCGATACCCTTACATGCGGCTTCGCCGACCTCGTGGGGTACA  
TTCCGCTCGTCGGCGCCCCCTAGGGGGCGCTGCCAGGGCCCTGGCGCATGGCGTCCG  
GGTTCTGGAGGACGGCGTGAACATGCAACAGGGAATTTGCCCGGTTGCTCTTTCTCT  
ATCTTCCTCTTGGCTTTGCTGTCCTGTCTGACCGTTCCAGCTTCCGCTTATGAAGTGCG  
CAACGTGTCCGGGATGTACCATGTACGAACGACTGCTCCAACCAAGCATTGTGTAT  
GAGGCAGCGGACATGATCATGCACACCCCGGGTGCGTGCCCTGCGTTCCGGGAGAAC  
AACTCTTCCCGCTGCTGGGTAGCGCTCACCCCCACGCTCGCAGCTAGGAACGCCAGCG  
TCCCCACCACGACAATACGACGCCACGTGATTCCCAGCTGTTACCATCTCGCCTCG  
CCGGCATGAGACGGTGCAGGACTGCAATTGCTCAATCTATCCCGGCCACATAACGGGT  
CACCGTATGGCTTGGGATATGATGATGAACTGGTCGCCTACAACGGCCCTGGTGGTAT  
CGCAGCTGCTCCGGATCGTGATCGAGGGCAGACACCATCACCACCATCACTAATAG

SEQ ID NO 29 (HCC162)

ATGGGTAAGGTCATCGATACCCTTACGTGCGGATTCGCCGATCTCATGGGGTACATCC  
CGCTCGTCGGCGCTCCCGTAGGAGGCGTCGCAAGAGCCCTTGCGCATGGCGTGAGGGC  
CCTTGAAGACGGGATAAATTCGCAACAGGGAATTTGCCCGGTTGCTCCTTTTCTATTT  
TCCTTCTCGCTCTGTTCTCTTGCTTAATTCATCCAGCAGCTAGTCTAGAGTGGCGGAAT  
ACGTCTGGCCTCTATGTCCTTACCAACGACTGTTCCAATAGCAGTATTGTGTACGAGGC  
CGATGACGTTATTCTGCACACACCCGGCTGCATACCTTGTGTCCAGGACGGCAATACA  
TCCACGTGCTGGACCCAGTGACACCTACAGTGGCAGTCAAGTACGTGCGGAGCAACCA  
CCGCTTCGATACGCAGTCATGTGGACCTATTAGTGGGCGCGGCCACGATGTGCTCTGC  
GCTCTACGTGGGTGACATGTGTGGGGCTGTCTTCTCGTGGGACAAGCCTTCACGTTCA  
GACCTCGTCGCCATCAAACGGTCCAGACCTGTAAGTCTGCTGTACCCAGGCCATCT  
TTCAGGACATCGAATGGCTTGGGATATGATGATGAACTGGTAATAG

26/65

## Fig. 21F

SEQ ID NO 31 (HCCI63)

ATGGGTAAGGTCATCGATACCCTAACGTGCGGATTCGCCGATCTCATGGGGTATATCC  
CGCTCGTAGGGCGGCCCCATTGGGGGCGTCGCAAGGGCTCTCGCACACGGTGTGAGGGT  
CCTTGAGGACGGGGTAACTATGCAACAGGGAATTTACCCGGTTGCTCTTTCTCTATCT  
TTATTCTTGCTCTTCTCTCGTGTCTGACCGTTCGGCCTCTGCAGTTCCTACCGAAATG  
CCTCTGGGATTTATCATGTTACCAATGATTGCCCAAACCTCTCCATAGTCTATGAGGCA  
GATAACCTGATCCTACACGCACCTGGTTGCGTGCCTTGTGTCTATGACAGGTAATGTGA  
GTAGATGCTGGGTCCAAATTACCCCTACACTGTCAGCCCCGAGCCTCGGAGCAGTCAC  
GGCTCCTCTTCGGAGAGCCGTTGACTACCTAGCGGGAGGGGCTGCCCTCTGCTCCGCG  
TTATACGTAGGAGACGCGTGTGGGGCACTATTCTTGGTAGGCCAAATGTTACCTATA  
GGCCTCGCCAGCACGCTACGGTGCAGAACTGCAACTGTTCCATTACAGTGGCCATGT  
TACCGGCCACCGGATGGCATGGGATATGATGATGAACTGGTAATAG

SEQ ID NO 33 (HCP109)

TGGGATATGATGATGAACTGGTC

SEQ ID NO 34 (HCP72)

CTATTATGGTGGTAAGCCARCARGAGCAGGAG

SEQ ID NO 35 (HCCL22A)

TGGGATATGATGATGAACTGGTCGCCTACAACGGCCCTGGTGGTATCGCAGCTGCTCC  
GGATCCCAAGCTGTCTGGACATGGTGGCGGGGGCCCCATTGGGGAGTCTGGCGG  
GCCTCGCCTACTATTCCATGGTGGGGAAGTGGGCTAAGGTTTGGTTGTGATGCTACTC  
TTTGCCGGCGTCGACGGGCATACCCGCGTGTGAGGAGGGGCAGCAGCCTCCGATACCA  
GGGGCCTTGTGTCCCTCTTTAGCCCCGGGTGCGCTCAGAAAATCCAGCTCGTAAACAC  
CAACGGCAGTTGGCACATCAACAGGACTGCCCTGAACTGCAACGACTCCCTCCAAAC  
AGGGTTCTTTGCCGCACTATTCTACAAACACAAATTCAACTCGTCTGGATGCCAGAG  
CGCTTGCCAGCTGTCTGCTCCATCGACAAGTTCGCTCAGGGGTGGGGTCCCCTCACTT  
ACACTGAGCCTAACAGCTCGGACCAGAGGCCCTACTGCTGGCACTACGCGCCTCGACC  
GTGTGGTATTGTACCCGCGTCTCAGGTGTGCGGTCCAGTGATTGCTTACCCCGAGCC  
CTGTTGTGGTGGGGACGACCGATCGGTTTGGTGTCCCCACGTATAACTGGGGGGCGAA  
CGACTCGGATGTGCTGATTCTCAACAACACGCGGCCGCCGCGAGGCAACTGGTTCGGC  
TGTACATGGATGAATGGCACTGGGTTACCAAGACGTGTGGGGGGCCCCCGTGCAACA  
TCGGGGGGGCGCGCAACAACACCTTGACCTGCCCCACTGACTGTTTTCGGAAGCACCC  
CGAGGCCACCTACGCCAGATGCGGTTCTGGGCCCTGGCTGACACCTAGGTGTATGGTT

27/65

## Fig. 21G

CATTACCCATATAGGCTCTGGCACTACCCCTGCACTGTCAACTTCACCATCTTCAAGGT  
TAGGATGTACGTGGGGGGCGTGGAGCACAGGTTTGAAGCCGCATGCAATTGGACTCG  
AGGAGAGCGTTGTGACTTGGAGGACAGGGATAGATCAGAGCTTAGCCCCGTGCTGCTG  
TCTACAACAGAGTGGCAGATACTGCCCTGTTCTTACCACCCTGCCGGCCCTATCCA  
CCGGCCTGATCCACCTCCATCAGAACATCGTGGACGTGCAATACCTGTACGGTGTAGG  
GTCGGCGGTTGTCTCCCTTGTCTATCAAATGGGAGTATGTCTGTGCTCTTCTCTCTCT  
GGCAGACGCGCGCATCTGCGCCTGCTTATGGATGATGCTGCTGATAGCTCAAGCTGAG  
GCCGCCCTTAGAGAACCTGGTGGTCTCAATGCGGCGGCCGTGGCCGGGGCGCATGGC  
ACTCTTCTCTCTTGTGTTCTTCTGTGCTGCCTGGTACATCAAGGGCAGGCTGGTCCC  
TGGTGCGGCATACGCCCTTCTATGGCGTGTGGCCGCTGCTCCTGCTTCTGCTGGCCTTAC  
CACCACGAGCTTATGCCTAGTAA

SEQ ID NO 37 (HCC141)

GATCCCAACAAGCTGTCTGGACATGGTGGCGGGGGCCCCATTGGGGAGTCCTGGCGGG  
CCTCGCCTACTATTCCATGGTGGGGAAGTGGGCTAAGGTTTTGGTTGTGATGCTACTCT  
TTGCCGGCGTCGACGGGCATACCCGCGTGTCAAGAGGGGCAGCAGCCTCCGATACCA  
GGGGCCTTGTGTCCGTCTTTAGCCCCGGGTGGGCTCAGAAAATCCAGCTCGTAAACAC  
CAACGGCAGTTGGCACATCAACAGGACTGCCCTGAACTGCAACGACTCCCTCCAAAC  
AGGGTTCTTTGCCGCACTATTCTACAAACACAAATCAACTCGTCTGGATGCCCAGAG  
CGCTTGGCCAGCTGTCTCCATCGACAAGTTCTGCTCAGGGGTGGGGTCCCCCTCACTT  
ACACTGAGCCTAACAGCTCGGACCAGAGGCCCTACTGCTGGCACTACGCGCCTCGACC  
GTGTGGTATTGTACCCGCGTCTCAGGTGTGCGGTCCAGTGTATTGCTTACCCCCGAGCC  
CTGTTGTGGTGGGGACGACCGATCGGTTTGGTGTCCCCACGTATAACTGGGGGGCGAA  
CGACTCGGATGTGCTGATTCTCAACAACACGCGGCCGCCGCGAGGCAACTGGTTCGGC  
TGTACATGGATGAATGGCACTGGGTTTACCAAGACGTGTGGGGGGCCCCCGTGCAACA  
TCGGGGGGGGCCGGCAACAACAGCTTGACCTGCCCCACTGACTGTTTTCGGAAGCACCC  
CGAGGCCACCTACGCCAGATGCGGTTCTGGGCCCTGGCTGACACCTAGGTGTATGGTT  
CATTACCCATATAGGCTCTGGCACTACCCCTGCACTGTCAACTTCACCATCTTCAAGGT  
TAGGATGTACGTGGGGGGCGTGGAGCACAGGTTTGAAGCCGCATGCAATTGGACTCG  
AGGAGAGCGTTGTGACTTGGAGGACAGGGATAGATCAGAGCTTAGCCCCGTGCTGCTG  
TCTACAACAGAGTGGCAGAGTGGCAGAGCTTAATTAATTAG

SEQ ID NO 39 (HCC142)

GATCCCAACAAGCTGTCTGGACATGGTGGCGGGGGCCCCATTGGGGAGTCCTGGCGGG  
CCTCGCCTACTATTCCATGGTGGGGAAGTGGGCTAAGGTTTTGGTTGTGATGCTACTCT

28/65

## Fig. 21H

TTGCCGGCGTCGACGGGCATACCCGCGTGTCTCAGGAGGGGCAGCAGCCTCCGATACCA  
GGGGCCTTGTGTCCCTCTTTAGCCCCGGGTCTGGCTCAGAAAATCCAGCTCGTAAACAC  
CAACGGCAGTTGGCACATCAACAGGACTGCCCTGAACTGCAACGACTCCCTCCAAAC  
AGGGTTCTTTGCCGCACTATTCTACAAACACAAATTCAACTCGTCTGGATGCCCAGAG  
CGCTTGCCAGCTGTCTGCTCCATCGACAAGTTCGCTCAGGGGTGGGGTCCCCTCACTT  
ACACTGAGCCTAACAGCTCGGACCAGAGGCCCTACTGCTGGCACTACGCGCCTCGACC  
GTGTGGTATTGTACCCGCGTCTCAGGTGTGCGGTCCAGTGATTGCTTCACCCCGAGCC  
CTGTTGTGGTGGGGACGACCGATCGGTTTGGTGTCCCCACGTATAACTGGGGGGCGAA  
CGACTCGGATGTGCTGATTCTCAACAACACGCGGCCGCCGCGAGGCAACTGGTTCGGC  
TGTACATGGATGAATGGCACTGGGTTTACCAAGACGTGTGGGGGCCCCCGTGCAACA  
TCGGGGGGGCCGGCAACAACACCTTGACCTGCCCCACTGACTGTTTTCGGAAGCACCC  
CGAGGCCACCTACGCCAGATGCGGTTCTGGGCCCTGGCTGACACCTAGGTGTATGGTT  
CATTACCCATATAGGCTCTGGCACTACCCCTGCACTGTCAACTTCACCATCTTCAAGGT  
TAGGATGTACGTGGGGGGCGTGGAGCACAGGTTCTGAAGCCGCATGCAATTGGACTCG  
AGGAGAGCGTTGTGACTTGGAGGACAGGGATAGATCAGAGCTTAGCCCGCTGCTGCTG  
TCTACAACAGGTGATCGAGGGCAGACACCATCACCACCATCACTAATAG

SEQ ID NO 41 (HCCI43)

ATGGTGGGGAACTGGGCTAAGGTTTTGGTTGTGATGCTACTCTTTGCCGGCGTCGACG  
GGCATACCCGCGTGTCTCAGGAGGGGCAGCAGCCTCCGATACCAGGGGCCTTGTGTCCCT  
CTTTAGCCCCGGGTCTGGCTCAGAAAATCCAGCTCGTAAACACCAACGGCAGTTGGCAC  
ATCAACAGGACTGCCCTGAACTGCAACGACTCCCTCCAAACAGGGTTCTTTGCCGCAC  
TATTCTACAAACACAAATTCAACTCGTCTGGATGCCCAGAGCGCTTGCCAGCTGTCTG  
CTCCATCGACAAGTTCGCTCAGGGGTGGGGTCCCCTCACTTACACTGAGCCTAACAGC  
TCGGACCAGAGGCCCTACTGCTGGCACTACGCGCCTCGACCGTGTGGTATTGTACCCG  
CGTCTCAGGTGTGCGGTCCAGTGATTGCTTCACCCCGAGCCCTGTTGTGGTGGGGAC  
GACCGATCGGTTTGGTGTCCCCACGTATAACTGGGGGGCGAACGACTCGGATGTGCTG  
ATTCTCAACAACACGCGGCCGCCGCGAGGCAACTGGTTCGGCTGTACATGGATGAATG  
GCACTGGGTTTACCAAGACGTGTGGGGGCCCCCGTGCAACATCGGGGGGGCCGGCA  
ACAACACCTTGACCTGCCCCACTGACTGTTTTCGGAAGCACCCCGAGGCCACCTACGC  
CAGATGCGGTTCTGGGCCCTGGCTGACACCTAGGTGTATGGTTTATTACCCATATAGG  
CTCTGGCACTACCCCTGCACTGTCAACTTCACCATCTTCAAGGTAGGATGTACGTGGG  
GGGCGTGGAGCACAGGTTCTGAAGCCGCATGCAATTGGACTCGAGGAGAGCGTTGTGA  
CTTGGAGGACAGGGATAGATCAGAGCTTAGCCCGCTGCTGCTGTCTACAACAGAGTGG  
CAGAGCTTAATTAATTAG

29/65

## Fig. 21I

SEQ ID NO 43 (HCCI44)

ATGGTGGGGAAC TGGGCTAAGGTTTTGGTTGTGATGCTACTCTTTGCCGGCGTCGACG  
GGCATACCCGCGTGT CAGGAGGGGCAGCAGCCTCCGATACCAGGGGCCTTGTGTCCCT  
CTTTAGCCCCGGGT CCGGCTCAGAAAATCCAGCTCGTAAACACCAACGGCAGTTGGCAC  
ATCAACAGGACTGCCCTGAACTGCAACGACTCCCTCCAAACAGGGTTCTTTGCCGCAC  
TATTCTACAAACACAAATTCAACTCGTCTGGATGCCCAGAGCGCTTGCCAGCTGTCTG  
CTCCATCGACAAGTTCGCTCAGGGGTGGGGTCCCCTCACTTACACTGAGCCTAACAGC  
TCGGACCAGAGGCCCTACTGCTGGCACTACGCGCCTCGACCGTGTGGTATTGTACCCG  
CGTCTCAGGTGTGCGGTCCAGTGTATTGCTTACCCCCGAGCCCTGTTGTGGTGGGGAC  
GACCGATCGGTTTTGGTGTCCCCACGTATAACTGGGGGGCGAACGACTCGGATGTGCTG  
ATTCTCAACAACACGCGGCCGCCGCGAGGCAACTGGTTCGGCTGTACATGGATGAATG  
GCACTGGGTTACCAAGACGTGTGGGGGCCCCCGTGCAACATCGGGGGGGCCGGCA  
ACAACACCTTGACCTGCCCCACTGACTGTTTTCGGAAGCACCCCGAGGCCACCTACGC  
CAGATGCGGTTCTGGGCCCTGGCTGACACCTAGGTGTATGGTTCATTACCCATATAGG  
CTCTGGCACTACCCCTGCACTGTCAACTTCACCATCTTCAAGGTTAGGATGTACGTGGG  
GGGCGTGGAGCACAGGTTGGAAGCCGCATGCAATTGGACTCGAGGAGAGCGTTGTGA  
CTTGGAGGACAGGGATAGATCAGAGCTTAGCCCGCTGCTGCTGTCTACAACAGGTGAT  
CGAGGGCAGACACCATCACCACCATCACTAATAG

SEQ ID NO 45 (HCCL64)

ATGGTGGCGGGGGCCCATTTGGGGAGTCTGGCGGGCCTCGCCTACTATTCCATGGTGG  
GGAAC TGGGCTAAGGTTTTGGTTGTGATGCTACTCTTTGCCGGCGTCGACGGGCATAC  
CCGCGTGT CAGGAGGGGCAGCAGCCTCCGATACCAGGGGCCTTGTGTCCCTCTTTAGC  
CCCGGGT CCGGCTCAGAAAATCCAGCTCGTAAACACCAACGGCAGTTGGCACATCAAC  
AGGACTGCCCTGAACTGCAACGACTCCCTCCAAACAGGGTTCTTTGCCGCACTATTCT  
ACAAACACAAAATTCAACTCGTCTGGATGCCCAGAGCGCTTGGCCAGCTGTGCTCCAT  
CGACAAGTTCGCTCAGGGGTGGGGTCCCCTCACTTACACTGAGCCTAACAGCTCGGAC  
CAGAGGCCCTACTGCTGGCACTACGCGCCTCGACCGTGTGGTATTGTACCCGCGTCTC  
AGGTGTGCGGTCCAGTGTATTGCTTACCCCGAGCCCTGTTGTGGTGGGGACGACCGA  
TCGGTTTGGTGTCCCCACGTATAACTGGGGGGCGAACGACTCGGATGTGCTGATTCTC  
AACAACACGCGGCCGCCGCGAGGCAACTGGTTCGGCTGTACATGGATGAATGGCACT  
GGGTTACCAAGACGTGTGGGGGCCCCCGTGCAACATCGGGGGGGCCGGCAACAAC  
ACCTTGACCTGCCCCACTGACTGTTTTCGGAAGCACCCCGAGGCCACCTACGCCAGAT  
GCGGTTCTGGGCCCTGGCTGACACCTAGGTGTATGGTTCATTACCCATATAGGCTCTGG  
CACTACCCCTGCACTGTCAACTTCACCATCTTCAAGGTTAGGATGTACGTGGGGGGCG

30/65

Fig. 21J

TGGAGCACAGGTTTGAAGCCGCATGCAATTGGACTCGAGGAGAGCGTTGTGACTTGGA  
GGACAGGGATAGATCAGAGCTTAGCCCCGCTGCTGCTGTCTACAACAGAGTGCCAGATA  
CTGCCCTGTTCTTCACCACCCTGCCGGCCCTATCCACCGCCTGATCCACCTCCATCA  
GAACATCGTGGACGTGCAATACCTGTACGGTGTAGGGTCGGCGGTTGTCTCCCTTGTC  
ATCAAATGGGAGTATGTCCTGTTGCTCTTCCTTCTCCTGGCAGACGCGCGCATCTGCGC  
CTGCTTATGGATGATGCTGCTGATAGCTCAAGCTGAGGCCGCCTTAGAGAACCTGGTG  
GTCCTCAATGCGGCGGCCGTGGCCGGGGCGCATGGCACTCTTTCCTTCTTGTGTTCTT  
CTGTGCTGCCTGGTACATCAAGGGCAGGCTGGTCCCTGGTGCGGCATACGCCTTCTAT  
GGCGTGTGGCCGCTGCTCCTGCTTCTGCTGGCCTTACCACCAGAGCTTATGCCTAGTAA

SEQ ID NO 47 (HCC165)

AATTGGGTAAGGTCATCGATACCCCTACATGCGGCTTCGCCGACCTCGTGGGGTACA  
TTCCGCTCGTCGGCGCCCCCTAGGGGGCGCTGCCAGGGCCCTGGCGCATGGCGTCCG  
GGTCTGGAGGACGGCGTGAACATGCAACAGGGAATTTGCCCGTTGCTCTTTCTCT  
ATCTTCCTCTTGGCTTTGCTGTCTGCTGACCGTTCCAGCTTCCGCTTATGAAGTGCG  
CAACGTGTCCGGATGTACCATGTACGAACGACTGCTCCAACTCAAGCATTGTGTAT  
GAGGCAGCGGACATGATCATGCACACCCCCGGGTGCGTGCCCTGCGTTCCGGGAGAAC  
AACTCTTCCCGCTGCTGGGTAGCGCTACCCCCACGCTCGCAGCTAGGAACGCCAGCG  
TCCCCACCACGACAATACGACGCCACGTGATTTGCTCGTTGGGGCGGCTGCTTTCTG  
TTCCGCTATGTACGTGGGGGACCTCTGCGGATCTGTCTTCTCGTCTCCAGCTGTTCA  
CCATCTCGCTCGCCGGCATGAGACGGTGCAGGACTGCAATTGCTCAATCTATCCCCG  
CCACATAACGGGTCACCGTATGGCTTGGGATATGATGATGAAGTGGTCGCCTACAACG  
GCCCTGGTGGTATCGCAGCTGCTCCGGATCCCACAAGCTGTGCGTGGACATGGTGGCGG  
GGGCCCATTTGGGGAGTCTTGGCGGGCCTCGCCTACTATTCCATGGTGGGGA'ACTGGGC  
TAAGGTTTTGGTTGTGATGCTACTCTTTGCCGGCGTCGACGGGCATACCCGCGTGTAG  
GAGGGGCAGCAGCCTCCGATACCAGGGGCCTTGTGTCCCTCTTTAGCCCCGGGTGCGC  
TCAGAAAATCCAGCTCGTAAACACCAACGGCAGTTGGCACATCAACAGGACTGCCCT  
GAACTGCAACGACTCCCTCCAAACAGGGTTCTTTGCCGCACTATTCT'ACAAACAAAA  
TTCAACTCCTCTGGATGCCCAGAGCGCTTGGCCAGCTGTGCTCCATCGACAAGTTCC  
CTCAGGGGTGGGGTCCCCTCACTTACACTGAGCCTAACAGCTCGGACAGAGGCCCTA  
CTGTGGCACTACGCGCCTCGACCGTGTGGTATTGTACCCGCGTCTCAGGTGTGCGGT  
CCAGTGTATTGCTTACCCCGAGCCCTGTTGTGGTGGGGACGACCGATCGGTTTGGTGT  
CCCCACGTATAACTGGGGGGCGAA'CGACTCGGATGTGCTGATTCTCAACAACACGCGG  
CCGCCGCGAGGCAACTGGTTCGGCTGTACATGGATGAATGGCACTGGGTTCACCAAGA  
CGTGTGGGGGGCCCCCGTGCAACATCGGGGGGGCCGGCAACAACACCTTGACCTGCC

31/65

## Fig. 21K

CCACTGACTGTTTTCGGAAGCACCCCGAGGCCACCTACGCCAGATGCGGTTCTGGGCC  
CTGGCTGACACCTAGGTGTATGGTTCATTACCCATATAGGCTCTGGCACTACCCCTGCA  
CTGTCAACTTCACCATCTTCAAGGTTAGGATGTACGTGGGGGGCGTGGAGCACAGGTT  
CGAAGCCGCATGCAATTGGA CTGAGGAGAGCGTTGTGACTTGGAGGACAGGGATAG  
ATCAGAGCTTAGCCCGCTGCTGCTGTCTACAACAGAGTGGCAGATACTGCCCTGTTCC  
TTCACCACCCTGCCGGCCCTATCCACCGGCTGATCCACCTCCATCAGAACATCGTGG  
ACGTGCAATACCTGTACGGTGTAGGGTCGGCGGTTGTCTCCCTTGTCAATAATGGGA  
GTATGTCCTGTTGCTCTTCTTCTCCTGGCAGACGCGCGCATCTGCGCCTGCTTATGGA  
TGATGCTGCTGATAGCTCAAGCTGAGGCCGCTTAGAGAACCTGGTGGTCCTCAATGC  
GGCGGCCGTGGCCGGGGCGCATGGCACTCTTTCCTTCTTGTGTTCTTCTGTGCTGCCT  
GGTACATCAAGGGCAGGCTGGTCCCTGGTGGGCATACGCCCTTCTATGGCGTGTGGCC  
GCTGCTCCTGCTTCTGCTGGCCTTACCACCACGAGCTTATGCCTAGTAAGCTT

SEQ ID NO 49 (HCC165)

ATGAGCACGAATCCTAAACCTCAAAGAAAAACCAAACGTAACACCAACCGCCGCCCA  
CAGGACGTCAAGTTCCCGGGCGGTGGTCAGATCGTTGGTGGAGTTTACCTGTTGCCGC  
GCAGGGGCCCCAGGTTGGGTGTGCGCGCGACTAGGAAGACTTCCGAGCGGTGCAAC  
CTCGTGGGAGGCGACAACCTATCCCCAAGGCTCGCCGACCCGAGGGTAGGGCCTGGG  
CTCAGCCCGGTACCCCTGGCCCTCTATGGCAATGAGGGCATGGGGTGGGCAGGATG  
GCTCCTGTACCCCGCGGCTCTCGGCCTAGTTGGGGCCCTACAGACCCCGGCGTAGG  
TCGCGTAATTTGGTAAGGTCATCGATACCCCTTACATGCGGCTTCGCCGACCTCGTGG  
GGTACATTCGCTCGTGGCGCCCCCTAGGGGGCGCTGCCAGGGCCCTGGCGCATGG  
CGTCCGGGTTCTGGAGGACGGCGTGAACATATGCAACAGGGAATTTGCCCGGTTGCTCT  
TTCTCTATCTTCTTCTTGGCTTTGCTGTCTGTGACCGTTCCAGCTTCCGCTTATGAA  
GTGCGCAACGTGTCCGGGATGTACCATGTACGAACGACTGCTCCA ACTCAAGCATTG  
TGTATGAGGCAGCGGACATGATCATGCACACCCCGGGTGGTGGCCTGCGTTCGGGA  
GAACA ACTCTTCCCGCTGCTGGGTAGCGCTACCCCCACGCTCGCAGCTAGGAACGCC  
AGCGTCCCCACACGACAATACGACGCCACGTCGATTTGCTCGTTGGGGCGGCTGCTT  
TCTGTTCCGCTATGTACGTGGGGGACCTCTGCGGATCTGTCTTCTCGTCTCCAGCTG  
TTCACCATCTCGCCTCGCCGGCATGAGACGGTGCAGGACTGCAATTGCTCAATCTATC  
CCGGCCACATAACGGGTCACCGTATGGCTTGGGATATGATGATGAACTGGTCGCCTAC  
AACGGCCCTGGTGGTATCGCAGCTGCTCCGGATCCCAACAAGCTGTCGTGGACATGGTG  
GCGGGGGCCCCATTGGGGAGTCTGGCGGGCCCTCGCCTACTATTCCATGGTGGGGAACT  
GGGCTAAGGTTTTGGTTGTGATGCTACTCTTTGCCGGCGTCGACGGGCATACCCGCGT  
GTCAGGAGGGGCAGCAGCCTCCGATACCAGGGGCCTTGTGTCCCTCTTTAGCCCCGGG

32/65

Fig. 21L

TCGGCTCAGAAAATCCAGCTCGTAAACACCAACGGCAGTTGGCACATCAACAGGACT  
GCCCTGAACTGCAACGACTCCCTCCAAACAGGGTTCTTTGCCGCACTATTCTACAAAC  
ACAAATTCAACTCGTCTGGATGCCCAGAGCGCTTGGCCAGCTGTCGCTCCATCGACAA  
GTTTCGCTCAGGGGTGGGGTCCCCTCACTTACACTGAGCCTAACAGCTCGGACCAGAGG  
CCCTACTGCTGGCACTACGCGCCTCGACCCTGTGGTATTGTACCCGCGTCTCAGGTGT  
GCGGTCCAGTGTATTGCTTACCCCGAGCCCTGTTGTGGTGGGGACGACCGATCGGTT  
TGGTGTCCCCACGTATAACTGGGGGGCGAACGACTCGGATGTGCTGATTCTCAACAAC  
ACGCGGCCCGCGGAGGCAACTGGTTCGGCTGTACATGGATGAATGGCACTGGGTTCA  
CCAAGACGTGTGGGGGCCCCCGTGCAACATCGGGGGGGCCGGCAACAACACCTTGA  
CCTGCCCCACTGACTGTTTTCGGAAGCACCCCGAGGCCACCTACGCCAGATGCGGTTT  
TGGGCCCTGGCTGACACCTAGGTGTATGGTTCATTACCCATATAGGCTCTGGCACTAC  
CCCTGCACTGTCAACTTCACCATCTTCAAGGTTAGGATGTACGTGGGGGGCGTGGAGC  
ACAGGTTCTGAAGCCGCATGCAATTGGAATCGAGGAGAGCGTTGTGACTTGGAGGACA  
GGGATAGATCAGAGCTTAGCCCGCTGCTGCTGTCTACAACAGAGTGGCAGATACTGCC  
CTGTTCTTCACCAACCCTGCCGGCCCTATCCACCGGCCTGATCCACCTCCATCAGAAC  
ATCGTGGACGTGCAATACCTGTACGGTGTAGGGTCGGCGGTTGTCTCCCTTGTCTATCA  
AATGGGAGTATGTCCTGTTGCTCTTCTCTCTGCGAGACGCGCGCATCTGCGCCTGC  
TTATGGATGATGCTGCTGATAAGCTCAAGCTGAGGCCGCTTAGAGAACCTGGTGGTCC  
TCAATGCGGCGGCCGTGGCCGGGGCGCATGGCACTCTTCTCTCTTGTGTTCTTCTGT  
GCTGCCTGGTACATCAAGGGCAGGCTGGTCCCTGGTGCGGCATACGCCTTCTATGGCG  
TGTGGCCGCTGCTCCTGCTTCTGCTGGCCTTACCACCACGAGCTTATGCCTAGTAA



33/65

Fig. 22

OD measured at 450 nm						
construct						
Fraction	volume	dilution	39 Type Ib	40 Type Ib	62 Type 3a	63 Type 5a
START	23 ml	1/20	2.517	1.954	1.426	1.142
FLOW THROUGH	23 ml	1/20	0.087	0.085	0.176	0.120
1	0.4 ml	1/200	0.102	0.051	0.048	0.050
2			0.396	0.550	0.090	0.067
3			2.627	2.603	2.481	2.372
4			3	2.967	3	2.694
5			3	2.810	2.640	2.154
6			2.694	2.499	1.359	1.561
7			2.408	2.481	0.347	1.390
8			2.176	1.970	1.624	0.865
9			1.461	1.422	0.887	0.604
10			1.286	0.926	0.543	0.519
11			0.981	0.781	0.294	0.294
12			0.812	0.650	0.249	0.199
13			0.373	0.432	0.239	0.209
14			0.653	0.371	0.145	0.134
15			0.441	0.348	0.151	0.151
16			0.321	0.374	0.098	0.106
17			0.525	0.186	0.099	0.108
18			0.351	0.171	0.083	0.090
19			0.192	0.164	0.084	0.087

34/65




Fig. 23

35/65

Figure 24

Fraction	volume	dilution	OD measured at 450 nm			
			construct			
			39 Type 1b	40 Type 1b	62 Type 3a	63 Type 5a
20	250 $\mu$ l	1/200	0.072	0.130	0.096	0.051
21			0.109	0.293	0.084	0.052
22			0.279	0.249	0.172	0.052
23			0.093	0.151	0.297	0.054
24			0.080	0.266	0.438	0.056
25			0.251	0.100	0.457	0.048
26			3	1.649	0.722	0.066
27			3	3	2.528	0.889
28			3	3	3	2.345
29			3	3	2.849	2.580
30			2.227	1.921	1.424	1.333
31			0.263	0.415	0.356	0.162
32			0.071	0.172	0.154	0.064
33			0.103	0.054	0.096	0.057
34			0.045	0.045	0.044	0.051
35			0.043	0.047	0.045	0.046
36			0.045	0.045	0.049	0.040
37			0.045	0.047	0.046	0.048
38			0.046	0.048	0.047	0.057
39			0.045	0.048	0.050	0.057
40			0.046	0.049	0.048	0.049

36/65




Fig. 25

37/65




Fig.26




Fig.27

38/65




Fig.28




Fig.29

Lane 1: Crude Lysate  
 Lane 2: Flow through Lentil Chromatography  
 Lane 3: Wash with EMPIGEN Lentil Chromatography  
 Lane 4: Eluate Lentil Chromatography  
 Lane 5: Flow through during concentration lentil eluate  
 Lane 6: Pool of Eluate after Size Exclusion Chromatography

39/65





Fig. 30

40/65

NON-REDUCED

Fig. 31A

E2 + CONTAMINANTS (AGGREGATES)



REDUCED




Fig. 31B



41/65




Fig. 32

42/65

## SILVER STAIN OF PURIFIED E2





Fig.33

43/65



No.	Ret. (min)	Peak start (min)	Peak end (min)	Dur (min)	Area (min* $\mu$ AU)	Height ( $\mu$ AU)
1	-0.45	-0.46	-0.43	0.04	0.0976	4.579
2	1.55	0.75	3.26	2.51	796.4167	889.377
3	3.27	3.26	3.31	0.05	0.0067	0.224
4	3.33	3.32	3.33	0.02	0.0002	0.018

Total number of detected peaks = 4  
 Total Area above baseline = 0.796522 ml\*AU  
 Total area in evaluated peaks = 0.796521 ml\*AU  
 Ratio peak area / total area = 0.999999  
 Total peak duration = 2.613583 ml

Fig. 34

44/65




Fig. 35A-1




Fig. 35A-2




Fig. 35A-3





Fig. 35A-4



35A-5




Fig. 35A-6

45/65




Fig. 35A-7




Fig. 35A-8

46/65




Fig. 35B-1




Fig. 35B-2




Fig. 35B-3





Fig. 35B-4



35B-5




Fig. 35B-6




Fig. 35B-7




Fig. 35B-8

48/65

Fig. 36A

E1 Ab




Fig. 36B

E2 Ab






Fig. 37B  
Long Term Responders




Fig. 37D  
Type 3a




Fig. 37A  
Non Responders






Fig. 37C  
Type 1b



50/65

Fig. 38

Relative Map Positions of  
anti-E2 monoclonal antibodies





PARTIAL TREATMENT OF HCV E2\E2s ENVELOPE PROTEINS  
BY PNGase F




Fig. 40

53/65

Fig. 41 *In Vitro* Mutagenesis of HCV E1 glycoprotein







Fig. 42A *In Vitro* Mutagenesis of HCV E1 glycoprotein



1. First step of PCR amplification (Gly-# and Ovr-# primers)









57/65




Fig. 44A




Fig. 44B

58/65




Fig. 45

KDa 119 67 43 29 18




Fig. 46

59/65

Fig. 47

	age (years)	HCV infection (years)	genotype
Marcel	17	9	1a
Peggy	21	16.5	1b
Fenna	15	9	1a
Yoran	12	none	
Marti	12	none	

chronic carriers (strong T-cell adjuvant)


↓ ↓ ↓ ↓ ↓ ↓      ↓ ↓ ↓ ↓ ↓ ↓ 50 µg E1 dose  
0 3 6 9 12 15      26 29 32 35 38 41 weeks

naive (alum)

↓ ↓ ↓ ↓ ↓ ↓      50 µg E1 dose  
0 3 6 9 12 15 weeks


60/65

Fig. 48




61/65

Fig. 49



62/65

Fig. 50



63/65




Fig 51





Fig 52




65/65

Fig 53

vaccinated



controls



## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: Maertens, Geert  
Bosman, Fons  
De Martynoff, Guy  
Buyse, Marie-Ange
- (ii) TITLE OF INVENTION: Purified hepatitis C virus envelope proteins for diagnostic and therapeutic use.
- (iii) NUMBER OF SEQUENCES: 111
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: NIXON & VANDERHYE
  - (B) STREET: 1100 North Glebe Road, 8th Floor
  - (C) CITY: Arlington, VA 22201
  - (E) COUNTRY: USA
  - (F) ZIP: 22201
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER: Unassigned
  - (B) FILING DATE: 01-DEC-2000
  - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Sadoff, B. J.
  - (B) REGISTRATION NUMBER: 36663
  - (C) REFERENCE/DOCKET NUMBER: 2551-53
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (703) 816-4000
  - (B) TELEFAX: (703) 816-4100

## (2) INFORMATION FOR SEQ ID NO: 1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iii) ANTI-SENSE: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

GGCATGCAAG CTTAATTAAT T

21

## (2) INFORMATION FOR SEQ ID NO: 2:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 68 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

CCGGGGAGGC CTGCACGTGA TCGAGGGCAG ACACCATCAC CACCATCACT AATAGTTAAT

60

TAACTGCA

68

## (2) INFORMATION FOR SEQ ID NO: 3:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 642 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..639

## (ix) FEATURE:

- (A) NAME/KEY: mat\_peptide
- (B) LOCATION: 1..636

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

ATG CCC GGT TGC TCT TTC TCT ATC TTC CTC TTG GCT TTA CTG TCC TGT  
 Met Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu Leu Ser Cys  
 1 5 10 15

48

CTG ACC ATT CCA GCT TCC GCT TAT GAG GTG CGC AAC GTG TCC GGG ATG  
 Leu Thr Ile Pro Ala Ser Ala Tyr Glu Val Arg Asn Val Ser Gly Met  
 20 25 30

96

TAC CAT GTC ACG AAC GAC TGC TCC AAC TCA AGC ATT GTG TAT GAG GCA  
 Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val Tyr Glu Ala

144

35	40	45	
GCG GAC ATG ATC ATG CAC ACC CCC GGG TGC GTG CCC TGC GTT CGG GAG			192
Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys Val Arg Glu			
50	55	60	
AAC AAC TCT TCC CGC TGC TGG GTA GCG CTC ACC CCC ACG CTC GCA GCT			240
Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr Leu Ala Ala			
65	70	75	80
AGG AAC GCC AGC GTC CCC ACC ACG ACA ATA CGA CGC CAC GTC GAT TTG			288
Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His Val Asp Leu			
	85	90	95
CTC GTT GGG GCG GCT GCT CTC TGT TCC GCT ATG TAC GTG GGG GAT CTC			336
Leu Val Gly Ala Ala Ala Leu Cys Ser Ala Met Tyr Val Gly Asp Leu			
	100	105	110
TGC GGA TCT GTC TTC CTC GTC TCC CAG CTG TTC ACC ATC TCG CCT CGC			384
Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Ile Ser Pro Arg			
	115	120	125
CGG CAT GAG ACG GTG CAG GAC TGC AAT TGC TCA ATC TAT CCC GGC CAC			432
Arg His Glu Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His			
	130	135	140
ATA ACA GGT CAC CGT ATG GCT TGG GAT ATG ATG ATG AAC TGG TCG CCT			480
Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp Ser Pro			
145	150	155	160
ACA ACG GCC CTG GTG GTA TCG CAG CTG CTC CGG ATC CCA CAA GCT GTC			528
Thr Thr Ala Leu Val Val Ser Gln Leu Leu Arg Ile Pro Gln Ala Val			
	165	170	175
GTG GAC ATG GTG GCG GGG GCC CAT TGG GGA GTC CTG GCG GGC CTC GCC			576
Val Asp Met Val Ala Gly Ala His Trp Gly Val Leu Ala Gly Leu Ala			
	180	185	190
TAC TAT TCC ATG GTG GGG AAC TGG GCT AAG GTT TTG ATT GTG ATG CTA			624
Tyr Tyr Ser Met Val Gly Asn Trp Ala Lys Val Leu Ile Val Met Leu			
	195	200	205
CTC TTT GCT CTC TAATAG			642
Leu Phe Ala Leu			
210			

## (2) INFORMATION FOR SEQ ID NO: 4:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 212 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Met	Pro	Gly	Cys	Ser	Phe	Ser	Ile	Phe	Leu	Leu	Ala	Leu	Leu	Ser	Cys
1					5				10					15	
Leu	Thr	Ile	Pro	Ala	Ser	Ala	Tyr	Glu	Val	Arg	Asn	Val	Ser	Gly	Met

20										25					30																
Tyr	His	Val	Thr	Asn	Asp	Cys	Ser	Asn	Ser	Ser	Ile	Val	Tyr	Glu	Ala																
		35						40					45																		
Ala	Asp	Met	Ile	Met	His	Thr	Pro	Gly	Cys	Val	Pro	Cys	Val	Arg	Glu																
		50					55				60																				
Asn	Asn	Ser	Ser	Arg	Cys	Trp	Val	Ala	Leu	Thr	Pro	Thr	Leu	Ala	Ala																
		65			70					75					80																
Arg	Asn	Ala	Ser	Val	Pro	Thr	Thr	Thr	Ile	Arg	Arg	His	Val	Asp	Leu																
				85					90					95																	
Leu	Val	Gly	Ala	Ala	Ala	Leu	Cys	Ser	Ala	Met	Tyr	Val	Gly	Asp	Leu																
			100					105					110																		
Cys	Gly	Ser	Val	Phe	Leu	Val	Ser	Gln	Leu	Phe	Thr	Ile	Ser	Pro	Arg																
		115					120					125																			
Arg	His	Glu	Thr	Val	Gln	Asp	Cys	Asn	Cys	Ser	Ile	Tyr	Pro	Gly	His																
		130				135					140																				
Ile	Thr	Gly	His	Arg	Met	Ala	Trp	Asp	Met	Met	Met	Asn	Trp	Ser	Pro																
145					150				155					160																	
Thr	Thr	Ala	Leu	Val	Val	Ser	Gln	Leu	Leu	Arg	Ile	Pro	Gln	Ala	Val																
			165						170					175																	
Val	Asp	Met	Val	Ala	Gly	Ala	His	Trp	Gly	Val	Leu	Ala	Gly	Leu	Ala																
			180					185					190																		
Tyr	Tyr	Ser	Met	Val	Gly	Asn	Trp	Ala	Lys	Val	Leu	Ile	Val	Met	Leu																
		195					200					205																			
Leu	Phe	Ala	Leu																												
		210																													

## (2) INFORMATION FOR SEQ ID NO: 5:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 795 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..792

## (ix) FEATURE:

- (A) NAME/KEY: mat\_peptide
- (B) LOCATION: 1..789

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

ATG TTG GGT AAG GTC ATC GAT ACC CTT ACA TGC GGC TTC GCC GAC CTC	48
Met Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu	
1 5 10 15	
GTG GGG TAC ATT CCG CTC GTC GGC GCC CCC CTA GGG GGC GCT GCC AGG	96
Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg	
20 25 30	
GCC CTG GCG CAT GGC GTC CGG GTT CTG GAG GAC GGC GTG AAC TAT GCA	144
Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala	
35 40 45	
ACA GGG AAT TTG CCC GGT TGC TCT TTC TCT ATC TTC CTC TTG GCT TTG	192
Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu	
50 55 60	
CTG TCC TGT CTG ACC GTT CCA GCT TCC GCT TAT GAA GTG CGC AAC GTG	240
Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr Glu Val Arg Asn Val	
65 70 75 80	
TCC GGG ATG TAC CAT GTC ACG AAC GAC TGC TCC AAC TCA AGC ATT GTG	288
Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val	
85 90 95	
TAT GAG GCA GCG GAC ATG ATC ATG CAC ACC CCC GGG TGC GTG CCC TGC	336
Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys	
100 105 110	
GTT CGG GAG AAC AAC TCT TCC CGC TGC TGG GTA GCG CTC ACC CCC ACG	384
Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr	
115 120 125	
CTC GCA GCT AGG AAC GCC AGC GTC CCC ACC ACG ACA ATA CGA CGC CAC	432
Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His	
130 135 140	
GTC GAT TTG CTC GTT GGG GCG GCT GCT TTC TGT TCC GCT ATG TAC GTG	480
Val Asp Leu Leu Val Gly Ala Ala Ala Phe Cys Ser Ala Met Tyr Val	
145 150 155 160	
GGG GAC CTC TGC GGA TCT GTC TTC CTC GTC TCC CAG CTG TTC ACC ATC	528
Gly Asp Leu Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Ile	
165 170 175	
TCG CCT CGC CGG CAT GAG ACG GTG CAG GAC TGC AAT TGC TCA ATC TAT	576
Ser Pro Arg Arg His Glu Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr	
180 185 190	
CCC GGC CAC ATA ACG GGT CAC CGT ATG GCT TGG GAT ATG ATG ATG AAC	624
Pro Gly His Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn	
195 200 205	
TGG TCG CCT ACA ACG GCC CTG GTG GTA TCG CAG CTG CTC CGG ATC CCA	672
Trp Ser Pro Thr Thr Ala Leu Val Val Ser Gln Leu Leu Arg Ile Pro	
210 215 220	
CAA GCT GTC GTG GAC ATG GTG GCG GGG GCC CAT TGG GGA GTC CTG GCG	720
Gln Ala Val Val Asp Met Val Ala Gly Ala His Trp Gly Val Leu Ala	
225 230 235 240	
GGT CTC GCC TAC TAT TCC ATG GTG GGG AAC TGG GCT AAG GTT TTG ATT	768

Gly Leu Ala Tyr Tyr Ser Met Val Gly Asn Trp Ala Lys Val Leu Ile  
245 250 255

GTG ATG CTA CTC TTT GCT CCC TAATAG  
Val Met Leu Leu Phe Ala Pro  
260

795

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 263 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Met 1	Leu	Gly	Lys	Val 5	Ile	Asp	Thr	Leu	Thr 10	Cys	Gly	Phe	Ala	Asp	Leu
Val	Gly	Tyr	Ile 20	Pro	Leu	Val	Gly	Ala 25	Pro	Leu	Gly	Gly	Ala 30	Ala	Arg
Ala	Leu	Ala 35	His	Gly	Val	Arg	Val 40	Leu	Glu	Asp	Gly	Val 45	Asn	Tyr	Ala
Thr	Gly 50	Asn	Leu	Pro	Gly	Cys 55	Ser	Phe	Ser	Ile	Phe 60	Leu	Leu	Ala	Leu
Leu 65	Ser	Cys	Leu	Thr	Val 70	Pro	Ala	Ser	Ala	Tyr 75	Glu	Val	Arg	Asn	Val 80
Ser	Gly	Met	Tyr	His 85	Val	Thr	Asn	Asp 90	Cys	Ser	Asn	Ser	Ser	Ile 95	Val
Tyr	Glu	Ala 100	Ala	Asp	Met	Ile	Met 105	His	Thr	Pro	Gly	Cys 110	Val	Pro	Cys
Val	Arg	Glu 115	Asn	Asn	Ser	Ser	Arg 120	Cys	Trp	Val	Ala	Leu 125	Thr	Pro	Thr
Leu 130	Ala	Ala	Arg	Asn	Ala	Ser 135	Val	Pro	Thr	Thr	Thr 140	Ile	Arg	Arg	His
Val 145	Asp	Leu	Leu	Val	Gly 150	Ala	Ala	Ala	Phe	Cys 155	Ser	Ala	Met	Tyr	Val 160
Gly	Asp	Leu	Cys 165	Gly	Ser	Val	Phe	Leu	Val 170	Ser	Gln	Leu	Phe 175	Thr	Ile
Ser	Pro	Arg 180	Arg	His	Glu	Thr	Val 185	Gln	Asp	Cys	Asn	Cys 190	Ser	Ile	Tyr
Pro	Gly	His 195	Ile	Thr	Gly	His	Arg 200	Met	Ala	Trp	Asp	Met 205	Met	Met	Asn
Trp 210	Ser	Pro	Thr	Thr	Ala	Leu 215	Val	Val	Ser	Gln	Leu 220	Leu	Arg	Ile	Pro
Gln	Ala	Val	Val	Asp	Met	Val	Ala	Gly	Ala	His	Trp	Gly	Val	Leu	Ala

```

225                230                235                240
Gly Leu Ala Tyr Tyr Ser Met Val Gly Asn Trp Ala Lys Val Leu Ile
                245                250                255
Val Met Leu Leu Phe Ala Pro
                260

```

## (2) INFORMATION FOR SEQ ID NO: 7:

```

(i) SEQUENCE CHARACTERISTICS:
    (A) LENGTH: 633 base pairs
    (B) TYPE: nucleic acid
    (C) STRANDEDNESS: single
    (D) TOPOLOGY: linear

```

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

```

(ix) FEATURE:
    (A) NAME/KEY: CDS
    (B) LOCATION: 1..630

```

```

(ix) FEATURE:
    (A) NAME/KEY: mat_peptide
    (B) LOCATION: 1..627

```

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

```

ATG TTG GGT AAG GTC ATC GAT ACC CTT ACG TGC GGC TTC GCC GAC CTC      48
Met Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu
  1                5                10                15

ATG GGG TAC ATT CCG CTC GTC GGC GCC CCC CTA GGG GGT GCT GCC AGA      96
Met Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg
                20                25                30

GCC CTG GCG CAT GGC GTC CGG GTT CTG GAA GAC GGC GTG AAC TAT GCA     144
Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala
                35                40                45

ACA GGG AAT TTG CCT GGT TGC TCT TTC TCT ATC TTC CTC TTG GCT TTA     192
Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu
                50                55                60

CTG TCC TGT CTG ACC ATT CCA GCT TCC GCT TAT GAG GTG CGC AAC GTG     240
Leu Ser Cys Leu Thr Ile Pro Ala Ser Ala Tyr Glu Val Arg Asn Val
                65                70                75                80

TCC GGG ATG TAC CAT GTC ACG AAC GAC TGC TCC AAC TCA AGC ATT GTG     288
Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val
                85                90                95

TAT GAG GCA GCG GAC ATG ATC ATG CAC ACC CCC GGG TGC GTG CCC TGC     336
Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys
                100                105                110

```



GTT CGG GAG AAC AAC TCT TCC CGC TGC TGG GTA GCG CTC ACC CCC ACG	384
Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr	
115 120 125	
CTC GCA GCT AGG AAC GCC AGC GTC CCC ACT ACG ACA ATA CGA CGC CAC	432
Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His	
130 135 140	
GTC GAT TTG CTC GTT GGG GCG GCT GCT TTC TGT TCC GCT ATG TAC GTG	480
Val Asp Leu Leu Val Gly Ala Ala Ala Phe Cys Ser Ala Met Tyr Val	
145 150 155 160	
GGG GAT CTC TGC GGA TCT GTC TTC CTC GTC TCC CAG CTG TTC ACC ATC	528
Gly Asp Leu Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Ile	
165 170 175	
TCG CCT CGC CGG CAT GAG ACG GTG CAG GAC TGC AAT TGC TCA ATC TAT	576
Ser Pro Arg Arg His Glu Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr	
180 185 190	
CCC GGC CAC ATA ACA GGT CAC CGT ATG GCT TGG GAT ATG ATG ATG AAC	624
Pro Gly His Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn	
195 200 205	
TGG TAATAG	633
Trp	
210	

## (2) INFORMATION FOR SEQ ID NO: 8:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 209 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Met Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu	
1 5 10 15	
Met Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg	
20 25 30	
Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala	
35 40 45	
Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu	
50 55 60	
Leu Ser Cys Leu Thr Ile Pro Ala Ser Ala Tyr Glu Val Arg Asn Val	
65 70 75 80	
Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val	
85 90 95	
Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys	
100 105 110	

Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr  
 115 120 125

Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His  
 130 135 140

Val Asp Leu Leu Val Gly Ala Ala Ala Phe Cys Ser Ala Met Tyr Val  
 145 150 155 160

Gly Asp Leu Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Ile  
 165 170 175

Ser Pro Arg Arg His Glu Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr  
 180 185 190

Pro Gly His Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn  
 195 200 205

Trp

## (2) INFORMATION FOR SEQ ID NO: 9:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 483 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

- (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 1..480

- (ix) FEATURE:  
 (A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 1..477

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

ATG CCC GGT TGC TCT TTC TCT ATC TTC CTC TTG GCC CTG CTG TCC TGT	48
Met Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu Leu Ser Cys	
1 5 10 15	
CTG ACC ATA CCA GCT TCC GCT TAT GAA GTG CGC AAC GTG TCC GGG GTG	96
Leu Thr Ile Pro Ala Ser Ala Tyr Glu Val Arg Asn Val Ser Gly Val	
20 25 30	
TAC CAT GTC ACG AAC GAC TGC TCC AAC TCA AGC ATA GTG TAT GAG GCA	144
Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val Tyr Glu Ala	
35 40 45	
GCG GAC ATG ATC ATG CAC ACC CCC GGG TGC GTG CCC TGC GTT CGG GAG	192
Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys Val Arg Glu	
50 55 60	

GGC AAC TCC TCC CGT TGC TGG GTG GCG CTC ACT CCC ACG CTC GCG GCC	240
Gly Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr Leu Ala Ala	
65 70 75 80	
AGG AAC GCC AGC GTC CCC ACA ACG ACA ATA CGA CGC CAC GTC GAT TTG	288
Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His Val Asp Leu	
85 90 95	
CTC GTT GGG GCT GCT GCT TTC TGT TCC GCT ATG TAC GTG GGG GAT CTC	336
Leu Val Gly Ala Ala Ala Phe Cys Ser Ala Met Tyr Val Gly Asp Leu	
100 105 110	
TGC GGA TCT GTT TTC CTT GTT TCC CAG CTG TTC ACC TTC TCA CCT CGC	384
Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Phe Ser Pro Arg	
115 120 125	
CGG CAT CAA ACA GTA CAG GAC TGC AAC TGC TCA ATC TAT CCC GGC CAT	432
Arg His Gln Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His	
130 135 140	
GTA TCA GGT CAC CGC ATG GCT TGG GAT ATG ATG ATG AAC TGG TCC TAATAG	483
Val Ser Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp Ser	
145 150 155 160	

## (2) INFORMATION FOR SEQ ID NO: 10:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 159 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Met Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu Leu Ser Cys	
1 5 10 15	
Leu Thr Ile Pro Ala Ser Ala Tyr Glu Val Arg Asn Val Ser Gly Val	
20 25 30	
Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val Tyr Glu Ala	
35 40 45	
Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys Val Arg Glu	
50 55 60	
Gly Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr Leu Ala Ala	
65 70 75 80	
Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His Val Asp Leu	
85 90 95	
Leu Val Gly Ala Ala Ala Phe Cys Ser Ala Met Tyr Val Gly Asp Leu	
100 105 110	
Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Phe Ser Pro Arg	
115 120 125	
Arg His Gln Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His	

130	135	140
Val Ser Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp Ser		
145	150	155

## (2) INFORMATION FOR SEQ ID NO: 11:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 480 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: CDS  
 (B) LOCATION: 1..477

## (ix) FEATURE:

- (A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 1..474

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

ATG TCC GGT TGC TCT TTC TCT ATC TTC CTC TTG GCC CTG CTG TCC TGT	48
Met Ser Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu Leu Ser Cys	
1 5 10 15	
CTG ACC ATA CCA GCT TCC GCT TAT GAA GTG CGC AAC GTG TCC GGG GTG	96
Leu Thr Ile Pro Ala Ser Ala Tyr Glu Val Arg Asn Val Ser Gly Val	
20 25 30	
TAC CAT GTC ACG AAC GAC TGC TCC AAC TCA AGC ATA GTG TAT GAG GCA	144
Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val Tyr Glu Ala	
35 40 45	
GCG GAC ATG ATC ATG CAC ACC CCC GGG TGC GTG CCC TGC GTT CGG GAG	192
Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys Val Arg Glu	
50 55 60	
GGC AAC TCC TCC CGT TGC TGG GTG GCG CTC ACT CCC ACG CTC GCG GCC	240
Gly Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr Leu Ala Ala	
65 70 75 80	
AGG AAC GCC AGC GTC CCC ACA ACG ACA ATA CGA CGC CAC GTC GAT TTG	288
Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His Val Asp Leu	
85 90 95	
CTC GTT GGG GCT GCT GCT TTC TGT TCC GCT ATG TAC GTG GGG GAT CTC	336
Leu Val Gly Ala Ala Phe Cys Ser Ala Met Tyr Val Gly Asp Leu	
100 105 110	
TGC GGA TCT GTT TTC CTT GTT TCC CAG CTG TTC ACC TTC TCA CCT CGC	384
Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Phe Ser Pro Arg	

115	120	125	
CGG CAT CAA ACA GTA CAG GAC TGC AAC TGC TCA ATC TAT CCC GGC CAT			432
Arg His Gln Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His			
130	135	140	
GTA TCA GGT CAC CGC ATG GCT TGG GAT ATG ATG ATG AAC TGG TAATAG			480
Val Ser Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp			
145	150	155	

## (2) INFORMATION FOR SEQ ID NO: 12:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 158 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

Met	Ser	Gly	Cys	Ser	Phe	Ser	Ile	Phe	Leu	Leu	Ala	Leu	Leu	Ser	Cys
1				5					10					15	
Leu	Thr	Ile	Pro	Ala	Ser	Ala	Tyr	Glu	Val	Arg	Asn	Val	Ser	Gly	Val
			20					25					30		
Tyr	His	Val	Thr	Asn	Asp	Cys	Ser	Asn	Ser	Ser	Ile	Val	Tyr	Glu	Ala
		35				40						45			
Ala	Asp	Met	Ile	Met	His	Thr	Pro	Gly	Cys	Val	Pro	Cys	Val	Arg	Glu
	50					55				60					
Gly	Asn	Ser	Ser	Arg	Cys	Trp	Val	Ala	Leu	Thr	Pro	Thr	Leu	Ala	Ala
65				70					75					80	
Arg	Asn	Ala	Ser	Val	Pro	Thr	Thr	Thr	Ile	Arg	Arg	His	Val	Asp	Leu
				85					90					95	
Leu	Val	Gly	Ala	Ala	Ala	Phe	Cys	Ser	Ala	Met	Tyr	Val	Gly	Asp	Leu
			100				105						110		
Cys	Gly	Ser	Val	Phe	Leu	Val	Ser	Gln	Leu	Phe	Thr	Phe	Ser	Pro	Arg
		115					120					125			
Arg	His	Gln	Thr	Val	Gln	Asp	Cys	Asn	Cys	Ser	Ile	Tyr	Pro	Gly	His
	130				135						140				
Val	Ser	Gly	His	Arg	Met	Ala	Trp	Asp	Met	Met	Met	Asn	Trp		
145					150					155					

## (2) INFORMATION FOR SEQ ID NO: 13:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 636 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..633

(ix) FEATURE:

(A) NAME/KEY: mat\_peptide

(B) LOCATION: 1..630

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

ATG CTG GGT AAG GCC ATC GAT ACC CTT ACG TGC GGC TTC GCC GAC CTC	48
Met Leu Gly Lys Ala Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu	
1 5 10 15	
GTG GGG TAC ATT CCG CTC GTC GGC GCC CCC CTA GGG GGC GCT GCC AGG	96
Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg	
20 25 30	
GCC CTG GCG CAT GGC GTC CGG GTT CTG GAA GAC GGC GTG AAC TAT GCA	144
Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala	
35 40 45	
ACA GGG AAT TTG CCT GGT TGC TCT TTC TCT ATC TTC CTC TTG GCT TTA	192
Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu	
50 55 60	
CTG TCC TGT CTA ACC ATT CCA GCT TCC GCT TAC GAG GTG CGC AAC GTG	240
Leu Ser Cys Leu Thr Ile Pro Ala Ser Ala Tyr Glu Val Arg Asn Val	
65 70 75 80	
TCC GGG ATG TAC CAT GTC ACG AAC GAC TGC TCC AAC TCA AGC ATT GTG	288
Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val	
85 90 95	
TAT GAG GCA GCG GAC ATG ATC ATG CAC ACC CCC GGG TGC GTG CCC TGC	336
Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys	
100 105 110	
GTT CGG GAG AAC AAC TCT TCC CGC TGC TGG GTA GCG CTC ACC CCC ACG	384
Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr	
115 120 125	
CTC GCG GCT AGG AAC GCC AGC ATC CCC ACT ACA ACA ATA CGA CGC CAC	432
Leu Ala Ala Arg Asn Ala Ser Ile Pro Thr Thr Thr Ile Arg Arg His	
130 135 140	
GTC GAT TTG CTC GTT GGG GCG GCT GCT TTC TGT TCC GCT ATG TAC GTG	480
Val Asp Leu Leu Val Gly Ala Ala Ala Phe Cys Ser Ala Met Tyr Val	
145 150 155 160	
GGG GAT CTC TGC GGA TCT GTC TTC CTC GTC TCC CAG CTG TTC ACC ATC	528
Gly Asp Leu Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Ile	
165 170 175	
TCG CCT CGC CGG CAT GAG ACG GTG CAG GAC TGC AAT TGC TCA ATC TAT	576
Ser Pro Arg Arg His Glu Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr	

14

180	185	190	
CCC GGC CAC ATA ACG GGT CAC CGT ATG GCT TGG GAT ATG ATG ATG AAC			624
Pro Gly His Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn			
195	200	205	
TGG TAC TAATAG			640
Trp Tyr			
210			

## (2) INFORMATION FOR SEQ ID NO: 14:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 210 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

Met	Leu	Gly	Lys	Ala	Ile	Asp	Thr	Leu	Thr	Cys	Gly	Phe	Ala	Asp	Leu	
1				5				10						15		
Val	Gly	Tyr	Ile	Pro	Leu	Val	Gly	Ala	Pro	Leu	Gly	Gly	Ala	Ala	Arg	
			20					25					30			
Ala	Leu	Ala	His	Gly	Val	Arg	Val	Leu	Glu	Asp	Gly	Val	Asn	Tyr	Ala	
		35					40					45				
Thr	Gly	Asn	Leu	Pro	Gly	Cys	Ser	Phe	Ser	Ile	Phe	Leu	Leu	Ala	Leu	
		50				55					60					
Leu	Ser	Cys	Leu	Thr	Ile	Pro	Ala	Ser	Ala	Tyr	Glu	Val	Arg	Asn	Val	
65					70					75					80	
Ser	Gly	Met	Tyr	His	Val	Thr	Asn	Asp	Cys	Ser	Asn	Ser	Ser	Ile	Val	
				85				90						95		
Tyr	Glu	Ala	Ala	Asp	Met	Ile	Met	His	Thr	Pro	Gly	Cys	Val	Pro	Cys	
			100					105					110			
Val	Arg	Glu	Asn	Asn	Ser	Ser	Arg	Cys	Trp	Val	Ala	Leu	Thr	Pro	Thr	
			115					120					125			
Leu	Ala	Ala	Arg	Asn	Ala	Ser	Ile	Pro	Thr	Thr	Thr	Ile	Arg	Arg	His	
			130				135					140				
Val	Asp	Leu	Leu	Val	Gly	Ala	Ala	Ala	Phe	Cys	Ser	Ala	Met	Tyr	Val	
145					150					155					160	
Gly	Asp	Leu	Cys	Gly	Ser	Val	Phe	Leu	Val	Ser	Gln	Leu	Phe	Thr	Ile	
			165						170					175		
Ser	Pro	Arg	Arg	His	Glu	Thr	Val	Gln	Asp	Cys	Asn	Cys	Ser	Ile	Tyr	
			180					185					190			
Pro	Gly	His	Ile	Thr	Gly	His	Arg	Met	Ala	Trp	Asp	Met	Met	Met	Asn	
			195				200					205				

Trp Tyr  
210

(2) INFORMATION FOR SEQ ID NO: 15:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 26 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

ATGCCCGGTT GCTCTTTCTC TATCTT

26

(2) INFORMATION FOR SEQ ID NO: 16:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 26 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

ATGTTGGGTA AGGTCATCGA TACCCT

26

(2) INFORMATION FOR SEQ ID NO: 17:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 30 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:



CTATTAGGAC CAGTTCATCA TCATATCCCA

30

(2) INFORMATION FOR SEQ ID NO: 18:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 27 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

CTATTACCAG TTCATCATCA TATCCCA

27

(2) INFORMATION FOR SEQ ID NO: 19:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 36 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

ATACGACGCC ACGTCGATTC CCAGCTGTTC ACCATC

36

(2) INFORMATION FOR SEQ ID NO: 20:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 36 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

GATGGTGAAC AGCTGGGAAT CGACGTGGCG TCGTAT

36

(2) INFORMATION FOR SEQ ID NO: 21:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 723 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(ix) FEATURE:

- (A) NAME/KEY: CDS  
 (B) LOCATION: 1..720

(ix) FEATURE:

- (A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 1..717

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

ATG TTG GGT AAG GTC ATC GAT ACC CTT ACA TGC GGC TTC GCC GAC CTC	48
Met Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu	
1 5 10 15	
GTG GGG TAC ATT CCG CTC GTC GGC GCC CCC CTA GGG GGC GCT GCC AGG	96
Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg	
20 25 30	
GCC CTG GCG CAT GGC GTC CGG GTT CTG GAG GAC GGC GTG AAC TAT GCA	144
Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala	
35 40 45	
ACA GGG AAT TTG CCC GGT TGC TCT TTC TCT ATC TTC CTC TTG GCT TTG	192
Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu	
50 55 60	
CTG TCC TGT CTG ACC GTT CCA GCT TCC GCT TAT GAA GTG CGC AAC GTG	240
Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr Glu Val Arg Asn Val	
65 70 75 80	
TCC GGG ATG TAC CAT GTC ACG AAC GAC TGC TCC AAC TCA AGC ATT GTG	288
Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val	
85 90 95	
TAT GAG GCA GCG GAC ATG ATC ATG CAC ACC CCC GGG TGC GTG CCC TGC	336
Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys	
100 105 110	
GTT CGG GAG AAC AAC TCT TCC CGC TGC TGG GTA GCG CTC ACC CCC ACG	384
Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr	
115 120 125	
CTC GCA GCT AGG AAC GCC AGC GTC CCC ACC ACG ACA ATA CGA CGC CAC	432
Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His	
130 135 140	
GTC GAT TCC CAG CTG TTC ACC ATC TCG CCT CGC CGG CAT GAG ACG GTG	480
Val Asp Ser Gln Leu Phe Thr Ile Ser Pro Arg Arg His Glu Thr Val	

145	150	155	160	
CAG GAC TGC AAT TGC TCA ATC TAT CCC GGC CAC ATA ACG GGT CAC CGT				528
Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His Ile Thr Gly His Arg	165	170	175	
ATG GCT TGG GAT ATG ATG ATG AAC TGG TCG CCT ACA ACG GCC CTG GTG				576
Met Ala Trp Asp Met Met Met Asn Trp Ser Pro Thr Thr Ala Leu Val	180	185	190	
GTA TCG CAG CTG CTC CGG ATC CCA CAA GCT GTC GTG GAC ATG GTG GCG				624
Val Ser Gln Leu Leu Arg Ile Pro Gln Ala Val Val Asp Met Val Ala	195	200	205	
GGG GCC CAT TGG GGA GTC CTG GCG GGT CTC GCC TAC TAT TCC ATG GTG				672
Gly Ala His Trp Gly Val Leu Ala Gly Leu Ala Tyr Tyr Ser Met Val	210	215	220	
GGG AAC TGG GCT AAG GTT TTG ATT GTG ATG CTA CTC TTT GCT CCC TAATAG				723
Gly Asn Trp Ala Lys Val Leu Ile Val Met Leu Leu Phe Ala Pro	225	230	235	240

## (2) INFORMATION FOR SEQ ID NO: 22:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 239 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

Met	Leu	Gly	Lys	Val	Ile	Asp	Thr	Leu	Thr	Cys	Gly	Phe	Ala	Asp	Leu
1				5					10					15	
Val	Gly	Tyr	Ile	Pro	Leu	Val	Gly	Ala	Pro	Leu	Gly	Gly	Ala	Ala	Arg
			20					25					30		
Ala	Leu	Ala	His	Gly	Val	Arg	Val	Leu	Glu	Asp	Gly	Val	Asn	Tyr	Ala
			35				40						45		
Thr	Gly	Asn	Leu	Pro	Gly	Cys	Ser	Phe	Ser	Ile	Phe	Leu	Leu	Ala	Leu
	50					55					60				
Leu	Ser	Cys	Leu	Thr	Val	Pro	Ala	Ser	Ala	Tyr	Glu	Val	Arg	Asn	Val
65				70					75					80	
Ser	Gly	Met	Tyr	His	Val	Thr	Asn	Asp	Cys	Ser	Asn	Ser	Ser	Ile	Val
			85						90					95	
Tyr	Glu	Ala	Ala	Asp	Met	Ile	Met	His	Thr	Pro	Gly	Cys	Val	Pro	Cys
			100					105					110		
Val	Arg	Glu	Asn	Asn	Ser	Ser	Arg	Cys	Trp	Val	Ala	Leu	Thr	Pro	Thr
			115				120					125			
Leu	Ala	Ala	Arg	Asn	Ala	Ser	Val	Pro	Thr	Thr	Thr	Ile	Arg	Arg	His
	130					135						140			
Val	Asp	Ser	Gln	Leu	Phe	Thr	Ile	Ser	Pro	Arg	Arg	His	Glu	Thr	Val

145		150		155		160
Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His Ile Thr Gly His Arg						
		165		170		175
Met Ala Trp Asp Met Met Met Asn Trp Ser Pro Thr Thr Ala Leu Val						
		180		185		190
Val Ser Gln Leu Leu Arg Ile Pro Gln Ala Val Val Asp Met Val Ala						
		195		200		205
Gly Ala His Trp Gly Val Leu Ala Gly Leu Ala Tyr Tyr Ser Met Val						
		210		215		220
Gly Asn Trp Ala Lys Val Leu Ile Val Met Leu Leu Phe Ala Pro						
		225		230		235

## (2) INFORMATION FOR SEQ ID NO: 23:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 561 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..558

## (ix) FEATURE:

- (A) NAME/KEY: mat\_peptide
- (B) LOCATION: 1..555

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

ATG TTG GGT AAG GTC ATC GAT ACC CTT ACA TGC GGC TTC GCC GAC CTC	48
Met Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu	
1 5 10 15	
GTG GGG TAC ATT CCG CTC GTC GGC GCC CCC CTA GGG GGC GCT GCC AGG	96
Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg	
20 25 30	
GCC CTG GCG CAT GGC GTC CGG GTT CTG GAG GAC GGC GTG AAC TAT GCA	144
Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala	
35 40 45	
ACA GGG AAT TTG CCC GGT TGC TCT TTC TCT ATC TTC CTC TTG GCT TTG	192
Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu	
50 55 60	
CTG TCC TGT CTG ACC GTT CCA GCT TCC GCT TAT GAA GTG CGC AAC GTG	240
Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr Glu Val Arg Asn Val	

65	70	75	80	
TCC GGG ATG TAC CAT GTC ACG AAC GAC TGC TCC AAC TCA AGC ATT GTG				288
Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val	85	90	95	
TAT GAG GCA GCG GAC ATG ATC ATG CAC ACC CCC GGG TGC GTG CCC TGC				336
Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys	100	105	110	
GTT CGG GAG AAC AAC TCT TCC CGC TGC TGG GTA GCG CTC ACC CCC ACG				384
Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr	115	120	125	
CTC GCA GCT AGG AAC GCC AGC GTC CCC ACC ACG ACA ATA CGA CGC CAC				432
Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His	130	135	140	
GTC GAT TCC CAG CTG TTC ACC ATC TCG CCT CGC CGG CAT GAG ACG GTG				480
Val Asp Ser Gln Leu Phe Thr Ile Ser Pro Arg Arg His Glu Thr Val	145	150	155	160
CAG GAC TGC AAT TGC TCA ATC TAT CCC GGC CAC ATA ACG GGT CAC CGT				528
Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His Ile Thr Gly His Arg	165	170	175	
ATG GCT TGG GAT ATG ATG ATG AAC TGG TAATAG				561
Met Ala Trp Asp Met Met Met Asn Trp	180	185		

## (2) INFORMATION FOR SEQ ID NO: 24:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 185 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

Met Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu  
 1 5 10 15

Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg  
 20 25 30

Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala  
 35 40 45

Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu  
 50 55 60

Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr Glu Val Arg Asn Val  
 65 70 75 80

Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val  
 85 90 95

Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys  
 100 105 110

```

Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr
    115                      120                      125
Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His
    130                      135                      140
Val Asp Ser Gln Leu Phe Thr Ile Ser Pro Arg Arg His Glu Thr Val
    145                      150                      155                      160
Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His Ile Thr Gly His Arg
    165                      170                      175
Met Ala Trp Asp Met Met Met Asn Trp
    180                      185

```

## (2) INFORMATION FOR SEQ ID NO: 25:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 606 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..603

## (ix) FEATURE:

- (A) NAME/KEY: mat\_peptide
- (B) LOCATION: 1..600

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

```

ATG TTG GGT AAG GTC ATC GAT ACC CTT ACA TGC GGC TTC GCC GAC CTC      48
Met Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu
  1                      5                      10                      15

GTG GGG TAC ATT CCG CTC GTC GGC GCC CCC CTA GGG GGC GCT GCC AGG      96
Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg
    20                      25                      30

GCC CTG GCG CAT GGC GTC CGG GTT CTG GAG GAC GGC GTG AAC TAT GCA      144
Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala
    35                      40                      45

ACA GGG AAT TTG CCC GGT TGC TCT TTC TCT ATC TTC CTC TTG GCT TTG      192
Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu
    50                      55                      60

CTG TCC TGT CTG ACC GTT CCA GCT TCC GCT TAT GAA GTG CGC AAC GTG      240
Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr Glu Val Arg Asn Val
    65                      70                      75                      80

```

TCC GGG ATG TAC CAT GTC ACG AAC GAC TGC TCC AAC TCA AGC ATT GTG	288
Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val	
85 90 95	
TAT GAG GCA GCG GAC ATG ATC ATG CAC ACC CCC GGG TGC GTG CCC TGC	336
Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys	
100 105 110	
GTT CGG GAG AAC AAC TCT TCC CGC TGC TGG GTA GCG CTC ACC CCC ACG	384
Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr	
115 120 125	
CTC GCA GCT AGG AAC GCC AGC GTC CCC ACC ACG ACA ATA CGA CGC CAC	432
Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His	
130 135 140	
GTC GAT TCC CAG CTG TTC ACC ATC TCG CCT CGC CGG CAT GAG ACG GTG	480
Val Asp Ser Gln Leu Phe Thr Ile Ser Pro Arg Arg His Glu Thr Val	
145 150 155 160	
CAG GAC TGC AAT TGC TCA ATC TAT CCC GGC CAC ATA ACG GGT CAC CGT	528
Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His Ile Thr Gly His Arg	
165 170 175	
ATG GCT TGG GAT ATG ATG ATG AAC TGG TCG CCT ACA ACG GCC CTG GTG	576
Met Ala Trp Asp Met Met Met Asn Trp Ser Pro Thr Thr Ala Leu Val	
180 185 190	
GTA TCG CAG CTG CTC CGG ATC CTC TAATAG	606
Val Ser Gln Leu Leu Arg Ile Leu	
195 200	

## (2) INFORMATION FOR SEQ ID NO: 26:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 200 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

Met Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu	
1 5 10 15	
Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg	
20 25 30	
Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala	
35 40 45	
Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu	
50 55 60	
Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr Glu Val Arg Asn Val	
65 70 75 80	
Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val	
85 90 95	

Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys  
 100 105 110  
 Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr  
 115 120 125  
 Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His  
 130 135 140  
 Val Asp Ser Gln Leu Phe Thr Ile Ser Pro Arg Arg His Glu Thr Val  
 145 150 155 160  
 Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His Ile Thr Gly His Arg  
 165 170 175  
 Met Ala Trp Asp Met Met Met Asn Trp Ser Pro Thr Thr Ala Leu Val  
 180 185 190  
 Val Ser Gln Leu Leu Arg Ile Leu  
 195 200

## (2) INFORMATION FOR SEQ ID NO: 27:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 636 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..633

## (ix) FEATURE:

- (A) NAME/KEY: mat\_peptide
- (B) LOCATION: 1..630

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

ATG TTG GGT AAG GTC ATC GAT ACC CTT ACA TGC GGC TTC GCC GAC CTC	48
Met Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu	
1 5 10 15	
GTG GGG TAC ATT CCG CTC GTC GGC GCC CCC CTA GGG GGC GCT GCC AGG	96
Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg	
20 25 30	
GCC CTG GCG CAT GGC GTC CGG GTT CTG GAG GAC GGC GTG AAC TAT GCA	144
Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala	
35 40 45	
ACA GGG AAT TTG CCC GGT TGC TCT TTC TCT ATC TTC CTC TTG GCT TTG	192



Thr	Gly	Asn	Leu	Pro	Gly	Cys	Ser	Phe	Ser	Ile	Phe	Leu	Leu	Ala	Leu		
50						55					60						
CTG	TCC	TGT	CTG	ACC	GTT	CCA	GCT	TCC	GCT	TAT	GAA	GTG	CGC	AAC	GTG	240	
Leu	Ser	Cys	Leu	Thr	Val	Pro	Ala	Ser	Ala	Tyr	Glu	Val	Arg	Asn	Val		
65					70					75					80		
TCC	GGG	ATG	TAC	CAT	GTC	ACG	AAC	GAC	TGC	TCC	AAC	TCA	AGC	ATT	GTG	288	
Ser	Gly	Met	Tyr	His	Val	Thr	Asn	Asp	Cys	Ser	Asn	Ser	Ser	Ile	Val		
				85				90						95			
TAT	GAG	GCA	GCG	GAC	ATG	ATC	ATG	CAC	ACC	CCC	GGG	TGC	GTG	CCC	TGC	336	
Tyr	Glu	Ala	Ala	Asp	Met	Ile	Met	His	Thr	Pro	Gly	Cys	Val	Pro	Cys		
			100					105					110				
GTT	CGG	GAG	AAC	AAC	TCT	TCC	CGC	TGC	TGG	GTA	GCG	CTC	ACC	CCC	ACG	384	
Val	Arg	Glu	Asn	Asn	Ser	Ser	Arg	Cys	Trp	Val	Ala	Leu	Thr	Pro	Thr		
		115					120					125					
CTC	GCA	GCT	AGG	AAC	GCC	AGC	GTC	CCC	ACC	ACG	ACA	ATA	CGA	CGC	CAC	432	
Leu	Ala	Ala	Arg	Asn	Ala	Ser	Val	Pro	Thr	Thr	Ile	Arg	Arg	His			
		130				135					140						
GTC	GAT	TCC	CAG	CTG	TTC	ACC	ATC	TCC	CCT	CGC	CGG	CAT	GAG	ACG	GTG	480	
Val	Asp	Ser	Gln	Leu	Phe	Thr	Ile	Ser	Pro	Arg	Arg	His	Glu	Thr	Val		
145					150					155					160		
CAG	GAC	TGC	AAT	TGC	TCA	ATC	TAT	CCC	GGC	CAC	ATA	ACG	GGT	CAC	CGT	528	
Gln	Asp	Cys	Asn	Cys	Ser	Ile	Tyr	Pro	Gly	His	Ile	Thr	Gly	His	Arg		
			165					170						175			
ATG	GCT	TGG	GAT	ATG	ATG	ATG	AAC	TGG	TCG	CCT	ACA	ACG	GCC	CTG	GTG	576	
Met	Ala	Trp	Asp	Met	Met	Met	Asn	Trp	Ser	Pro	Thr	Thr	Ala	Leu	Val		
			180					185					190				
GTA	TCG	CAG	CTG	CTC	CGG	ATC	GTG	ATC	GAG	GGC	AGA	CAC	CAT	CAC	CAC	624	
Val	Ser	Gln	Leu	Leu	Arg	Ile	Val	Ile	Glu	Gly	Arg	His	His	His	His		
		195				200					205						
CAT	CAC	TAATAG														636	
His	His																
210																	

## (2) INFORMATION FOR SEQ ID NO: 28:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 210 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:

Met	Leu	Gly	Lys	Val	Ile	Asp	Thr	Leu	Thr	Cys	Gly	Phe	Ala	Asp	Leu		
1				5					10					15			
Val	Gly	Tyr	Ile	Pro	Leu	Val	Gly	Ala	Pro	Leu	Gly	Gly	Ala	Ala	Arg		
			20					25					30				
Ala	Leu	Ala	His	Gly	Val	Arg	Val	Leu	Glu	Asp	Gly	Val	Asn	Tyr	Ala		

25

```

          35              40              45
Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu
  50              55              60
Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr Glu Val Arg Asn Val
  65              70              75              80
Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val
              85              90              95
Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys
              100              105              110
Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr
              115              120              125
Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His
              130              135              140
Val Asp Ser Gln Leu Phe Thr Ile Ser Pro Arg Arg His Glu Thr Val
              145              150              155              160
Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His Ile Thr Gly His Arg
              165              170              175
Met Ala Trp Asp Met Met Met Asn Trp Ser Pro Thr Thr Ala Leu Val
              180              185              190
Val Ser Gln Leu Leu Arg Ile Val Ile Glu Gly Arg His His His His
              195              200              205
His His
              210

```

## (2) INFORMATION FOR SEQ ID NO: 29:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 630 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..627

## (ix) FEATURE:

- (A) NAME/KEY: mat\_peptide
- (B) LOCATION: 1..624

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:

ATG GGT AAG GTC ATC GAT ACC CTT ACG TGC GGA TTC GCC GAT CTC ATG

48

Met	Gly	Lys	Val	Ile	Asp	Thr	Leu	Thr	Cys	Gly	Phe	Ala	Asp	Leu	Met	
1				5					10					15		
GGG	TAC	ATC	CCG	CTC	GTC	GGC	GCT	CCC	GTA	GGA	GGC	CTC	GCA	AGA	GCC	96
Gly	Tyr	Ile	Pro	Leu	Val	Gly	Ala	Pro	Val	Gly	Gly	Val	Ala	Arg	Ala	
			20					25					30			
CTT	GCG	CAT	GGC	GTG	AGG	GCC	CTT	GAA	GAC	GGG	ATA	AAT	TTC	GCA	ACA	144
Leu	Ala	His	Gly	Val	Arg	Ala	Leu	Glu	Asp	Gly	Ile	Asn	Phe	Ala	Thr	
		35					40					45				
GGG	AAT	TTG	CCC	GGT	TGC	TCC	TTT	TCT	ATT	TTC	CTT	CTC	GCT	CTG	TTC	192
Gly	Asn	Leu	Pro	Gly	Cys	Ser	Phe	Ser	Ile	Phe	Leu	Leu	Ala	Leu	Phe	
	50					55					60					
TCT	TGC	TTA	ATT	CAT	CCA	GCA	GCT	AGT	CTA	GAG	TGG	CGG	AAT	ACG	TCT	240
Ser	Cys	Leu	Ile	His	Pro	Ala	Ala	Ser	Leu	Glu	Trp	Arg	Asn	Thr	Ser	
	65				70					75					80	
GGC	CTC	TAT	GTC	CTT	ACC	AAC	GAC	TGT	TCC	AAT	AGC	AGT	ATT	GTG	TAC	288
Gly	Leu	Tyr	Val	Leu	Thr	Asn	Asp	Cys	Ser	Asn	Ser	Ser	Ile	Val	Tyr	
			85						90					95		
GAG	GCC	GAT	GAC	GTT	ATT	CTG	CAC	ACA	CCC	GGC	TGC	ATA	CCT	TGT	GTC	336
Glu	Ala	Asp	Val	Ile	Leu	His	Thr	Pro	Gly	Cys	Ile	Pro	Cys	Val		
			100				105						110			
CAG	GAC	GGC	AAT	ACA	TCC	ACG	TGC	TGG	ACC	CCA	GTG	ACA	CCT	ACA	GTG	384
Gln	Asp	Gly	Asn	Thr	Ser	Thr	Cys	Trp	Thr	Pro	Val	Thr	Pro	Thr	Val	
		115					120					125				
GCA	GTC	AAG	TAC	GTC	GGA	GCA	ACC	ACC	GCT	TCG	ATA	CGC	AGT	CAT	GTG	432
Ala	Val	Lys	Tyr	Val	Gly	Ala	Thr	Thr	Ala	Ser	Ile	Arg	Ser	His	Val	
		130				135					140					
GAC	CTA	TTA	GTG	GGC	GCG	GCC	ACG	ATG	TGC	TCT	GCG	CTC	TAC	GTG	GGT	480
Asp	Leu	Leu	Val	Gly	Ala	Ala	Thr	Met	Cys	Ser	Ala	Leu	Tyr	Val	Gly	
	145				150					155				160		
GAC	ATG	TGT	GGG	GCT	GTC	TTC	CTC	GTG	GGA	CAA	GCC	TTC	ACG	TTC	AGA	528
Asp	Met	Cys	Gly	Ala	Val	Phe	Leu	Val	Gly	Gln	Ala	Phe	Thr	Phe	Arg	
			165						170					175		
CCT	CGT	CGC	CAT	CAA	ACG	GTC	CAG	ACC	TGT	AAC	TGC	TCG	CTG	TAC	CCA	576
Pro	Arg	Arg	His	Gln	Thr	Val	Gln	Thr	Cys	Asn	Cys	Ser	Leu	Tyr	Pro	
			180					185					190			
GGC	CAT	CTT	TCA	GGA	CAT	CGA	ATG	GCT	TGG	GAT	ATG	ATG	ATG	AAC	TGG	624
Gly	His	Leu	Ser	Gly	His	Arg	Met	Ala	Trp	Asp	Met	Met	Met	Asn	Trp	
		195					200					205				
TAATAG																634

(2) INFORMATION FOR SEQ ID NO: 30:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 208 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30:

Met Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu Met  
 1 5 10 15  
 Gly Tyr Ile Pro Leu Val Gly Ala Pro Val Gly Gly Val Ala Arg Ala  
 20 25 30  
 Leu Ala His Gly Val Arg Ala Leu Glu Asp Gly Ile Asn Phe Ala Thr  
 35 40 45  
 Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu Phe  
 50 55 60  
 Ser Cys Leu Ile His Pro Ala Ala Ser Leu Glu Trp Arg Asn Thr Ser  
 65 70 75 80  
 Gly Leu Tyr Val Leu Thr Asn Asp Cys Ser Asn Ser Ser Ile Val Tyr  
 85 90 95  
 Glu Ala Asp Asp Val Ile Leu His Thr Pro Gly Cys Ile Pro Cys Val  
 100 105 110  
 Gln Asp Gly Asn Thr Ser Thr Cys Trp Thr Pro Val Thr Pro Thr Val  
 115 120 125  
 Ala Val Lys Tyr Val Gly Ala Thr Thr Ala Ser Ile Arg Ser His Val  
 130 135 140  
 Asp Leu Leu Val Gly Ala Ala Thr Met Cys Ser Ala Leu Tyr Val Gly  
 145 150 155 160  
 Asp Met Cys Gly Ala Val Phe Leu Val Gly Gln Ala Phe Thr Phe Arg  
 165 170 175  
 Pro Arg Arg His Gln Thr Val Gln Thr Cys Asn Cys Ser Leu Tyr Pro  
 180 185 190  
 Gly His Leu Ser Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp  
 195 200 205

## (2) INFORMATION FOR SEQ ID NO: 31:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 630 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: CDS  
(B) LOCATION: 1..627

## (ix) FEATURE:

(A) NAME/KEY: mat\_peptide  
(B) LOCATION: 1..624

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31:

ATG GGT AAG GTC ATC GAT ACC CTA ACG TGC GGA TTC GCC GAT CTC ATG	48
Met Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu Met	
1 5 10 15	
GGG TAT ATC CCG CTC GTA GGC GGC CCC ATT GGG GGC GTC GCA AGG GCT	96
Gly Tyr Ile Pro Leu Val Gly Gly Pro Ile Gly Gly Val Ala Arg Ala	
20 25 30	
CTC GCA CAC GGT GTG AGG GTC CTT GAG GAC GGG GTA AAC TAT GCA ACA	144
Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala Thr	
35 40 45	
GGG AAT TTA CCC GGT TGC TCT TTC TCT ATC TTT ATT CTT GCT CTT CTC	192
Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Ile Leu Ala Leu Leu	
50 55 60	
TCG TGT CTG ACC GTT CCG GCC TCT GCA GTT CCC TAC CGA AAT GCC TCT	240
Ser Cys Leu Thr Val Pro Ala Ser Ala Val Pro Tyr Arg Asn Ala Ser	
65 70 75 80	
GGG ATT TAT CAT GTT ACC AAT GAT TGC CCA AAC TCT TCC ATA GTC TAT	288
Gly Ile Tyr His Val Thr Asn Asp Cys Pro Asn Ser Ser Ile Val Tyr	
85 90 95	
GAG GCA GAT AAC CTG ATC CTA CAC GCA CCT GGT TGC GTG CCT TGT GTC	336
Glu Ala Asp Asn Leu Ile Leu His Ala Pro Gly Cys Val Pro Cys Val	
100 105 110	
ATG ACA GGT AAT GTG AGT AGA TGC TGG GTC CAA ATT ACC CCT ACA CTG	384
Met Thr Gly Asn Val Ser Arg Cys Trp Val Gln Ile Thr Pro Thr Leu	
115 120 125	
TCA GCC CCG AGC CTC GGA GCA GTC ACG GCT CCT CTT CGG AGA GCC GTT	432
Ser Ala Pro Ser Leu Gly Ala Val Thr Ala Pro Leu Arg Arg Ala Val	
130 135 140	
GAC TAC CTA GCG GGA GGG GCT GCC CTC TGC TCC GCG TTA TAC GTA GGA	480
Asp Tyr Leu Ala Gly Gly Ala Ala Leu Cys Ser Ala Leu Tyr Val Gly	
145 150 155 160	
GAC GCG TGT GGG GCA CTA TTC TTG GTA GGC CAA ATG TTC ACC TAT AGG	528
Asp Ala Cys Gly Ala Leu Phe Leu Val Gly Gln Met Phe Thr Tyr Arg	
165 170 175	
CCT CGC CAG CAC GCT ACG GTG CAG AAC TGC AAC TGT TCC ATT TAC AGT	576
Pro Arg Gln His Ala Thr Val Gln Asn Cys Asn Cys Ser Ile Tyr Ser	
180 185 190	
GGC CAT GTT ACC GGC CAC CGG ATG GCA TGG GAT ATG ATG ATG AAC TGG	624
Gly His Val Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp	
195 200 205	

TAATAG

630

## (2) INFORMATION FOR SEQ ID NO: 32:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 208 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32:

```

Met Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu Met
 1           5           10           15
Gly Tyr Ile Pro Leu Val Gly Gly Pro Ile Gly Gly Val Ala Arg Ala
 20           25           30
Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala Thr
 35           40           45
Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Ile Leu Ala Leu Leu
 50           55           60
Ser Cys Leu Thr Val Pro Ala Ser Ala Val Pro Tyr Arg Asn Ala Ser
 65           70           75           80
Gly Ile Tyr His Val Thr Asn Asp Cys Pro Asn Ser Ser Ile Val Tyr
 85           90           95
Glu Ala Asp Asn Leu Ile Leu His Ala Pro Gly Cys Val Pro Cys Val
100           105           110
Met Thr Gly Asn Val Ser Arg Cys Trp Val Gln Ile Thr Pro Thr Leu
115           120           125
Ser Ala Pro Ser Leu Gly Ala Val Thr Ala Pro Leu Arg Arg Ala Val
130           135           140
Asp Tyr Leu Ala Gly Gly Ala Ala Leu Cys Ser Ala Leu Tyr Val Gly
145           150           155           160
Asp Ala Cys Gly Ala Leu Phe Leu Val Gly Gln Met Phe Thr Tyr Arg
165           170           175
Pro Arg Gln His Ala Thr Val Gln Asn Cys Asn Cys Ser Ile Tyr Ser
180           185           190
Gly His Val Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp
195           200           205

```

## (2) INFORMATION FOR SEQ ID NO: 33:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23 base pairs

30

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33:  
TGGGATATGA TGATGAACTG GTC

23

(2) INFORMATION FOR SEQ ID NO: 34:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 30 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 34:  
CTATTATGGT GGTAAGCCAC AGAGCAGGAG

30

(2) INFORMATION FOR SEQ ID NO: 35:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1476 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 1..1473

- (ix) FEATURE:
  - (A) NAME/KEY: mat\_peptide
  - (B) LOCATION: 1..1470

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 35:  
TGG GAT ATG ATG ATG AAC TGG TCG CCT ACA ACG GCC CTG GTG GTA TCG

48

Trp	Asp	Met	Met	Met	Asn	Trp	Ser	Pro	Thr	Thr	Ala	Leu	Val	Val	Ser	
1				5					10					15		
CAG	CTG	CTC	CGG	ATC	CCA	CAA	GCT	GTC	GTG	GAC	ATG	GTG	GCG	GGG	GCC	96
Gln	Leu	Leu	Arg	Ile	Pro	Gln	Ala	Val	Val	Asp	Met	Val	Ala	Gly	Ala	
			20					25					30			
CAT	TGG	GGA	GTC	CTG	GCG	GGC	CTC	GCC	TAC	TAT	TCC	ATG	GTG	GGG	AAC	144
His	Trp	Gly	Val	Leu	Ala	Gly	Leu	Ala	Tyr	Tyr	Ser	Met	Val	Gly	Asn	
		35					40					45				
TGG	GCT	AAG	GTT	TTG	GTT	GTG	ATG	CTA	CTC	TTT	GCC	GGC	GTC	GAC	GGG	192
Trp	Ala	Lys	Val	Leu	Val	Val	Met	Leu	Leu	Phe	Ala	Gly	Val	Asp	Gly	
	50					55					60					
CAT	ACC	CGC	GTG	TCA	GGA	GGG	GCA	GCA	GCC	TCC	GAT	ACC	AGG	GGC	CTT	240
His	Thr	Arg	Val	Ser	Gly	Gly	Ala	Ala	Ala	Ser	Asp	Thr	Arg	Gly	Leu	
	65				70					75					80	
GTG	TCC	CTC	TTT	AGC	CCC	GGG	TCG	GCT	CAG	AAA	ATC	CAG	CTC	GTA	AAC	288
Val	Ser	Leu	Phe	Ser	Pro	Gly	Ser	Ala	Gln	Lys	Ile	Gln	Leu	Val	Asn	
				85					90					95		
ACC	AAC	GGC	AGT	TGG	CAC	ATC	AAC	AGG	ACT	GCC	CTG	AAC	TGC	AAC	GAC	336
Thr	Asn	Gly	Ser	Trp	His	Ile	Asn	Arg	Thr	Ala	Leu	Asn	Cys	Asn	Asp	
			100					105					110			
TCC	CTC	CAA	ACA	GGG	TTC	TTT	GCC	GCA	CTA	TTC	TAC	AAA	CAC	AAA	TTC	384
Ser	Leu	Gln	Thr	Gly	Phe	Phe	Ala	Ala	Leu	Phe	Tyr	Lys	His	Lys	Phe	
		115					120					125				
AAC	TCG	TCT	GGA	TGC	CCA	GAG	CGC	TTG	GCC	AGC	TGT	CGC	TCC	ATC	GAC	432
Asn	Ser	Ser	Gly	Cys	Pro	Glu	Arg	Leu	Ala	Ser	Cys	Arg	Ser	Ile	Asp	
	130					135					140					
AAG	TTT	GCT	CAG	GGG	TGG	GGT	CCC	CTC	ACT	TAC	ACT	GAG	CCT	AAC	AGC	480
Lys	Phe	Ala	Gln	Gly	Trp	Gly	Pro	Leu	Thr	Tyr	Thr	Glu	Pro	Asn	Ser	
	145				150					155					160	
TCG	GAC	CAG	AGG	CCC	TAC	TGC	TGG	CAC	TAC	GCG	CCT	CGA	CCG	TGT	GGT	528
Ser	Asp	Gln	Arg	Pro	Tyr	Cys	Trp	His	Tyr	Ala	Pro	Arg	Pro	Cys	Gly	
				165					170					175		
ATT	GTA	CCC	GCG	TCT	CAG	GTG	TGC	GGT	CCA	GTG	TAT	TGC	TTC	ACC	CCG	576
Ile	Val	Pro	Ala	Ser	Gln	Val	Cys	Gly	Pro	Val	Tyr	Cys	Phe	Thr	Pro	
			180					185					190			
AGC	CCT	GTT	GTG	GTG	GGG	ACG	ACC	GAT	CGG	TTT	GGT	GTC	CCC	ACG	TAT	624
Ser	Pro	Val	Val	Val	Gly	Thr	Thr	Asp	Arg	Phe	Gly	Val	Pro	Thr	Tyr	
		195					200					205				
AAC	TGG	GGG	GCG	AAC	GAC	TCG	GAT	GTG	CTG	ATT	CTC	AAC	AAC	ACG	CGG	672
Asn	Trp	Gly	Ala	Asn	Asp	Ser	Asp	Val	Leu	Ile	Leu	Asn	Asn	Thr	Arg	
	210					215					220					
CCG	CCG	CGA	GGC	AAC	TGG	TTC	GGC	TGT	ACA	TGG	ATG	AAT	GGC	ACT	GGG	720
Pro	Pro	Arg	Gly	Asn	Trp	Phe	Gly	Cys	Thr	Trp	Met	Asn	Gly	Thr	Gly	
	225				230					235				240		
TTC	ACC	AAG	ACG	TGT	GGG	GGC	CCC	CCG	TGC	AAC	ATC	GGG	GGG	GCC	GGC	768
Phe	Thr	Lys	Thr	Cys	Gly	Gly	Pro	Pro	Cys	Asn	Ile	Gly	Gly	Ala	Gly	
				245					250					255		



AAC AAC ACC TTG ACC TGC CCC ACT GAC TGT TTT CGG AAG CAC CCC GAG Asn Asn Thr Leu Thr Cys Pro Thr Asp Cys Phe Arg Lys His Pro Glu 260 265 270	816
GCC ACC TAC GCC AGA TGC GGT TCT GGG CCC TGG CTG ACA CCT AGG TGT Ala Thr Tyr Ala Arg Cys Gly Ser Gly Pro Trp Leu Thr Pro Arg Cys 275 280 285	864
ATG GTT CAT TAC CCA TAT AGG CTC TGG CAC TAC CCC TGC ACT GTC AAC Met Val His Tyr Pro Tyr Arg Leu Trp His Tyr Pro Cys Thr Val Asn 290 295 300	912
TTC ACC ATC TTC AAG GTT AGG ATG TAC GTG GGG GGC GTG GAG CAC AGG Phe Thr Ile Phe Lys Val Arg Met Tyr Val Gly Gly Val Glu His Arg 305 310 315 320	960
TTC GAA GCC GCA TGC AAT TGG ACT CGA GGA GAG CGT TGT GAC TTG GAG Phe Glu Ala Ala Cys Asn Trp Thr Arg Gly Glu Arg Cys Asp Leu Glu 325 330 335	1008
GAC AGG GAT AGA TCA GAG CTT AGC CCG CTG CTG CTG TCT ACA ACA GAG Asp Arg Asp Arg Ser Glu Leu Ser Pro Leu Leu Leu Ser Thr Thr Glu 340 345 350	1056
TGG CAG ATA CTG CCC TGT TCC TTC ACC ACC CTG CCG GCC CTA TCC ACC Trp Gln Ile Leu Pro Cys Ser Phe Thr Thr Leu Pro Ala Leu Ser Thr 355 360 365	1104
GGC CTG ATC CAC CTC CAT CAG AAC ATC GTG GAC GTG CAA TAC CTG TAC Gly Leu Ile His Leu His Gln Asn Ile Val Asp Val Gln Tyr Leu Tyr 370 375 380	1152
GGT GTA GGG TCG GCG GTT GTC TCC CTT GTC ATC AAA TGG GAG TAT GTC Gly Val Gly Ser Ala Val Val Ser Leu Val Ile Lys Trp Glu Tyr Val 385 390 395 400	1200
CTG TTG CTC TTC CTT CTC CTG GCA GAC GCG CGC ATC TGC GCC TGC TTA Leu Leu Leu Phe Leu Leu Leu Ala Asp Ala Arg Ile Cys Ala Cys Leu 405 410 415	1248
TGG ATG ATG CTG CTG ATA GCT CAA GCT GAG GCC GCC TTA GAG AAC CTG Trp Met Met Leu Leu Ile Ala Gln Ala Glu Ala Ala Leu Glu Asn Leu 420 425 430	1296
GTG GTC CTC AAT GCG GCG GCC GTG GCC GGG GCG CAT GGC ACT CTT TCC Val Val Leu Asn Ala Ala Ala Val Ala Gly Ala His Gly Thr Leu Ser 435 440 445	1344
TTC CTT GTG TTC TTC TGT GCT GCC TGG TAC ATC AAG GGC AGG CTG GTC Phe Leu Val Phe Phe Cys Ala Ala Trp Tyr Ile Lys Gly Arg Leu Val 450 455 460	1392
CCT GGT GCG GCA TAC GCC TTC TAT GGC GTG TGG CCG CTG CTC CTG CTT Pro Gly Ala Ala Tyr Ala Phe Tyr Gly Val Trp Pro Leu Leu Leu Leu 465 470 475 480	1440
CTG CTG GCC TTA CCA CCA CGA GCT TAT GCC TAGTAA Leu Leu Ala Leu Pro Pro Arg Ala Tyr Ala 485 490	1476

## (2) INFORMATION FOR SEQ ID NO: 36:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 490 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 36:

```

Trp Asp Met Met Met Asn Trp Ser Pro Thr Thr Ala Leu Val Val Ser
 1           5           10
Gln Leu Leu Arg Ile Pro Gln Ala Val Val Asp Met Val Ala Gly Ala
      20           25           30
His Trp Gly Val Leu Ala Gly Leu Ala Tyr Tyr Ser Met Val Gly Asn
      35           40           45
Trp Ala Lys Val Leu Val Val Met Leu Leu Phe Ala Gly Val Asp Gly
      50           55           60
His Thr Arg Val Ser Gly Gly Ala Ala Ala Ser Asp Thr Arg Gly Leu
      65           70           75           80
Val Ser Leu Phe Ser Pro Gly Ser Ala Gln Lys Ile Gln Leu Val Asn
      85           90           95
Thr Asn Gly Ser Trp His Ile Asn Arg Thr Ala Leu Asn Cys Asn Asp
      100          105          110
Ser Leu Gln Thr Gly Phe Phe Ala Ala Leu Phe Tyr Lys His Lys Phe
      115          120          125
Asn Ser Ser Gly Cys Pro Glu Arg Leu Ala Ser Cys Arg Ser Ile Asp
      130          135          140
Lys Phe Ala Gln Gly Trp Gly Pro Leu Thr Tyr Thr Glu Pro Asn Ser
      145          150          155          160
Ser Asp Gln Arg Pro Tyr Cys Trp His Tyr Ala Pro Arg Pro Cys Gly
      165          170          175
Ile Val Pro Ala Ser Gln Val Cys Gly Pro Val Tyr Cys Phe Thr Pro
      180          185          190
Ser Pro Val Val Val Gly Thr Thr Asp Arg Phe Gly Val Pro Thr Tyr
      195          200          205
Asn Trp Gly Ala Asn Asp Ser Asp Val Leu Ile Leu Asn Asn Thr Arg
      210          215          220
Pro Pro Arg Gly Asn Trp Phe Gly Cys Thr Trp Met Asn Gly Thr Gly
      225          230          235          240
Phe Thr Lys Thr Cys Gly Gly Pro Pro Cys Asn Ile Gly Gly Ala Gly
      245          250          255
Asn Asn Thr Leu Thr Cys Pro Thr Asp Cys Phe Arg Lys His Pro Glu
      260          265          270
Ala Thr Tyr Ala Arg Cys Gly Ser Gly Pro Trp Leu Thr Pro Arg Cys

```

275	280	285
Met Val His Tyr Pro Tyr	Arg Leu Trp His Tyr	Pro Cys Thr Val Asn
290	295	300
Phe Thr Ile Phe Lys Val	Arg Met Tyr Val Gly Gly Val	Glu His Arg
305	310	315 320
Phe Glu Ala Ala Cys Asn Trp Thr Arg Gly Glu Arg Cys Asp Leu Glu		
	325	330 335
Asp Arg Asp Arg Ser Glu Leu Ser Pro Leu Leu Leu Ser Thr Thr Glu		
	340	345 350
Trp Gln Ile Leu Pro Cys Ser Phe Thr Thr Leu Pro Ala Leu Ser Thr		
	355	360 365
Gly Leu Ile His Leu His Gln Asn Ile Val Asp Val Gln Tyr Leu Tyr		
	370	375 380
Gly Val Gly Ser Ala Val Val Ser Leu Val Ile Lys Trp Glu Tyr Val		
	385	390 395 400
Leu Leu Leu Phe Leu Leu Leu Ala Asp Ala Arg Ile Cys Ala Cys Leu		
	405	410 415
Trp Met Met Leu Leu Ile Ala Gln Ala Glu Ala Ala Leu Glu Asn Leu		
	420	425 430
Val Val Leu Asn Ala Ala Ala Val Ala Gly Ala His Gly Thr Leu Ser		
	435	440 445
Phe Leu Val Phe Phe Cys Ala Ala Trp Tyr Ile Lys Gly Arg Leu Val		
	450	455 460
Pro Gly Ala Ala Tyr Ala Phe Tyr Gly Val Trp Pro Leu Leu Leu Leu		
	465	470 475 480
Leu Leu Ala Leu Pro Pro Arg Ala Tyr Ala		
	485	490

## (2) INFORMATION FOR SEQ ID NO: 37:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1021 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: cDNA

- (iii) HYPOTHETICAL: NO

- (iii) ANTI-SENSE: NO

- (ix) FEATURE:

- (A) NAME/KEY: CDS
  - (B) LOCATION: 2..1018

- (ix) FEATURE:

- (A) NAME/KEY: mat\_peptide

(B) LOCATION: 2..1015

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 37:

G ATC CCA CAA GCT GTC GTG GAC ATG GTG GCG GGG GCC CAT TGG GGA	46
Ile Pro Gln Ala Val Val Asp Met Val Ala Gly Ala His Trp Gly	
1 5 10 15	
GTC CTG GCG GGC CTC GCC TAC TAT TCC ATG GTG GGG AAC TGG GCT AAG	94
Val Leu Ala Gly Leu Ala Tyr Tyr Ser Met Val Gly Asn Trp Ala Lys	
20 25 30	
GTT TTG GTT GTG ATG CTA CTC TTT GCC GGC GTC GAC GGG CAT ACC CGC	142
Val Leu Val Val Met Leu Leu Phe Ala Gly Val Asp Gly His Thr Arg	
35 40 45	
GTG TCA GGA GGG GCA GCA GCC TCC GAT ACC AGG GGC CTT GTG TCC CTC	190
Val Ser Gly Gly Ala Ala Ala Ser Asp Thr Arg Gly Leu Val Ser Leu	
50 55 60	
TTT AGC CCC GGG TCG GCT CAG AAA ATC CAG CTC GTA AAC ACC AAC GGC	238
Phe Ser Pro Gly Ser Ala Gln Lys Ile Gln Leu Val Asn Thr Asn Gly	
65 70 75	
AGT TGG CAC ATC AAC AGG ACT GCC CTG AAC TGC AAC GAC TCC CTC CAA	286
Ser Trp His Ile Asn Arg Thr Ala Leu Asn Cys Asn Asp Ser Leu Gln	
80 85 90 95	
ACA GGG TTC TTT GCC GCA CTA TTC TAC AAA CAC AAA TTC AAC TCG TCT	334
Thr Gly Phe Phe Ala Ala Leu Phe Tyr Lys His Lys Phe Asn Ser Ser	
100 105 110	
GGA TGC CCA GAG CGC TTG GCC AGC TGT CGC TCC ATC GAC AAG TTC GCT	382
Gly Cys Pro Glu Arg Leu Ala Ser Cys Arg Ser Ile Asp Lys Phe Ala	
115 120 125	
CAG GGG TGG GGT CCC CTC ACT TAC ACT GAG CCT AAC AGC TCG GAC CAG	430
Gln Gly Trp Gly Pro Leu Thr Tyr Thr Glu Pro Asn Ser Ser Asp Gln	
130 135 140	
AGG CCC TAC TGC TGG CAC TAC GCG CCT CGA CCG TGT GGT ATT GTA CCC	478
Arg Pro Tyr Cys Trp His Tyr Ala Pro Arg Pro Cys Gly Ile Val Pro	
145 150 155	
GCG TCT CAG GTG TGC GGT CCA GTG TAT TGC TTC ACC CCG AGC CCT GTT	526
Ala Ser Gln Val Cys Gly Pro Val Tyr Cys Phe Thr Pro Ser Pro Val	
160 165 170 175	
GTG GTG GGG ACG ACC GAT CGG TTT GGT GTC CCC ACG TAT AAC TGG GGG	574
Val Val Gly Thr Thr Asp Arg Phe Gly Val Pro Thr Tyr Asn Trp Gly	
180 185 190	
GCG AAC GAC TCG GAT GTG CTG ATT CTC AAC AAC ACG CGG CCG CCG CGA	622
Ala Asn Asp Ser Asp Val Leu Ile Leu Asn Asn Thr Arg Pro Pro Arg	
195 200 205	
GGC AAC TGG TTC GGC TGT ACA TGG ATG AAT GGC ACT GGG TTC ACC AAG	670
Gly Asn Trp Phe Gly Cys Thr Trp Met Asn Gly Thr Gly Phe Thr Lys	
210 215 220	
ACG TGT GGG GGC CCC CCG TGC AAC ATC GGG GGG GCC GGC AAC AAC ACC	718
Thr Cys Gly Gly Pro Pro Cys Asn Ile Gly Gly Ala Gly Asn Asn Thr	

225	230	235	
TTG ACC TGC CCC ACT GAC TGT TTT CGG AAG CAC CCC GAG GCC ACC TAC			766
Leu Thr Cys Pro Thr Asp Cys Phe Arg Lys His Pro Glu Ala Thr Tyr			
240	245	250	255
GCC AGA TGC GGT TCT GGG CCC TGG CTG ACA CCT AGG TGT ATG GTT CAT			814
Ala Arg Cys Gly Ser Gly Pro Trp Leu Thr Pro Arg Cys Met Val His			
	260	265	270
TAC CCA TAT AGG CTC TGG CAC TAC CCC TGC ACT GTC AAC TTC ACC ATC			862
Tyr Pro Tyr Arg Leu Trp His Tyr Pro Cys Thr Val Asn Phe Thr Ile			
	275	280	285
TTC AAG GTT AGG ATG TAC GTG GGG GGC GTG GAG CAC AGG TTC GAA GCC			910
Phe Lys Val Arg Met Tyr Val Gly Gly Val Glu His Arg Phe Glu Ala			
	290	295	300
GCA TGC AAT TGG ACT CGA GGA GAG CGT TGT GAC TTG GAG GAC AGG GAT			958
Ala Cys Asn Trp Thr Arg Gly Glu Arg Cys Asp Leu Glu Asp Arg Asp			
	305	310	315
AGA TCA GAG CTT AGC CCG CTG CTG CTG TCT ACA ACA GAG TGG CAG AGT			1006
Arg Ser Glu Leu Ser Pro Leu Leu Leu Ser Thr Thr Glu Trp Gln Ser			
	320	325	330
GGC AGA GCT TAATTA			1021
Gly Arg Ala			

## (2) INFORMATION FOR SEQ ID NO: 38:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 338 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 38:

Ile Pro Gln Ala Val Val Asp Met Val Ala Gly Ala His Trp Gly Val  
 1 5 10 15  
 Leu Ala Gly Leu Ala Tyr Tyr Ser Met Val Gly Asn Trp Ala Lys Val  
 20 25 30  
 Leu Val Val Met Leu Leu Phe Ala Gly Val Asp Gly His Thr Arg Val  
 35 40 45  
 Ser Gly Gly Ala Ala Ala Ser Asp Thr Arg Gly Leu Val Ser Leu Phe  
 50 55 60  
 Ser Pro Gly Ser Ala Gln Lys Ile Gln Leu Val Asn Thr Asn Gly Ser  
 65 70 75 80  
 Trp His Ile Asn Arg Thr Ala Leu Asn Cys Asn Asp Ser Leu Gln Thr  
 85 90 95  
 Gly Phe Phe Ala Ala Leu Phe Tyr Lys His Lys Phe Asn Ser Ser Gly  
 100 105 110

Cys Pro Glu Arg Leu Ala Ser Cys Arg Ser Ile Asp Lys Phe Ala Gln  
           115                          120                          125  
 Gly Trp Gly Pro Leu Thr Tyr Thr Glu Pro Asn Ser Ser Asp Gln Arg  
           130                          135                          140  
 Pro Tyr Cys Trp His Tyr Ala Pro Arg Pro Cys Gly Ile Val Pro Ala  
   145                          150                          155                          160  
 Ser Gln Val Cys Gly Pro Val Tyr Cys Phe Thr Pro Ser Pro Val Val  
                           165                          170                          175  
 Val Gly Thr Thr Asp Arg Phe Gly Val Pro Thr Tyr Asn Trp Gly Ala  
                           180                          185                          190  
 Asn Asp Ser Asp Val Leu Ile Leu Asn Asn Thr Arg Pro Pro Arg Gly  
           195                          200                          205  
 Asn Trp Phe Gly Cys Thr Trp Met Asn Gly Thr Gly Phe Thr Lys Thr  
           210                          215                          220  
 Cys Gly Gly Pro Pro Cys Asn Ile Gly Gly Ala Gly Asn Asn Thr Leu  
   225                          230                          235                          240  
 Thr Cys Pro Thr Asp Cys Phe Arg Lys His Pro Glu Ala Thr Tyr Ala  
                           245                          250                          255  
 Arg Cys Gly Ser Gly Pro Trp Leu Thr Pro Arg Cys Met Val His Tyr  
           260                          265                          270  
 Pro Tyr Arg Leu Trp His Tyr Pro Cys Thr Val Asn Phe Thr Ile Phe  
           275                          280                          285  
 Lys Val Arg Met Tyr Val Gly Gly Val Glu His Arg Phe Glu Ala Ala  
           290                          295                          300  
 Cys Asn Trp Thr Arg Gly Glu Arg Cys Asp Leu Glu Asp Arg Asp Arg  
   305                          310                          315                          320  
 Ser Glu Leu Ser Pro Leu Leu Leu Ser Thr Thr Glu Trp Gln Ser Gly  
           325                          330                          335  
 Arg Ala

## (2) INFORMATION FOR SEQ ID NO: 39:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1034 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: CDS  
(B) LOCATION: 2..1032

(ix) FEATURE:

(A) NAME/KEY: mat\_peptide  
(B) LOCATION: 2..1029

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 39:

G ATC CCA CAA GCT GTC GTG GAC ATG GTG GCG GGG GCC CAT TGG GGA	46
Ile Pro Gln Ala Val Val Asp Met Val Ala Gly Ala His Trp Gly	
1 5 10 15	
GTC CTG GCG GGC CTC GCC TAC TAT TCC ATG GTG GGG AAC TGG GCT AAG	94
Val Leu Ala Gly Leu Ala Tyr Tyr Ser Met Val Gly Asn Trp Ala Lys	
20 25 30	
GTT TTG GTT GTG ATG CTA CTC TTT GCC GGC GTC GAC GGG CAT ACC CGC	142
Val Leu Val Val Met Leu Leu Phe Ala Gly Val Asp Gly His Thr Arg	
35 40 45	
GTG TCA GGA GGG GCA GCA GCC TCC GAT ACC AGG GGC CTT GTG TCC CTC	190
Val Ser Gly Gly Ala Ala Ala Ser Asp Thr Arg Gly Leu Val Ser Leu	
50 55 60	
TTT AGC CCC GGG TCG GCT CAG AAA ATC CAG CTC GTA AAC ACC AAC GGC	238
Phe Ser Pro Gly Ser Ala Gln Lys Ile Gln Leu Val Asn Thr Asn Gly	
65 70 75	
AGT TGG CAC ATC AAC AGG ACT GCC CTG AAC TGC AAC GAC TCC CTC CAA	286
Ser Trp His Ile Asn Arg Thr Ala Leu Asn Cys Asn Asp Ser Leu Gln	
80 85 90 95	
ACA GGG TTC TTT GCC GCA CTA TTC TAC AAA CAC AAA TTC AAC TCG TCT	334
Thr Gly Phe Phe Ala Ala Leu Phe Tyr Lys His Lys Phe Asn Ser Ser	
100 105 110	
GGA TGC CCA GAG CGC TTG GCC AGC TGT CGC TCC ATC GAC AAG TTC GCT	382
Gly Cys Pro Glu Arg Leu Ala Ser Cys Arg Ser Ile Asp Lys Phe Ala	
115 120 125	
CAG GGG TGG GGT CCC CTC ACT TAC ACT GAG CCT AAC AGC TCG GAC CAG	430
Gln Gly Trp Gly Pro Leu Thr Tyr Thr Glu Pro Asn Ser Ser Asp Gln	
130 135 140	
AGG CCC TAC TGC TGG CAC TAC GCG CCT CGA CCG TGT GGT ATT GTA CCC	478
Arg Pro Tyr Cys Trp His Tyr Ala Pro Arg Pro Cys Gly Ile Val Pro	
145 150 155	
GCG TCT CAG GTG TGC GGT CCA GTG TAT TGC TTC ACC CCG AGC CCT GTT	526
Ala Ser Gln Val Cys Gly Pro Val Tyr Cys Phe Thr Pro Ser Pro Val	
160 165 170 175	
GTG GTG GGG ACG ACC GAT CGG TTT GGT GTC CCC ACG TAT AAC TGG GGG	574
Val Val Gly Thr Thr Asp Arg Phe Gly Val Pro Thr Tyr Asn Trp Gly	
180 185 190	
GCG AAC GAC TCG GAT GTG CTG ATT CTC AAC AAC ACG CGG CCG CCG CGA	622
Ala Asn Asp Ser Asp Val Leu Ile Leu Asn Asn Thr Arg Pro Pro Arg	
195 200 205	

GGC AAC TGG TTC GGC TGT ACA TGG ATG AAT GGC ACT GGG TTC ACC AAG Gly Asn Trp Phe Gly Cys Thr Trp Met Asn Gly Thr Gly Phe Thr Lys 210 215 220	670
ACG TGT GGG GGC CCC CCG TGC AAC ATC GGG GGG GCC GGC AAC AAC ACC Thr Cys Gly Gly Pro Pro Cys Asn Ile Gly Gly Ala Gly Asn Asn Thr 225 230 235	718
TTG ACC TGC CCC ACT GAC TGT TTT CGG AAG CAC CCC GAG GCC ACC TAC Leu Thr Cys Pro Thr Asp Cys Phe Arg Lys His Pro Glu Ala Thr Tyr 240 245 250 255	766
GCC AGA TGC GGT TCT GGG CCC TGG CTG ACA CCT AGG TGT ATG GTT CAT Ala Arg Cys Gly Ser Gly Pro Trp Leu Thr Pro Arg Cys Met Val His 260 265 270	814
TAC CCA TAT AGG CTC TGG CAC TAC CCC TGC ACT GTC AAC TTC ACC ATC Tyr Pro Tyr Arg Leu Trp His Tyr Pro Cys Thr Val Asn Phe Thr Ile 275 280 285	862
TTC AAG GTT AGG ATG TAC GTG GGG GGC GTG GAG CAC AGG TTC GAA GCC Phe Lys Val Arg Met Tyr Val Gly Gly Val Glu His Arg Phe Glu Ala 290 295 300	910
GCA TGC AAT TGG ACT CGA GGA GAG CGT TGT GAC TTG GAG GAC AGG GAT Ala Cys Asn Trp Thr Arg Gly Glu Arg Cys Asp Leu Glu Asp Arg Asp 305 310 315	958
AGA TCA GAG CTT AGC CCG CTG CTG CTG TCT ACA ACA GGT GAT CGA GGG Arg Ser Glu Leu Ser Pro Leu Leu Leu Ser Thr Thr Gly Asp Arg Gly 320 325 330 335	1006
CAG ACA CCA TCA CCA CCA TCA CTA AT AG Gln Thr Pro Ser Pro Pro Ser Leu 340	1034

## (2) INFORMATION FOR SEQ ID NO: 40:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 343 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 40:

Ile Pro Gln Ala Val Val Asp Met Val Ala Gly Ala His Trp Gly Val 1 5 10 15
Leu Ala Gly Leu Ala Tyr Tyr Ser Met Val Gly Asn Trp Ala Lys Val 20 25 30
Leu Val Val Met Leu Leu Phe Ala Gly Val Asp Gly His Thr Arg Val 35 40 45
Ser Gly Gly Ala Ala Ala Ser Asp Thr Arg Gly Leu Val Ser Leu Phe 50 55 60
Ser Pro Gly Ser Ala Gln Lys Ile Gln Leu Val Asn Thr Asn Gly Ser 65 70 75 80



Trp	His	Ile	Asn	Arg 85	Thr	Ala	Leu	Asn	Cys 90	Asn	Asp	Ser	Leu	Gln	Thr 95
Gly	Phe	Phe	Ala 100	Ala	Leu	Phe	Tyr	Lys 105	His	Lys	Phe	Asn	Ser	Ser	Gly
Cys	Pro	Glu	Arg 115	Leu	Ala	Ser	Cys 120	Arg	Ser	Ile	Asp	Lys 125	Phe	Ala	Gln
Gly	Trp	Gly	Pro	Leu	Thr	Tyr	Thr 135	Glu	Pro	Asn	Ser	Ser	Asp	Gln	Arg
Pro 145	Tyr	Cys	Trp	His	Tyr 150	Ala	Pro	Arg	Pro	Cys 155	Gly	Ile	Val	Pro	Ala 160
Ser	Gln	Val	Cys 165	Gly	Pro	Val	Tyr	Cys 170	Phe	Thr	Pro	Ser	Pro	Val 175	Val
Val	Gly	Thr	Thr 180	Asp	Arg	Phe	Gly	Val 185	Pro	Thr	Tyr	Asn	Trp	Gly	Ala
Asn	Asp	Ser 195	Asp	Val	Leu	Ile	Leu 200	Asn	Asn	Thr	Arg	Pro 205	Pro	Arg	Gly
Asn	Trp 210	Phe	Gly	Cys	Thr	Trp 215	Met	Asn	Gly	Thr	Gly 220	Phe	Thr	Lys	Thr
Cys 225	Gly	Gly	Pro	Pro	Cys 230	Asn	Ile	Gly	Gly	Ala 235	Gly	Asn	Asn	Thr	Leu 240
Thr	Cys	Pro	Thr	Asp 245	Cys	Phe	Arg	Lys	His 250	Pro	Glu	Ala	Thr	Tyr 255	Ala
Arg	Cys	Gly	Ser 260	Gly	Pro	Trp	Leu	Thr 265	Pro	Arg	Cys	Met	Val	His	Tyr
Pro	Tyr 275	Arg	Leu	Trp	His	Tyr	Pro 280	Cys	Thr	Val	Asn	Phe 285	Thr	Ile	Phe
Lys 290	Val	Arg	Met	Tyr	Val	Gly 295	Gly	Val	Glu	His 300	Arg	Phe	Glu	Ala	Ala
Cys 305	Asn	Trp	Thr	Arg	Gly 310	Glu	Arg	Cys	Asp	Leu 315	Glu	Asp	Arg	Asp	Arg
Ser	Glu	Leu	Ser	Pro 325	Leu	Leu	Leu	Ser	Thr 330	Thr	Gly	Asp	Arg	Gly 335	Gln
Thr	Pro	Ser	Pro 340	Pro	Ser	Leu									

(2) INFORMATION FOR SEQ ID NO: 41:

- ```
(i) SEQUENCE CHARACTERISTICS:
      (A) LENGTH: 945 base pairs
      (B) TYPE: nucleic acid
      (C) STRANDEDNESS: single
      (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO
```

(iii) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..942

(ix) FEATURE:

(A) NAME/KEY: mat\_peptide

(B) LOCATION: 1..939

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 41:

|                                                                 |     |
|-----------------------------------------------------------------|-----|
| ATG GTG GGG AAC TGG GCT AAG GTT TTG GTT GTG ATG CTA CTC TTT GCC | 48  |
| Met Val Gly Asn Trp Ala Lys Val Leu Val Val Met Leu Leu Phe Ala |     |
| 1 5 10 15                                                       |     |
| GGC GTC GAC GGG CAT ACC CGC GTG TCA GGA GGG GCA GCA GCC TCC GAT | 96  |
| Gly Val Asp Gly His Thr Arg Val Ser Gly Gly Ala Ala Ala Ser Asp |     |
| 20 25 30                                                        |     |
| ACC AGG GGC CTT GTG TCC CTC TTT AGC CCC GGG TCG GCT CAG AAA ATC | 144 |
| Thr Arg Gly Leu Val Ser Leu Phe Ser Pro Gly Ser Ala Gln Lys Ile |     |
| 35 40 45                                                        |     |
| CAG CTC GTA AAC ACC AAC GGC AGT TGG CAC ATC AAC AGG ACT GCC CTG | 192 |
| Gln Leu Val Asn Thr Asn Gly Ser Trp His Ile Asn Arg Thr Ala Leu |     |
| 50 55 60                                                        |     |
| AAC TGC AAC GAC TCC CTC CAA ACA GGG TTC TTT GCC GCA CTA TTC TAC | 240 |
| Asn Cys Asn Asp Ser Leu Gln Thr Gly Phe Phe Ala Ala Leu Phe Tyr |     |
| 65 70 75 80                                                     |     |
| AAA CAC AAA TTC AAC TCG TCT GGA TGC CCA GAG CGC TTG GCC AGC TGT | 288 |
| Lys His Lys Phe Asn Ser Ser Gly Cys Pro Glu Arg Leu Ala Ser Cys |     |
| 85 90 95                                                        |     |
| CGC TCC ATC GAC AAG TTC GCT CAG GGG TGG GGT CCC CTC ACT TAC ACT | 336 |
| Arg Ser Ile Asp Lys Phe Ala Gln Gly Trp Gly Pro Leu Thr Tyr Thr |     |
| 100 105 110                                                     |     |
| GAG CCT AAC AGC TCG GAC CAG AGG CCC TAC TGC TGG CAC TAC GCG CCT | 384 |
| Glu Pro Asn Ser Ser Asp Gln Arg Pro Tyr Cys Trp His Tyr Ala Pro |     |
| 115 120 125                                                     |     |
| CGA CCG TGT GGT ATT GTA CCC GCG TCT CAG GTG TGC GGT CCA GTG TAT | 432 |
| Arg Pro Cys Gly Ile Val Pro Ala Ser Gln Val Cys Gly Pro Val Tyr |     |
| 130 135 140                                                     |     |
| TGC TTC ACC CCG AGC CCT GTT GTG GTG GGG ACG ACC GAT CGG TTT GGT | 480 |
| Cys Phe Thr Pro Ser Pro Val Val Val Gly Thr Thr Asp Arg Phe Gly |     |
| 145 150 155 160                                                 |     |
| GTC CCC ACG TAT AAC TGG GGG GCG AAC GAC TCG GAT GTG CTG ATT CTC | 528 |
| Val Pro Thr Tyr Asn Trp Gly Ala Asn Asp Ser Asp Val Leu Ile Leu |     |
| 165 170 175                                                     |     |
| AAC AAC ACG CGG CCG CCG CGA GGC AAC TGG TTC GGC TGT ACA TGG ATG | 576 |
| Asn Asn Thr Arg Pro Pro Arg Gly Asn Trp Phe Gly Cys Thr Trp Met |     |
| 180 185 190                                                     |     |

```

AAT GGC ACT GGG TTC ACC AAG ACG TGT GGG GGC CCC CCG TGC AAC ATC      624
Asn Gly Thr Gly Phe Thr Lys Thr Cys Gly Gly Pro Pro Cys Asn Ile
      195                      200                      205

GGG GGG GCC GGC AAC AAC ACC TTG ACC TGC CCC ACT GAC TGT TTT CGG      672
Gly Gly Ala Gly Asn Asn Thr Leu Thr Cys Pro Thr Asp Cys Phe Arg
      210                      215                      220

AAG CAC CCC GAG GCC ACC TAC GCC AGA TGC GGT TCT GGG CCC TGG CTG      720
Lys His Pro Glu Ala Thr Tyr Ala Arg Cys Gly Ser Gly Pro Trp Leu
      225                      230                      235

ACA CCT AGG TGT ATG GTT CAT TAC CCA TAT AGG CTC TGG CAC TAC CCC      768
Thr Pro Arg Cys Met Val His Tyr Pro Tyr Arg Leu Trp His Tyr Pro
      245                      250                      255

TGC ACT GTC AAC TTC ACC ATC TTC AAG GTT AGG ATG TAC GTG GGG GGC      816
Cys Thr Val Asn Phe Thr Ile Phe Lys Val Arg Met Tyr Val Gly Gly
      260                      265                      270

GTG GAG CAC AGG TTC GAA GCC GCA TGC AAT TGG ACT CGA GGA GAG CGT      864
Val Glu His Arg Phe Glu Ala Ala Cys Asn Trp Thr Arg Gly Glu Arg
      275                      280                      285

TGT GAC TTG GAG GAC AGG GAT AGA TCA GAG CTT AGC CCG CTG CTG CTG      912
Cys Asp Leu Glu Asp Arg Asp Arg Ser Glu Leu Ser Pro Leu Leu Leu
      290                      295                      300

TCT ACA ACA GAG TGG CAG AGC TTA ATT AAT TAG      945
Ser Thr Thr Glu Trp Gln Ser Leu Ile Asn
      305                      310

```

## (2) INFORMATION FOR SEQ ID NO: 42:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 314 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 42:

```

Met Val Gly Asn Trp Ala Lys Val Leu Val Val Met Leu Leu Phe Ala
  1           5           10           15

Gly Val Asp Gly His Thr Arg Val Ser Gly Gly Ala Ala Ala Ser Asp
      20           25           30

Thr Arg Gly Leu Val Ser Leu Phe Ser Pro Gly Ser Ala Gln Lys Ile
      35           40           45

Gln Leu Val Asn Thr Asn Gly Ser Trp His Ile Asn Arg Thr Ala Leu
      50           55           60

Asn Cys Asn Asp Ser Leu Gln Thr Gly Phe Phe Ala Ala Leu Phe Tyr
      65           70           75           80

Lys His Lys Phe Asn Ser Ser Gly Cys Pro Glu Arg Leu Ala Ser Cys
      85           90           95

```

43

```

Arg Ser Ile Asp Lys Phe Ala Gln Gly Trp Gly Pro Leu Thr Tyr Thr
      100                      105                      110

Glu Pro Asn Ser Ser Asp Gln Arg Pro Tyr Cys Trp His Tyr Ala Pro
      115                      120                      125

Arg Pro Cys Gly Ile Val Pro Ala Ser Gln Val Cys Gly Pro Val Tyr
      130                      135                      140

Cys Phe Thr Pro Ser Pro Val Val Val Gly Thr Thr Asp Arg Phe Gly
      145                      150                      155                      160

Val Pro Thr Tyr Asn Trp Gly Ala Asn Asp Ser Asp Val Leu Ile Leu
      165                      170                      175

Asn Asn Thr Arg Pro Pro Arg Gly Asn Trp Phe Gly Cys Thr Trp Met
      180                      185                      190

Asn Gly Thr Gly Phe Thr Lys Thr Cys Gly Gly Pro Pro Cys Asn Ile
      195                      200                      205

Gly Gly Ala Gly Asn Asn Thr Leu Thr Cys Pro Thr Asp Cys Phe Arg
      210                      215                      220

Lys His Pro Glu Ala Thr Tyr Ala Arg Cys Gly Ser Gly Pro Trp Leu
      225                      230                      235                      240

Thr Pro Arg Cys Met Val His Tyr Pro Tyr Arg Leu Trp His Tyr Pro
      245                      250                      255

Cys Thr Val Asn Phe Thr Ile Phe Lys Val Arg Met Tyr Val Gly Gly
      260                      265                      270

Val Glu His Arg Phe Glu Ala Ala Cys Asn Trp Thr Arg Gly Glu Arg
      275                      280                      285

Cys Asp Leu Glu Asp Arg Asp Arg Ser Glu Leu Ser Pro Leu Leu Leu
      290                      295                      300

Ser Thr Thr Glu Trp Gln Ser Leu Ile Asn
      305                      310

```

## (2) INFORMATION FOR SEQ ID NO: 43:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 961 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..958

## (ix) FEATURE:

- (A) NAME/KEY: mat\_peptide

(B) LOCATION: 1..955

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 43:

|                                                                 |     |
|-----------------------------------------------------------------|-----|
| ATG GTG GGG AAC TGG GCT AAG GCT TGG GTT GTG ATG CTA CTC TTT GCC | 48  |
| Met Val Gly Asn Trp Ala Lys Val Leu Val Val Met Leu Leu Phe Ala |     |
| 1 5 10 15                                                       |     |
| GGC GTC GAC GGG CAT ACC CGC GTG TCA GGA GGG GCA GCA GCC TCC GAT | 96  |
| Gly Val Asp Gly His Thr Arg Val Ser Gly Gly Ala Ala Ala Ser Asp |     |
| 20 25 30                                                        |     |
| ACC AGG GGC CTT GTG TCC CTC TTT AGC CCC GGG TCG GCT CAG AAA ATC | 144 |
| Thr Arg Gly Leu Val Ser Leu Phe Ser Pro Gly Ser Ala Gln Lys Ile |     |
| 35 40 45                                                        |     |
| CAG CTC GTA AAC ACC AAC GGC AGT TGG CAC ATC AAC AGG ACT GCC CTG | 192 |
| Gln Leu Val Asn Thr Asn Gly Ser Trp His Ile Asn Arg Thr Ala Leu |     |
| 50 55 60                                                        |     |
| AAC TGC AAC GAC TCC CTC CAA ACA GGG TTC TTT GCC GCA CTA TTC TAC | 240 |
| Asn Cys Asn Asp Ser Leu Gln Thr Gly Phe Phe Ala Ala Leu Phe Tyr |     |
| 65 70 75 80                                                     |     |
| AAA CAC AAA TTC AAC TCG TCT GGA TGC CCA GAG CGC TTG GCC AGC TGT | 288 |
| Lys His Lys Phe Asn Ser Ser Gly Cys Pro Glu Arg Leu Ala Ser Cys |     |
| 85 90 95                                                        |     |
| CGC TCC ATC GAC AAG TTC GCT CAG GGG TGG GGT CCC CTC ACT TAC ACT | 336 |
| Arg Ser Ile Asp Lys Phe Ala Gln Gly Trp Gly Pro Leu Thr Tyr Thr |     |
| 100 105 110                                                     |     |
| GAG CCT AAC AGC TCG GAC CAG AGG CCC TAC TGC TGG CAC TAC GCG CCT | 384 |
| Glu Pro Asn Ser Ser Asp Gln Arg Pro Tyr Cys Trp His Tyr Ala Pro |     |
| 115 120 125                                                     |     |
| CGA CCG TGT GGT ATT GTA CCC GCG TCT CAG GTG TGC GGT CCA GTG TAT | 432 |
| Arg Pro Cys Gly Ile Val Pro Ala Ser Gln Val Cys Gly Pro Val Tyr |     |
| 130 135 140                                                     |     |
| TGC TTC ACC CCG AGC CCT GTT GTG GTG GGG ACG ACC GAT CGG TTT GGT | 480 |
| Cys Phe Thr Pro Ser Pro Val Val Val Gly Thr Thr Asp Arg Phe Gly |     |
| 145 150 155 160                                                 |     |
| GTC CCC ACG TAT AAC TGG GGG GCG AAC GAC TCG GAT GTG CTG ATT CTC | 528 |
| Val Pro Thr Tyr Asn Trp Gly Ala Asn Asp Ser Asp Val Leu Ile Leu |     |
| 165 170 175                                                     |     |
| AAC AAC ACG CGG CCG CCG CGA GCG AAC TGG TTC GGC TGT ACA TGG ATG | 576 |
| Asn Asn Thr Arg Pro Pro Arg Gly Asn Trp Phe Gly Cys Thr Trp Met |     |
| 180 185 190                                                     |     |
| AAT GGC ACT GGG TTC ACC AAG ACG TGT GGG GGC CCC CCG TGC AAC ATC | 624 |
| Asn Gly Thr Gly Phe Thr Lys Thr Cys Gly Gly Pro Pro Cys Asn Ile |     |
| 195 200 205                                                     |     |
| GGG GGG GCC GGC AAC AAC ACC TTG ACC TGC CCC ACT GAC TGT TTT CGG | 672 |
| Gly Gly Ala Gly Asn Asn Thr Leu Thr Cys Pro Thr Asp Cys Phe Arg |     |
| 210 215 220                                                     |     |
| AAG CAC CCC GAG GCC ACC TAC GCC AGA TGC GGT TCT GGG CCC TGG CTG | 720 |

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |  |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|-----|
| Lys | His | Pro | Glu | Ala | Thr | Tyr | Ala | Arg | Cys | Gly | Ser | Gly | Pro | Trp | Leu |  |     |
| 225 |     |     |     |     | 230 |     |     |     |     | 235 |     |     |     |     | 240 |  |     |
| ACA | CCT | AGG | TGT | ATG | GTT | CAT | TAC | CCA | TAT | AGG | CTC | TGG | CAC | TAC | CCC |  | 768 |
| Thr | Pro | Arg | Cys | Met | Val | His | Tyr | Pro | Tyr | Arg | Leu | Trp | His | Tyr | Pro |  |     |
|     |     |     |     | 245 |     |     |     |     | 250 |     |     |     |     | 255 |     |  |     |
| TGC | ACT | GTC | AAC | TTC | ACC | ATC | TTC | AAG | GTT | AGG | ATG | TAC | GTG | GGG | GGC |  | 816 |
| Cys | Thr | Val | Asn | Phe | Thr | Ile | Phe | Lys | Val | Arg | Met | Tyr | Val | Gly | Gly |  |     |
|     |     |     | 260 |     |     |     |     | 265 |     |     |     |     | 270 |     |     |  |     |
| GTG | GAG | CAC | AGG | TTC | GAA | GCC | GCA | TGC | AAT | TGG | ACT | CGA | GGA | GAG | CGT |  | 864 |
| Val | Glu | His | Arg | Phe | Glu | Ala | Ala | Cys | Asn | Trp | Thr | Arg | Gly | Glu | Arg |  |     |
|     |     |     | 275 |     |     |     | 280 |     |     |     |     |     | 285 |     |     |  |     |
| TGT | GAC | TTG | GAG | GAC | AGG | GAT | AGA | TCA | GAG | CTT | AGC | CCG | CTG | CTG | CTG |  | 912 |
| Cys | Asp | Leu | Glu | Asp | Arg | Asp | Arg | Ser | Glu | Leu | Ser | Pro | Leu | Leu | Leu |  |     |
|     | 290 |     |     |     |     | 295 |     |     |     |     | 300 |     |     |     |     |  |     |
| TCT | ACA | ACA | GGT | GAT | CGA | GGG | CAG | ACA | CCA | TCA | CCA | CCA | TCA | CTA | A   |  | 958 |
| Ser | Thr | Thr | Gly | Asp | Arg | Gly | Gln | Thr | Pro | Ser | Pro | Pro | Ser | Leu |     |  |     |
| 305 |     |     |     |     | 310 |     |     |     |     | 315 |     |     |     |     |     |  |     |
| TAG |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |  | 961 |

## (2) INFORMATION FOR SEQ ID NO: 44:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 319 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 44:

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |  |  |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|--|
| Met | Val | Gly | Asn | Trp | Ala | Lys | Val | Leu | Val | Val | Met | Leu | Leu | Phe | Ala |  |  |
| 1   |     |     |     | 5   |     |     |     | 10  |     |     |     |     |     | 15  |     |  |  |
| Gly | Val | Asp | Gly | His | Thr | Arg | Val | Ser | Gly | Gly | Ala | Ala | Ala | Ser | Asp |  |  |
|     |     | 20  |     |     |     |     |     | 25  |     |     |     |     | 30  |     |     |  |  |
| Thr | Arg | Gly | Leu | Val | Ser | Leu | Phe | Ser | Pro | Gly | Ser | Ala | Gln | Lys | Ile |  |  |
|     |     | 35  |     |     |     | 40  |     |     |     |     |     | 45  |     |     |     |  |  |
| Gln | Leu | Val | Asn | Thr | Asn | Gly | Ser | Trp | His | Ile | Asn | Arg | Thr | Ala | Leu |  |  |
|     | 50  |     |     |     |     | 55  |     |     |     |     | 60  |     |     |     |     |  |  |
| Asn | Cys | Asn | Asp | Ser | Leu | Gln | Thr | Gly | Phe | Phe | Ala | Ala | Leu | Phe | Tyr |  |  |
| 65  |     |     |     | 70  |     |     |     |     | 75  |     |     |     |     | 80  |     |  |  |
| Lys | His | Lys | Phe | Asn | Ser | Ser | Gly | Cys | Pro | Glu | Arg | Leu | Ala | Ser | Cys |  |  |
|     |     |     | 85  |     |     |     |     | 90  |     |     |     |     |     | 95  |     |  |  |
| Arg | Ser | Ile | Asp | Lys | Phe | Ala | Gln | Gly | Trp | Gly | Pro | Leu | Thr | Tyr | Thr |  |  |
|     |     | 100 |     |     |     |     | 105 |     |     |     |     |     | 110 |     |     |  |  |
| Glu | Pro | Asn | Ser | Ser | Asp | Gln | Arg | Pro | Tyr | Cys | Trp | His | Tyr | Ala | Pro |  |  |
|     | 115 |     |     |     |     | 120 |     |     |     |     |     | 125 |     |     |     |  |  |
| Arg | Pro | Cys | Gly | Ile | Val | Pro | Ala | Ser | Gln | Val | Cys | Gly | Pro | Val | Tyr |  |  |

|                                                                 |     |         |
|-----------------------------------------------------------------|-----|---------|
| 130                                                             | 135 | 140     |
| Cys Phe Thr Pro Ser Pro Val Val Val Gly Thr Thr Asp Arg Phe Gly |     |         |
| 145                                                             | 150 | 155 160 |
| Val Pro Thr Tyr Asn Trp Gly Ala Asn Asp Ser Asp Val Leu Ile Leu |     |         |
|                                                                 | 165 | 170 175 |
| Asn Asn Thr Arg Pro Pro Arg Gly Asn Trp Phe Gly Cys Thr Trp Met |     |         |
|                                                                 | 180 | 185 190 |
| Asn Gly Thr Gly Phe Thr Lys Thr Cys Gly Gly Pro Pro Cys Asn Ile |     |         |
|                                                                 | 195 | 200 205 |
| Gly Gly Ala Gly Asn Asn Thr Leu Thr Cys Pro Thr Asp Cys Phe Arg |     |         |
|                                                                 | 210 | 215 220 |
| Lys His Pro Glu Ala Thr Tyr Ala Arg Cys Gly Ser Gly Pro Trp Leu |     |         |
| 225                                                             | 230 | 235 240 |
| Thr Pro Arg Cys Met Val His Tyr Pro Tyr Arg Leu Trp His Tyr Pro |     |         |
|                                                                 | 245 | 250 255 |
| Cys Thr Val Asn Phe Thr Ile Phe Lys Val Arg Met Tyr Val Gly Gly |     |         |
|                                                                 | 260 | 265 270 |
| Val Glu His Arg Phe Glu Ala Ala Cys Asn Trp Thr Arg Gly Glu Arg |     |         |
|                                                                 | 275 | 280 285 |
| Cys Asp Leu Glu Asp Arg Asp Arg Ser Glu Leu Ser Pro Leu Leu Leu |     |         |
|                                                                 | 290 | 295 300 |
| Ser Thr Thr Gly Asp Arg Gly Glu Thr Pro Ser Pro Pro Ser Leu     |     |         |
| 305                                                             | 310 | 315     |

## (2) INFORMATION FOR SEQ ID NO: 45:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1395 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..1392

## (ix) FEATURE:

- (A) NAME/KEY: mat\_peptide
- (B) LOCATION: 1..1389

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 45:

ATG GTG GCG GGG GCC CAT TGG GGA GTC CTG GCG GGC CTC GCC TAC TAT

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Met | Val | Ala | Gly | Ala | His | Trp | Gly | Val | Leu | Ala | Gly | Leu | Ala | Tyr | Tyr |     |
| 1   |     |     |     | 5   |     |     |     |     | 10  |     |     |     |     | 15  |     |     |
| TCC | ATG | GTG | GGG | AAC | TGG | GCT | AAG | GTT | TTG | GTT | GTG | ATG | CTA | CTC | TTT | 96  |
| Ser | Met | Val | Gly | Asn | Trp | Ala | Lys | Val | Leu | Val | Val | Met | Leu | Leu | Phe |     |
|     |     |     | 20  |     |     |     |     | 25  |     |     |     |     | 30  |     |     |     |
| GCC | GGC | GTC | GAC | GGG | CAT | ACC | CGC | GTG | TCA | GGA | GGG | GCA | GCA | GCC | TCC | 144 |
| Ala | Gly | Val | Asp | Gly | His | Thr | Arg | Val | Ser | Gly | Gly | Ala | Ala | Ala | Ser |     |
|     |     | 35  |     |     |     |     | 40  |     |     |     |     | 45  |     |     |     |     |
| GAT | ACC | AGG | GGC | CTT | GTG | TCC | CTC | TTT | AGC | CCC | GGG | TCG | GCT | CAG | AAA | 192 |
| Asp | Thr | Arg | Gly | Leu | Val | Ser | Leu | Phe | Ser | Pro | Gly | Ser | Ala | Gln | Lys |     |
|     | 50  |     |     |     |     | 55  |     |     |     |     | 60  |     |     |     |     |     |
| ATC | CAG | CTC | GTA | AAC | ACC | AAC | GGC | AGT | TGG | CAC | ATC | AAC | AGG | ACT | GCC | 240 |
| Ile | Gln | Leu | Val | Asn | Thr | Asn | Gly | Ser | Trp | His | Ile | Asn | Arg | Thr | Ala |     |
| 65  |     |     |     | 70  |     |     |     |     | 75  |     |     |     |     | 80  |     |     |
| CTG | AAC | TGC | AAC | GAC | TCC | CTC | CAA | ACA | GGG | TTC | TTT | GCC | GCA | CTA | TTC | 288 |
| Leu | Asn | Cys | Asn | Asp | Ser | Leu | Gln | Thr | Gly | Phe | Phe | Ala | Ala | Leu | Phe |     |
|     |     |     | 85  |     |     |     |     |     | 90  |     |     |     |     | 95  |     |     |
| TAC | AAA | CAC | AAA | TTC | AAC | TCG | TCT | GGA | TGC | CCA | GAG | CGC | TTG | GCC | AGC | 336 |
| Tyr | Lys | His |     | Phe | Asn | Ser | Ser | Gly | Cys | Pro | Glu | Arg | Leu | Ala | Ser |     |
|     |     | 100 |     |     |     |     |     | 105 |     |     |     |     | 110 |     |     |     |
| TGT | CGC | TCC | ATC | GAC | AAG | TTC | GCT | CAG | GGG | TGG | GGT | CCC | CTC | ACT | TAC | 384 |
| Cys | Arg | Ser | Ile | Asp | Lys | Phe | Ala | Gln | Gly | Trp | Gly | Pro | Leu | Thr | Tyr |     |
|     |     | 115 |     |     |     | 120 |     |     |     |     |     | 125 |     |     |     |     |
| ACT | GAG | CCT | AAC | AGC | TCG | GAC | CAG | AGG | CCC | TAC | TGC | TGG | CAC | TAC | GCG | 432 |
| Thr | Glu | Pro | Asn | Ser | Ser |     | Gln | Arg | Pro | Tyr | Cys | Trp | His | Tyr | Ala |     |
|     | 130 |     |     |     |     | 135 |     |     |     |     |     | 140 |     |     |     |     |
| CCT | CGA | CCG | TGT | GGT | ATT | GTA | CCC | GCG | TCT | CAG | GTG | TGC | GGT | CCA | GTG | 480 |
| Pro | Arg | Pro | Cys | Gly | Ile | Val | Pro | Ala | Ser | Gln | Val | Cys | Gly | Pro | Val |     |
| 145 |     |     |     | 150 |     |     |     |     |     | 155 |     |     |     |     | 160 |     |
| TAT | TGC | TTC | ACC | CCG | AGC | CCT | GTT | GTG | GTG | GGG | ACG | ACC | GAT | CGG | TTT | 528 |
| Tyr | Cys | Phe | Thr | Pro | Ser | Pro | Val | Val | Gly | Thr | Thr | Asp | Arg | Phe |     |     |
|     |     |     | 165 |     |     |     |     | 170 |     |     |     |     |     | 175 |     |     |
| GGT | GTC | CCC | ACG | TAT | AAC | TGG | GGG | GCG | AAC | GAC | TCG | GAT | GTG | CTG | ATT | 576 |
| Gly | Val | Pro | Thr | Tyr | Asn | Trp | Gly | Ala | Asn | Asp | Ser | Asp | Val | Leu | Ile |     |
|     |     | 180 |     |     |     |     |     | 185 |     |     |     |     | 190 |     |     |     |
| CTC | AAC | AAC | ACG | CGG | CCG | CCG | CGA | GGC | AAC | TGG | TTC | GGC | TGT | ACA | TGG | 624 |
| Leu | Asn | Asn | Thr | Arg | Pro | Pro | Arg | Gly | Asn | Trp | Phe | Gly | Cys | Thr | Trp |     |
|     |     | 195 |     |     |     | 200 |     |     |     |     |     | 205 |     |     |     |     |
| ATG | AAT | GGC | ACT | GGG | TTC | ACC | AAG | ACG | TGT | GGG | GGC | CCC | CCG | TGC | AAC | 672 |
| Met | Asn | Gly | Thr | Gly | Phe | Thr | Lys | Thr | Cys | Gly | Gly | Pro | Pro | Cys | Asn |     |
|     | 210 |     |     |     |     | 215 |     |     |     |     | 220 |     |     |     |     |     |
| ATC | GGG | GGG | GCC | GGC | AAC | AAC | ACC | TTG | ACC | TGC | CCC | ACT | GAC | TGT | TTT | 720 |
| Ile | Gly | Gly | Ala | Gly | Asn | Asn | Thr | Leu | Thr | Cys | Pro | Thr | Asp | Cys | Phe |     |
| 225 |     |     |     | 230 |     |     |     |     |     | 235 |     |     |     | 240 |     |     |
| CGG | AAG | CAC | CCC | GAG | GCC | ACC | TAC | GCC | AGA | TGC | GGT | TCT | GGG | CCC | TGG | 768 |
| Arg | Lys | His | Pro | Glu | Ala | Thr | Tyr | Ala | Arg | Cys | Gly | Ser | Gly | Pro | Trp |     |
|     |     |     | 245 |     |     |     |     | 250 |     |     |     |     |     | 255 |     |     |



|                                                                    |      |
|--------------------------------------------------------------------|------|
| CTG ACA CCT AGG TGT ATG GTT CAT TAC CCA TAT AGG CTC TGG CAC TAC    | 816  |
| Leu Thr Pro Arg Cys Met Val His Tyr Pro Tyr Arg Leu Trp His Tyr    |      |
| 260 265 270                                                        |      |
| CCC TGC ACT GTC AAC TTC ACC ATC TTC AAG GTT AGG ATG TAC GTG GGG    | 864  |
| Pro Cys Thr Val Asn Phe Thr Ile Phe Lys Val Arg Met Tyr Val Gly    |      |
| 275 280 285                                                        |      |
| GGC GTG GAG CAC AGG TTC GAA GCC GCA TGC AAT TGG ACT CGA GGA GAG    | 912  |
| Gly Val Glu His Arg Phe Glu Ala Ala Cys Asn Trp Thr Arg Gly Glu    |      |
| 290 295 300                                                        |      |
| CGT TGT GAC TTG GAG GAC AGG GAT AGA TCA GAG CTT AGC CCG CTG CTG    | 960  |
| Arg Cys Asp Leu Glu Asp Arg Asp Arg Ser Glu Leu Ser Pro Leu Leu    |      |
| 305 310 315 320                                                    |      |
| CTG TCT ACA ACA GAG TGG CAG ATA CTG CCC TGT TCC TTC ACC ACC CTG    | 1008 |
| Leu Ser Thr Thr Glu Trp Gln Ile Leu Pro Cys Ser Phe Thr Thr Leu    |      |
| 325 330 335                                                        |      |
| CCG GCC CTA TCC ACC GGC CTG ATC CAC CTC CAT CAG AAC ATC GTG GAC    | 1056 |
| Pro Ala Leu Ser Thr Gly Leu Ile His Leu His Gln Asn Ile Val Asp    |      |
| 340 345 350                                                        |      |
| GTG CAA TAC CTG TAC GGT GTA GGG TCG GCG GTT GTC TCC CTT GTC ATC    | 1104 |
| Val Gln Tyr Leu Tyr Gly Val Gly Ser Ala Val Val Ser Leu Val Ile    |      |
| 355 360 365                                                        |      |
| AAA TGG GAG TAT GTC CTG TTG CTC TTC CTT CTC CTG GCA GAC GCG CGC    | 1152 |
| Lys Trp Glu Tyr Val Leu Leu Leu Phe Leu Leu Leu Ala Asp Ala Arg    |      |
| 370 375 380                                                        |      |
| ATC TGC GCC TGC TTA TGG ATG ATG CTG CTG ATA GCT CAA GCT GAG GCC    | 1200 |
| Ile Cys Ala Cys Leu Trp Met Met Leu Leu Ile Ala Gln Ala Glu Ala    |      |
| 385 390 395 400                                                    |      |
| GCC TTA GAG AAC CTG GTG GTC CTC AAT GCG GCG GCC GTG GCC GGG GCG    | 1248 |
| Ala Leu Glu Asn Leu Val Val Leu Asn Ala Ala Ala Val Ala Gly Ala    |      |
| 405 410 415                                                        |      |
| CAT GGC ACT CTT TCC TTC CTT GTG TTC TTC TGT GCT GCC TGG TAC ATC    | 1296 |
| His Gly Thr Leu Ser Phe Leu Val Phe Phe Cys Ala Ala Trp Tyr Ile    |      |
| 420 425 430                                                        |      |
| AAG GGC AGG CTG GTC CCT GGT GCG GCA TAC GCC TTC TAT GGC GTG TGG    | 1344 |
| Lys Gly Arg Leu Val Pro Gly Ala Ala Tyr Ala Phe Tyr Gly Val Trp    |      |
| 435 440 445                                                        |      |
| CCG CTG CTC CTG CTT CTG CTG GCC TTA CCA CCA CGA GCT TAT GCC TAGTAA | 1395 |
| Pro Leu Leu Leu Leu Leu Ala Leu Pro Pro Arg Ala Tyr Ala            |      |
| 450 455 460                                                        |      |

## (2) INFORMATION FOR SEQ ID NO: 46:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 463 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 46:

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Met | Val | Ala | Gly | Ala | His | Trp | Gly | Val | Leu | Ala | Gly | Leu | Ala | Tyr | Tyr | 1   | 5   | 10  | 15  |
| Ser | Met | Val | Gly | Asn | Trp | Ala | Lys | Val | Leu | Val | Val | Met | Leu | Leu | Phe | 20  | 25  | 30  |     |
| Ala | Gly | Val | Asp | Gly | His | Thr | Arg | Val | Ser | Gly | Gly | Ala | Ala | Ala | Ser | 35  | 40  | 45  |     |
| Asp | Thr | Arg | Gly | Leu | Val | Ser | Leu | Phe | Ser | Pro | Gly | Ser | Ala | Gln | Lys | 50  | 55  | 60  |     |
| Ile | Gln | Leu | Val | Asn | Thr | Asn | Gly | Ser | Trp | His | Ile | Asn | Arg | Thr | Ala | 65  | 70  | 75  | 80  |
| Leu | Asn | Cys | Asn | Asp | Ser | Leu | Gln | Thr | Gly | Phe | Phe | Ala | Ala | Leu | Phe | 85  | 90  | 95  |     |
| Tyr | Lys | His | Lys | Phe | Asn | Ser | Ser | Gly | Cys | Pro | Glu | Arg | Leu | Ala | Ser | 100 | 105 | 110 |     |
| Cys | Arg | Ser | Ile | Asp | Lys | Phe | Ala | Gln | Gly | Trp | Gly | Pro | Leu | Thr | Tyr | 115 | 120 | 125 |     |
| Thr | Glu | Pro | Asn | Ser | Ser | Asp | Gln | Arg | Pro | Tyr | Cys | Trp | His | Tyr | Ala | 130 | 135 | 140 |     |
| Pro | Arg | Pro | Cys | Gly | Ile | Val | Pro | Ala | Ser | Gln | Val | Cys | Gly | Pro | Val | 145 | 150 | 155 | 160 |
| Tyr | Cys | Phe | Thr | Pro | Ser | Pro | Val | Val | Val | Gly | Thr | Thr | Asp | Arg | Phe | 165 | 170 | 175 |     |
| Gly | Val | Pro | Thr | Tyr | Asn | Trp | Gly | Ala | Asn | Asp | Ser | Asp | Val | Leu | Ile | 180 | 185 | 190 |     |
| Leu | Asn | Asn | Thr | Arg | Pro | Pro | Arg | Gly | Asn | Trp | Phe | Gly | Cys | Thr | Trp | 195 | 200 | 205 |     |
| Met | Asn | Gly | Thr | Gly | Phe | Thr | Lys | Thr | Cys | Gly | Gly | Pro | Pro | Cys | Asn | 210 | 215 | 220 |     |
| Ile | Gly | Gly | Ala | Gly | Asn | Asn | Thr | Leu | Thr | Cys | Pro | Thr | Asp | Cys | Phe | 225 | 230 | 235 | 240 |
| Arg | Lys | His | Pro | Glu | Ala | Thr | Tyr | Ala | Arg | Cys | Gly | Ser | Gly | Pro | Trp | 245 | 250 | 255 |     |
| Leu | Thr | Pro | Arg | Cys | Met | Val | His | Tyr | Pro | Tyr | Arg | Leu | Trp | His | Tyr | 260 | 265 | 270 |     |
| Pro | Cys | Thr | Val | Asn | Phe | Thr | Ile | Phe | Lys | Val | Arg | Met | Tyr | Val | Gly | 275 | 280 | 285 |     |
| Gly | Val | Glu | His | Arg | Phe | Glu | Ala | Ala | Cys | Asn | Trp | Thr | Arg | Gly | Glu | 290 | 295 | 300 |     |
| Arg | Cys | Asp | Leu | Glu | Asp | Arg | Asp | Arg | Ser | Glu | Leu | Ser | Pro | Leu | Leu | 305 | 310 | 315 | 320 |

50

```

Leu Ser Thr Thr Glu Trp Gln Ile Leu Pro Cys Ser Phe Thr Thr Leu
      325                      330                      335
Pro Ala Leu Ser Thr Gly Leu Ile His Leu His Gln Asn Ile Val Asp
      340                      345                      350
Val Gln Tyr Leu Tyr Gly Val Gly Ser Ala Val Val Ser Leu Val Ile
      355                      360                      365
Lys Trp Glu Tyr Val Leu Leu Leu Phe Leu Leu Leu Ala Asp Ala Arg
      370                      375                      380
Ile Cys Ala Cys Leu Trp Met Met Leu Leu Ile Ala Gln Ala Glu Ala
      385                      390                      395                      400
Ala Leu Glu Asn Leu Val Val Leu Asn Ala Ala Ala Val Ala Gly Ala
      405                      410                      415
His Gly Thr Leu Ser Phe Leu Val Phe Phe Cys Ala Ala Trp Tyr Ile
      420                      425                      430
Lys Gly Arg Leu Val Pro Gly Ala Ala Tyr Ala Phe Tyr Gly Val Trp
      435                      440                      445
Pro Leu Leu Leu Leu Leu Leu Ala Leu Pro Pro Arg Ala Tyr Ala
      450                      455                      460

```

## (2) INFORMATION FOR SEQ ID NO: 47:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 2082 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..2079

## (ix) FEATURE:

- (A) NAME/KEY: mat\_peptide
- (B) LOCATION: 1..2076

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 47:

```

AAT TTG GGT AAG GTC ATC GAT ACC CTT ACA TGC GGC TTC GCC GAC CTC      48
Asn Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu
  1                      5                      10                      15

GTG GGG TAC ATT CCG CTC GTC GGC GCC CCC CTA GGG GGC GCT GCC AGG      96
Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg
      20                      25                      30

```

|                                                                                                                                                       |     |
|-------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| GCC CTG GCG CAT GGC GTC CGG GTT CTG GAG GAC GGC GTG AAC TAT GCA<br>Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala<br>35 40 45        | 144 |
| ACA GGG AAT TTG CCC GGT TGC TCT TTC TCT ATC TTC CTC TTG GCT TTG<br>Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu<br>50 55 60        | 192 |
| CTG TCC TGT CTG ACC GTT CCA GCT TCC GCT TAT GAA GTG CGC AAC GTG<br>Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr Glu Val Arg Asn Val<br>65 70 75 80     | 240 |
| TCC GGG ATG TAC CAT GTC ACG AAC GAC TGC TCC AAC TCA AGC ATT GTG<br>Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser Asn Ser Ser Ile Val<br>85 90 95        | 288 |
| TAT GAG GCA GCG GAC ATG ATC ATG CAC ACC CCC GGG TGC GTG CCC TGC<br>Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys Val Pro Cys<br>100 105 110     | 336 |
| GTT CGG GAG AAC AAC TCT TCC CGC TGC TGG GTA GCG CTC ACC CCC ACG<br>Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val Ala Leu Thr Pro Thr<br>115 120 125     | 384 |
| CTC GCA GCT AGG AAC GCC AGC GTC CCC ACC ACG ACA ATA CGA CGC CAC<br>Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr Thr Ile Arg Arg His<br>130 135 140     | 432 |
| GTC GAT TTG CTC GTT GGG GCG GCT GCT TTC TGT TCC GCT ATG TAC GTG<br>Val Asp Leu Leu Val Gly Ala Ala Phe Cys Ser Ala Met Tyr Val<br>145 150 155 160     | 480 |
| GGG GAC CTC TGC GGA TCT GTC TTC CTC GTC TCC CAG CTG TTC ACC ATC<br>Gly Asp Leu Cys Gly Ser Val Phe Leu Val Ser Gln Leu Phe Thr Ile<br>165 170 175     | 528 |
| TCG CCT CGC CGG CAT GAG ACG GTG CAG GAC TGC AAT TGC TCA ATC TAT<br>Ser Pro Arg Arg His Glu Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr<br>180 185 190     | 576 |
| CCC GGC CAC ATA ACG GGT CAC CGT ATG GCT TGG GAT ATG ATG ATG AAC<br>Pro Gly His Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn<br>195 200 205     | 624 |
| TGG TCG CCT ACA ACG GCC CTG GTG GTA TCG CAG CTG CTC CGG ATC CCA<br>Trp Ser Pro Thr Thr Ala Leu Val Ser Gln Leu Leu Arg Ile Pro<br>210 215 220         | 672 |
| CAA GCT GTC GTG GAC ATG GTG GCG GGG GCC CAT TGG GGA GTC CTG GCG<br>Gln Ala Val Val Asp Met Val Ala Gly Ala His Trp Gly Val Leu Ala<br>225 230 235 240 | 720 |
| GGC CTC GCC TAC TAT TCC ATG GTG GGG AAC TGG GCT AAG GTT TTG GTT<br>Gly Leu Ala Tyr Tyr Ser Met Val Gly Asn Trp Ala Lys Val Leu Val<br>245 250 255     | 768 |
| GTG ATG CTA CTC TTT GCC GGC GTC GAC GGG CAT ACC CGC GTG TCA GGA<br>Val Met Leu Leu Phe Ala Gly Val Asp Gly His Thr Arg Val Ser Gly<br>260 265 270     | 816 |
| GGG GCA GCA GCC TCC GAT ACC AGG GGC CTT GTG TCC CTC TTT AGC CCC<br>Gly Ala Ala Ser Asp Thr Arg Gly Leu Val Ser Leu Phe Ser Pro                        | 864 |

| 275                                                                                                                                                   | 280 | 285 |      |
|-------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----|------|
| GGG TCG GCT CAG AAA ATC CAG CTC GTA AAC ACC AAC GGC AGT TGG CAC<br>Gly Ser Ala Gln Lys Ile Gln Leu Val Asn Thr Asn Gly Ser Trp His<br>290 295 300     |     |     | 912  |
| ATC AAC AGG ACT GCC CTG AAC TGC AAC GAC TCC CTC CAA ACA GGG TTC<br>Ile Asn Arg Thr Ala Leu Asn Cys Asn Asp Ser Leu Gln Thr Gly Phe<br>305 310 315 320 |     |     | 960  |
| TTT GCC GCA CTA TTC TAC AAA CAC AAA TTC AAC TCG TCT GGA TGC CCA<br>Phe Ala Ala Leu Phe Tyr Lys His Lys Phe Asn Ser Ser Gly Cys Pro<br>325 330 335     |     |     | 1008 |
| GAG CGC TTG GCC AGC TGT CGC TCC ATC GAC AAG TTC GCT CAG GGG TGG<br>Glu Arg Leu Ala Ser Cys Arg Ser Ile Asp Lys Phe Ala Gln Gly Trp<br>340 345 350     |     |     | 1056 |
| GGT CCC CTC ACT TAC ACT GAG CCT AAC AGC TCG GAC CAG AGG CCC TAC<br>Gly Pro Leu Thr Tyr Thr Glu Pro Asn Ser Ser Asp Gln Arg Pro Tyr<br>355 360 365     |     |     | 1104 |
| TGC TGG CAC TAC GCG CCT CGA CCG TGT GGT ATT GTA CCC GCG TCT CAG<br>Cys Trp His Tyr Ala Pro Arg Pro Cys Gly Ile Val Pro Ala Ser Gln<br>370 375 380     |     |     | 1152 |
| GTG TGC GGT CCA GTG TAT TGC TTC ACC CCG AGC CCT GTT GTG GTG GGG<br>Val Cys Gly Pro Val Tyr Cys Phe Thr Pro Ser Pro Val Val Val Gly<br>385 390 395 400 |     |     | 1200 |
| ACG ACC GAT CGG TTT GGT GTC CCC ACG TAT AAC TGG GGG GCG AAC GAC<br>Thr Thr Asp Arg Phe Gly Val Pro Thr Tyr Asn Trp Gly Ala Asn Asp<br>405 410 415     |     |     | 1248 |
| TCG GAT GTG CTG ATT CTC AAC AAC ACG CGG CCG CCG CGA GGC AAC TGG<br>Ser Asp Val Leu Ile Leu Asn Asn Thr Arg Pro Pro Arg Gly Asn Trp<br>420 425 430     |     |     | 1296 |
| TTC GGC TGT ACA TGG ATG AAT GGC ACT GGG TTC ACC AAG ACG TGT GGG<br>Phe Gly Cys Thr Trp Met Asn Gly Thr Gly Phe Thr Lys Thr Cys Gly<br>435 440 445     |     |     | 1344 |
| GGC CCC CCG TGC AAC ATC GGG GGG GCC GGC AAC AAC ACC TTG ACC TGC<br>Gly Pro Pro Cys Asn Ile Gly Gly Ala Gly Asn Asn Thr Leu Thr Cys<br>450 455 460     |     |     | 1392 |
| CCC ACT GAC TGT TTT CGG AAG CAC CCC GAG GCC ACC TAC GCC AGA TGC<br>Pro Thr Asp Cys Phe Arg Lys His Pro Glu Ala Thr Tyr Ala Arg Cys<br>465 470 475 480 |     |     | 1440 |
| GGT TCT GGG CCC TGG CTG ACA CCT AGG TGT ATG GTT CAT TAC CCA TAT<br>Gly Ser Gly Pro Trp Leu Thr Pro Arg Cys Met Val His Tyr Pro Tyr<br>485 490 495     |     |     | 1488 |
| AGG CTC TGG CAC TAC CCC TGC ACT GTC AAC TTC ACC ATC TTC AAG GTT<br>Arg Leu Trp His Tyr Pro Cys Thr Val Asn Phe Thr Ile Phe Lys Val<br>500 505 510     |     |     | 1536 |
| AGG ATG TAC GTG GGG GGC GTG GAG CAC AGG TTC GAA GCC GCA TGC AAT<br>Arg Met Tyr Val Gly Gly Val Glu His Arg Phe Glu Ala Ala Cys Asn<br>515 520 525     |     |     | 1584 |
| TGG ACT CGA GGA GAG CGT TGT GAC TTG GAG GAC AGG GAT AGA TCA GAG                                                                                       |     |     | 1632 |

|     |     |     |     |        |     |     |     |     |     |     |     |     |     |     |     |      |
|-----|-----|-----|-----|--------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|
| Trp | Thr | Arg | Gly | Glu    | Arg | Cys | Asp | Leu | Glu | Asp | Arg | Asp | Arg | Ser | Glu |      |
| 530 |     |     |     |        |     | 535 |     |     |     |     | 540 |     |     |     |     |      |
| CTT | AGC | CCG | CTG | CTG    | CTG | TCT | ACA | ACA | GAG | TGG | CAG | ATA | CTG | CCC | TGT | 1680 |
| Leu | Ser | Pro | Leu | Leu    | Leu | Ser | Thr | Thr | Glu | Trp | Gln | Ile | Leu | Pro | Cys |      |
| 545 |     |     |     |        | 550 |     |     |     | 555 |     |     |     |     |     | 560 |      |
| TCC | TTC | ACC | ACC | CTG    | CCG | GCC | CTA | TCC | ACC | GGC | CTG | ATC | CAC | CTC | CAT | 1728 |
| Ser | Phe | Thr | Thr | Leu    | Pro | Ala | Leu | Ser | Thr | Gly | Leu | Ile | His | Leu | His |      |
|     |     |     |     | 565    |     |     |     |     | 570 |     |     |     |     | 575 |     |      |
| CAG | AAC | ATC | GTG | GAC    | GTG | CAA | TAC | CTG | TAC | GGT | GTA | GGG | TCG | GCG | GTT | 1776 |
| Gln | Asn | Ile | Val | Asp    | Val | Gln | Tyr | Leu | Tyr | Gly | Val | Gly | Ser | Ala | Val |      |
|     |     |     | 580 |        |     |     |     | 585 |     |     |     |     | 590 |     |     |      |
| GTC | TCC | CTT | GTC | ATC    | AAA | TGG | GAG | TAT | GTC | CTG | TTG | CTC | TTC | CTT | CTC | 1824 |
| Val | Ser | Leu | Val | Ile    | Lys | Trp | Glu | Tyr | Val | Leu | Leu | Leu | Phe | Leu | Leu |      |
|     |     | 595 |     |        |     | 600 |     |     |     |     |     | 605 |     |     |     |      |
| CTG | GCA | GAC | GCG | CGC    | ATC | TGC | GCC | TGC | TTA | TGG | ATG | ATG | CTG | CTG | ATA | 1872 |
| Leu | Ala | Asp | Ala | Arg    | Ile | Cys | Ala | Cys | Leu | Trp | Met | Met | Leu | Leu | Ile |      |
|     | 610 |     |     |        |     | 615 |     |     |     |     | 620 |     |     |     |     |      |
| GCT | CAA | GCT | GAG | GCC    | GCC | TTA | GAG | AAC | CTG | GTG | GTC | CTC | AAT | GCG | GCG | 1920 |
| Ala | Gln | Ala | Glu | Ala    | Ala | Leu | Glu | Asn | Leu | Val | Val | Leu | Asn | Ala | Ala |      |
| 625 |     |     |     |        | 630 |     |     |     | 635 |     |     |     |     |     | 640 |      |
| GCC | GTG | GCC | GGG | GCG    | CAT | GGC | ACT | CTT | TCC | TTC | CTT | GTG | TTC | TTC | TGT | 1968 |
| Ala | Val | Ala | Gly | Ala    | His | Gly | Thr | Leu | Ser | Phe | Leu | Val | Phe | Phe | Cys |      |
|     |     |     |     | 645    |     |     |     |     | 650 |     |     |     |     | 655 |     |      |
| GCT | GCC | TGG | TAC | ATC    | AAG | GGC | AGG | CTG | GTC | CCT | GGT | GCG | GCA | TAC | GCC | 2016 |
| Ala | Ala | Trp | Tyr | Ile    | Lys | Gly | Arg | Leu | Val | Pro | Gly | Ala | Ala | Tyr | Ala |      |
|     |     |     | 660 |        |     |     |     | 665 |     |     |     |     | 670 |     |     |      |
| TTC | TAT | GGC | GTG | TGG    | CCG | CTG | CTC | CTG | CTT | CTG | CTG | GCC | TTA | CCA | CCA | 2064 |
| Phe | Tyr | Gly | Val | Trp    | Pro | Leu | Leu | Leu | Leu | Leu | Leu | Ala | Leu | Pro | Pro |      |
|     |     | 675 |     |        |     | 680 |     |     |     |     |     | 685 |     |     |     |      |
| CGA | GCT | TAT | GCC | TAGTAA |     |     |     |     |     |     |     |     |     |     |     | 2082 |
| Arg | Ala | Tyr | Ala |        |     |     |     |     |     |     |     |     |     |     |     |      |
|     |     | 690 |     |        |     |     |     |     |     |     |     |     |     |     |     |      |

## (2) INFORMATION FOR SEQ ID NO: 48:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 692 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 48:

Asn Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu  
 1 5 10 15  
 Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg  
 20 25 30  
 Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala

| 35  |     |     |     |     | 40  |     |     |     |     | 45  |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Thr | Gly | Asn | Leu | Pro | Gly | Cys | Ser | Phe | Ser | Ile | Phe | Leu | Leu | Ala | Leu |
| 50  |     |     |     |     |     | 55  |     |     |     |     | 60  |     |     |     |     |
| Leu | Ser | Cys | Leu | Thr | Val | Pro | Ala | Ser | Ala | Tyr | Glu | Val | Arg | Asn | Val |
| 65  |     |     |     |     | 70  |     |     |     |     | 75  |     |     |     |     | 80  |
| Ser | Gly | Met | Tyr | His | Val | Thr | Asn | Asp | Cys | Ser | Asn | Ser | Ser | Ile | Val |
|     |     |     |     | 85  |     |     |     |     | 90  |     |     |     |     | 95  |     |
| Tyr | Glu | Ala | Ala | Asp | Met | Ile | Met | His | Thr | Pro | Gly | Cys | Val | Pro | Cys |
|     |     |     | 100 |     |     |     |     | 105 |     |     |     |     | 110 |     |     |
| Val | Arg | Glu | Asn | Asn | Ser | Ser | Arg | Cys | Trp | Val | Ala | Leu | Thr | Pro | Thr |
|     |     | 115 |     |     |     |     | 120 |     |     |     |     | 125 |     |     |     |
| Leu | Ala | Ala | Arg | Asn | Ala | Ser | Val | Pro | Thr | Thr | Thr | Ile | Arg | Arg | His |
|     |     | 130 |     |     |     |     | 135 |     |     |     |     | 140 |     |     |     |
| Val | Asp | Leu | Leu | Val | Gly | Ala | Ala | Ala | Phe | Cys | Ser | Ala | Met | Tyr | Val |
| 145 |     |     |     |     | 150 |     |     |     |     | 155 |     |     |     |     | 160 |
| Gly | Asp | Leu | Cys | Gly | Ser | Val | Phe | Leu | Val | Ser | Gln | Leu | Phe | Thr | Ile |
|     |     |     | 165 |     |     |     |     |     | 170 |     |     |     |     | 175 |     |
| Ser | Pro | Arg | Arg | His | Glu | Thr | Val | Gln | Asp | Cys | Asn | Cys | Ser | Ile | Tyr |
|     |     |     | 180 |     |     |     |     | 185 |     |     |     |     | 190 |     |     |
| Pro | Gly | His | Ile | Thr | Gly | His | Arg | Met | Ala | Trp | Asp | Met | Met | Met | Asn |
|     |     | 195 |     |     |     |     | 200 |     |     |     |     | 205 |     |     |     |
| Trp | Ser | Pro | Thr | Thr | Ala | Leu | Val | Val | Ser | Gln | Leu | Leu | Arg | Ile | Pro |
|     |     | 210 |     |     |     | 215 |     |     |     |     | 220 |     |     |     |     |
| Gln | Ala | Val | Val | Asp | Met | Val | Ala | Gly | Ala | His | Trp | Gly | Val | Leu | Ala |
| 225 |     |     |     |     | 230 |     |     |     |     | 235 |     |     |     |     | 240 |
| Gly | Leu | Ala | Tyr | Tyr | Ser | Met | Val | Gly | Asn | Trp | Ala | Lys | Val | Leu | Val |
|     |     |     |     | 245 |     |     |     |     | 250 |     |     |     |     | 255 |     |
| Val | Met | Leu | Leu | Phe | Ala | Gly | Val | Asp | Gly | His | Thr | Arg | Val | Ser | Gly |
|     |     |     | 260 |     |     |     |     | 265 |     |     |     |     | 270 |     |     |
| Gly | Ala | Ala | Ala | Ser | Asp | Thr | Arg | Gly | Leu | Val | Ser | Leu | Phe | Ser | Pro |
|     |     | 275 |     |     |     |     | 280 |     |     |     |     | 285 |     |     |     |
| Gly | Ser | Ala | Gln | Lys | Ile | Gln | Leu | Val | Asn | Thr | Asn | Gly | Ser | Trp | His |
|     |     | 290 |     |     |     | 295 |     |     |     |     | 300 |     |     |     |     |
| Ile | Asn | Arg | Thr | Ala | Leu | Asn | Cys | Asn | Asp | Ser | Leu | Gln | Thr | Gly | Phe |
| 305 |     |     |     |     | 310 |     |     |     |     | 315 |     |     |     |     | 320 |
| Phe | Ala | Ala | Leu | Phe | Tyr | Lys | His | Lys | Phe | Asn | Ser | Ser | Gly | Cys | Pro |
|     |     |     |     | 325 |     |     |     |     | 330 |     |     |     |     | 335 |     |
| Glu | Arg | Leu | Ala | Ser | Cys | Arg | Ser | Ile | Asp | Lys | Phe | Ala | Gln | Gly | Trp |
|     |     |     | 340 |     |     |     |     | 345 |     |     |     |     | 350 |     |     |
| Gly | Pro | Leu | Thr | Tyr | Thr | Glu | Pro | Asn | Ser | Ser | Asp | Gln | Arg | Pro | Tyr |
|     |     | 355 |     |     |     | 360 |     |     |     |     |     | 365 |     |     |     |
| Cys | Trp | His | Tyr | Ala | Pro | Arg | Pro | Cys | Gly | Ile | Val | Pro | Ala | Ser | Gln |

| 370                                                                                | 375 | 380 |
|------------------------------------------------------------------------------------|-----|-----|
| Val Cys Gly Pro Val Tyr Cys Phe Thr Pro Ser Pro Val Val Val Gly<br>385 390 395 400 |     |     |
| Thr Thr Asp Arg Phe Gly Val Pro Thr Tyr Asn Trp Gly Ala Asn Asp<br>405 410 415     |     |     |
| Ser Asp Val Leu Ile Leu Asn Asn Thr Arg Pro Pro Arg Gly Asn Trp<br>420 425 430     |     |     |
| Phe Gly Cys Thr Trp Met Asn Gly Thr Gly Phe Thr Lys Thr Cys Gly<br>435 440 445     |     |     |
| Gly Pro Pro Cys Asn Ile Gly Gly Ala Gly Asn Asn Thr Leu Thr Cys<br>450 455 460     |     |     |
| Pro Thr Asp Cys Phe Arg Lys His Pro Glu Ala Thr Tyr Ala Arg Cys<br>465 470 475 480 |     |     |
| Gly Ser Gly Pro Trp Leu Thr Pro Arg Cys Met Val His Tyr Pro Tyr<br>485 490 495     |     |     |
| Arg Leu Trp His Tyr Pro Cys Thr Val Asn Phe Thr Ile Phe Lys Val<br>500 505 510     |     |     |
| Arg Met Tyr Val Gly Gly Val Glu His Arg Phe Glu Ala Ala Cys Asn<br>515 520 525     |     |     |
| Trp Thr Arg Gly Glu Arg Cys Asp Leu Glu Asp Arg Asp Arg Ser Glu<br>530 535 540     |     |     |
| Leu Ser Pro Leu Leu Leu Ser Thr Thr Glu Trp Gln Ile Leu Pro Cys<br>545 550 555 560 |     |     |
| Ser Phe Thr Thr Leu Pro Ala Leu Ser Thr Gly Leu Ile His Leu His<br>565 570 575     |     |     |
| Gln Asn Ile Val Asp Val Gln Tyr Leu Tyr Gly Val Gly Ser Ala Val<br>580 585 590     |     |     |
| Val Ser Leu Val Ile Lys Trp Glu Tyr Val Leu Leu Leu Phe Leu Leu<br>595 600 605     |     |     |
| Leu Ala Asp Ala Arg Ile Cys Ala Cys Leu Trp Met Met Leu Leu Ile<br>610 615 620     |     |     |
| Ala Gln Ala Glu Ala Ala Leu Glu Asn Leu Val Val Leu Asn Ala Ala<br>625 630 635 640 |     |     |
| Ala Val Ala Gly Ala His Gly Thr Leu Ser Phe Leu Val Phe Phe Cys<br>645 650 655     |     |     |
| Ala Ala Trp Tyr Ile Lys Gly Arg Leu Val Pro Gly Ala Ala Tyr Ala<br>660 665 670     |     |     |
| Phe Tyr Gly Val Trp Pro Leu Leu Leu Leu Leu Ala Leu Pro Pro<br>675 680 685         |     |     |
| Arg Ala Tyr Ala<br>690                                                             |     |     |



## (2) INFORMATION FOR SEQ ID NO: 49:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 2433 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: CDS  
 (B) LOCATION: 1..2430

## (ix) FEATURE:

- (A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 1..2427

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 49:

|                                                                 |     |
|-----------------------------------------------------------------|-----|
| ATG AGC ACG AAT CCT AAA CCT CAA AGA AAA ACC AAA CGT AAC ACC AAC | 48  |
| Met Ser Thr Asn Pro Lys Pro Gln Arg Lys Thr Lys Arg Asn Thr Asn |     |
| 1 5 10 15                                                       |     |
| CGC CGC CCA CAG GAC GTC AAG TTC CCG GGC GGT GGT CAG ATC GTT GGT | 96  |
| Arg Arg Pro Gln Asp Val Lys Phe Pro Gly Gly Gly Gln Ile Val Gly |     |
| 20 25 30                                                        |     |
| GGA GTT TAC CTG TTG CCG CGC AGG GGC CCC AGG TTG GGT GTG CGC GCG | 144 |
| Gly Val Tyr Leu Leu Pro Arg Arg Gly Pro Arg Leu Gly Val Arg Ala |     |
| 35 40 45                                                        |     |
| ACT AGG AAG ACT TCC GAG CGG TCG CAA CCT CGT GGG AGG CGA CAA CCT | 192 |
| Thr Arg Lys Thr Ser Glu Arg Ser Gln Pro Arg Gly Arg Arg Gln Pro |     |
| 50 55 60                                                        |     |
| ATC CCC AAG GCT CGC CGA CCC GAG GGT AGG GCC TGG GCT CAG CCC GGG | 240 |
| Ile Pro Lys Ala Arg Arg Pro Glu Gly Arg Ala Trp Ala Gln Pro Gly |     |
| 65 70 75 80                                                     |     |
| TAC CCT TGG CCC CTC TAT GGC AAT GAG GGC ATG GGG TGG GCA GGA TGG | 288 |
| Tyr Pro Trp Pro Leu Tyr Gly Asn Glu Gly Met Gly Trp Ala Gly Trp |     |
| 85 90 95                                                        |     |
| CTC CTG TCA CCC CGC GGC TCT CGG CCT AGT TGG GGC CCT ACA GAC CCC | 336 |
| Leu Leu Ser Pro Arg Gly Ser Arg Pro Ser Trp Gly Pro Thr Asp Pro |     |
| 100 105 110                                                     |     |
| CGG CGT AGG TCG CGT AAT TTG GGT AAG GTC ATC GAT ACC CTT ACA TGC | 384 |
| Arg Arg Arg Ser Arg Asn Leu Gly Lys Val Ile Asp Thr Leu Thr Cys |     |
| 115 120 125                                                     |     |
| GGC TTC GCC GAC CTC GTG GGG TAC ATT CCG CTC GTC GGC GCC CCC CTA | 432 |
| Gly Phe Ala Asp Leu Val Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu |     |
| 130 135 140                                                     |     |
| GGG GGC GCT GCC AGG GCC CTG GCG CAT GGC GTC CGG GTT CTG GAG GAC | 480 |

|                                                                                                                                                       |      |
|-------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| Gly Gly Ala Ala Arg Ala Leu Ala His Gly Val Arg Val Leu Glu Asp<br>145 150 155 160                                                                    |      |
| GGC GTG AAC TAT GCA ACA GGG AAT TTG CCC GGT TGC TCT TTC TCT ATC<br>Gly Val Asn Tyr Ala Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile<br>165 170 175     | 528  |
| TTC CTC TTG GCT TTG CTG TCC TGT CTG ACC GTT CCA GCT TCC GCT TAT<br>Phe Leu Leu Ala Leu Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr<br>180 185 190     | 576  |
| GAA GTG CGC AAC GTG TCC GGG ATG TAC CAT GTC ACG AAC GAC TGC TCC<br>Glu Val Arg Asn Val Ser Gly Met Tyr His Val Thr Asn Asp Cys Ser<br>195 200 205     | 624  |
| AAC TCA AGC ATT GTG TAT GAG GCA GCG GAC ATG ATC ATG CAC ACC CCC<br>Asn Ser Ile Val Tyr Glu Ala Ala Asp Met Ile Met His Thr Pro<br>210 215 220         | 672  |
| GGG TGC GTG CCC TGC GTT CGG GAG AAC AAC TCT TCC CGC TGC TGG GTA<br>Gly Cys Val Pro Cys Val Arg Glu Asn Asn Ser Ser Arg Cys Trp Val<br>225 230 235 240 | 720  |
| GCG CTC ACC CCC ACG CTC GCA GCT AGG AAC GCC AGC GTC CCC ACC ACG<br>Ala Leu Thr Pro Thr Leu Ala Ala Arg Asn Ala Ser Val Pro Thr Thr<br>245 250 255     | 768  |
| ACA ATA CGA CGC CAC GTC GAT TTG CTC GTT GGG GCG GCT GCT TTC TGT<br>Thr Ile Arg His Val Asp Leu Leu Val Gly Ala Ala Ala Phe Cys<br>260 265 270         | 816  |
| TCC GCT ATG TAC GTG GGG GAC CTC TGC GGA TCT GTC TTC CTC GTC TCC<br>Ser Ala Met Tyr Val Gly Asp Leu Cys Gly Ser Val Phe Leu Val Ser<br>275 280 285     | 864  |
| CAG CTG TTC ACC ATC TCG CCT CGC CGG CAT GAG ACG GTG CAG GAC TGC<br>Gln Leu Phe Thr Ile Ser Pro Arg Arg His Glu Thr Val Gln Asp Cys<br>290 295 300     | 912  |
| AAT TGC TCA ATC TAT CCC GGC CAC ATA ACG GGT CAC CGT ATG GCT TGG<br>Asn Cys Ser Ile Tyr Pro Gly His Ile Thr Gly His Arg Met Ala Trp<br>305 310 315 320 | 960  |
| GAT ATG ATG ATG AAC TGG TCG CCT ACA ACG GCC CTG GTG GTA TCG CAG<br>Asp Met Met Met Asn Trp Ser Pro Thr Thr Ala Leu Val Val Ser Gln<br>325 330 335     | 1008 |
| CTG CTC CGG ATC CCA CAA GCT GTC GTG GAC ATG GTG GCG GGG GCC CAT<br>Leu Leu Arg Ile Pro Gln Ala Val Val Asp Met Val Ala Gly Ala His<br>340 345 350     | 1056 |
| TGG GGA GTC CTG GCG GGC CTC GCC TAC TAT TCC ATG GTG GGG AAC TGG<br>Trp Gly Val Leu Ala Gly Leu Ala Tyr Tyr Ser Met Val Gly Asn Trp<br>355 360 365     | 1104 |
| GCT AAG GTT TTG GTT GTG ATG CTA CTC TTT GCC GGC GTC GAC GGG CAT<br>Ala Lys Val Leu Val Val Met Leu Leu Phe Ala Gly Val Asp Gly His<br>370 375 380     | 1152 |
| ACC CGC GTG TCA GGA GGG GCA GCA GCC TCC GAT ACC AGG GGC CTT GTG<br>Thr Arg Val Ser Gly Gly Ala Ala Ala Ser Asp Thr Arg Gly Leu Val<br>385 390 395 400 | 1200 |

|                                                                                                                                                       |      |
|-------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| TCC CTC TTT AGC CCC GGG TCG GCT CAG AAA ATC CAG CTC GTA AAC ACC<br>Ser Leu Phe Ser Pro Gly Ser Ala Gln Lys Ile Gln Leu Val Asn Thr<br>405 410 415     | 1248 |
| AAC GGC AGT TGG CAC ATC AAC AGG ACT GCC CTG AAC TGC AAC GAC TCC<br>Asn Gly Ser Trp His Ile Asn Arg Thr Ala Leu Asn Cys Asn Asp Ser<br>420 425 430     | 1296 |
| CTC CAA ACA GGG TTC TTT GCC GCA CTA TTC TAC AAA CAC AAA TTC AAC<br>Leu Gln Thr Gly Phe Phe Ala Ala Leu Phe Tyr Lys His Lys Phe Asn<br>435 440 445     | 1344 |
| TCG TCT GGA TGC CCA GAG CGC TTG GCC AGC TGT CGC TCC ATC GAC AAG<br>Ser Ser Gly Cys Pro Glu Arg Leu Ala Ser Cys Arg Ser Ile Asp Lys<br>450 455 460     | 1392 |
| TTC GCT CAG GGG TGG GGT CCC CTC ACT TAC ACT GAG CCT AAC AGC TCG<br>Phe Ala Gln Gly Trp Gly Pro Leu Thr Tyr Thr Glu Pro Asn Ser Ser<br>465 470 475 480 | 1440 |
| GAC CAG AGG CCC TAC TGC TGG CAC TAC GCG CCT CGA CCG TGT GGT ATT<br>Asp Gln Arg Pro Tyr Cys Trp His Tyr Ala Pro Arg Pro Cys Gly Ile<br>485 490 495     | 1488 |
| GTA CCC GCG TCT CAG GTG TGC GGT CCA GTG TAT TGC TTC ACC CCG AGC<br>Val Pro Ala Ser Gln Val Cys Gly Pro Val Tyr Cys Phe Thr Pro Ser<br>500 505 510     | 1536 |
| CCT GTT GTG GTG GGG ACG ACC GAT CGG TTT GGT GTC CCC ACG TAT AAC<br>Pro Val Val Val Gly Thr Thr Asp Arg Phe Gly Val Pro Thr Tyr Asn<br>515 520 525     | 1584 |
| TGG GGG GCG AAC GAC TCG GAT GTG CTG ATT CTC AAC AAC ACG CGG CCG<br>Trp Gly Ala Asn Asp Ser Asp Val Leu Ile Leu Asn Asn Thr Arg Pro<br>530 535 540     | 1632 |
| CCG CGA GGC AAC TGG TTC GGC TGT ACA TGG ATG AAT GGC ACT GGG TTC<br>Pro Arg Gly Asn Trp Phe Gly Cys Thr Trp Met Asn Gly Thr Gly Phe<br>545 550 555 560 | 1680 |
| ACC AAG ACG TGT GGG GGC CCC CCG TGC AAC ATC GGG GGG GCC GGC AAC<br>Thr Lys Thr Cys Gly Gly Pro Pro Cys Asn Ile Gly Gly Ala Gly Asn<br>565 570 575     | 1728 |
| AAC ACC TTG ACC TGC CCC ACT GAC TGT TTT CGG AAG CAC CCC GAG GCC<br>Asn Thr Leu Thr Cys Pro Thr Asp Cys Phe Arg Lys His Pro Glu Ala<br>580 585 590     | 1776 |
| ACC TAC GCC AGA TGC GGT TCT GGG CCC TGG CTG ACA CCT AGG TGT ATG<br>Thr Tyr Ala Arg Cys Gly Ser Gly Pro Trp Leu Thr Pro Arg Cys Met<br>595 600 605     | 1824 |
| GTT CAT TAC CCA TAT AGG CTC TGG CAC TAC CCC TGC ACT GTC AAC TTC<br>Val His Tyr Pro Tyr Arg Leu Trp His Tyr Pro Cys Thr Val Asn Phe<br>610 615 620     | 1872 |
| ACC ATC TTC AAG GTT AGG ATG TAC GTG GGG GGC GTG GAG CAC AGG TTC<br>Thr Ile Phe Lys Val Arg Met Tyr Val Gly Gly Val Glu His Arg Phe<br>625 630 635 640 | 1920 |
| GAA GCC GCA TGC AAT TGG ACT CGA GGA GAG CGT TGT GAC TTG GAG GAC                                                                                       | 1968 |

|     |     |     |     |     |     |     |     |     |        |     |     |     |     |     |     |      |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|--------|-----|-----|-----|-----|-----|-----|------|
| Glu | Ala | Ala | Cys | Asn | Trp | Thr | Arg | Gly | Glu    | Arg | Cys | Asp | Leu | Glu | Asp |      |
|     |     |     |     | 645 |     |     |     |     | 650    |     |     |     |     | 655 |     |      |
| AGG | GAT | AGA | TCA | GAG | CTT | AGC | CCG | CTG | CTG    | CTG | TCT | ACA | ACA | GAG | TGG | 2016 |
| Arg | Asp | Arg | Ser | Glu | Leu | Ser | Pro | Leu | Leu    | Leu | Ser | Thr | Thr | Glu | Trp |      |
|     |     |     | 660 |     |     |     |     | 665 |        |     |     |     |     | 670 |     |      |
| CAG | ATA | CTG | CCC | TGT | TCC | TTC | ACC | ACC | CTG    | CCG | GCC | CTA | TCC | ACC | GGC | 2064 |
| Gln | Ile | Leu | Pro | Cys | Ser | Phe | Thr | Thr | Leu    | Pro | Ala | Leu | Ser | Thr | Gly |      |
|     |     | 675 |     |     |     |     | 680 |     |        |     |     | 685 |     |     |     |      |
| CTG | ATC | CAC | CTC | CAT | CAG | AAC | ATC | GTG | GAC    | GTG | CAA | TAC | CTG | TAC | GGT | 2112 |
| Leu | Ile | His | Leu | His | Gln | Asn | Ile | Val | Asp    | Val | Gln | Tyr | Leu | Tyr | Gly |      |
|     | 690 |     |     |     |     | 695 |     |     |        | 700 |     |     |     |     |     |      |
| GTA | GGG | TCG | GCG | GTT | GTC | TCC | CTT | GTC | ATC    | AAA | TGG | GAG | TAT | GTC | CTG | 2160 |
| Val | Gly | Ser | Ala | Val | Val | Ser | Leu | Val | Ile    | Lys | Trp | Glu | Tyr | Val | Leu |      |
| 705 |     |     |     | 710 |     |     |     |     |        | 715 |     |     |     | 720 |     |      |
| TTG | CTC | TTC | CTT | CTC | CTG | GCA | GAC | GCG | CGC    | ATC | TGC | GCC | TGC | TTA | TGG | 2208 |
| Leu | Leu | Phe | Leu | Leu | Leu | Ala | Asp | Ala | Arg    | Ile | Cys | Ala | Cys | Leu | Trp |      |
|     |     |     |     | 725 |     |     |     | 730 |        |     |     |     |     | 735 |     |      |
| ATG | ATG | CTG | CTG | ATA | GCT | CAA | GCT | GAG | GCC    | GCC | TTA | GAG | AAC | CTG | GTG | 2256 |
| Met | Met | Leu | Leu | Ile | Ala | Gln | Ala | Glu | Ala    | Ala | Leu | Glu | Asn | Leu | Val |      |
|     |     |     | 740 |     |     |     | 745 |     |        |     |     |     | 750 |     |     |      |
| GTC | CTC | AAT | GCG | GCG | GCC | GTG | GCC | GGG | GCG    | CAT | GGC | ACT | CTT | TCC | TTC | 2304 |
| Val | Leu | Asn | Ala | Ala | Ala | Val | Ala | Gly | Ala    | His | Gly | Thr | Leu | Ser | Phe |      |
|     | 755 |     |     |     |     | 760 |     |     |        |     | 765 |     |     |     |     |      |
| CTT | GTG | TTC | TTC | TGT | GCT | GCC | TGG | TAC | ATC    | AAG | GGC | AGG | CTG | GTC | CCT | 2352 |
| Leu | Val | Phe | Phe | Cys | Ala | Ala | Trp | Tyr | Ile    | Lys | Gly | Arg | Leu | Val | Pro |      |
|     | 770 |     |     |     |     | 775 |     |     |        |     | 780 |     |     |     |     |      |
| GGT | GCG | GCA | TAC | GCC | TTC | TAT | GGC | GTG | TGG    | CCG | CTG | CTC | CTG | CTT | CTG | 2400 |
| Gly | Ala | Ala | Tyr | Ala | Phe | Tyr | Gly | Val | Trp    | Pro | Leu | Leu | Leu | Leu | Leu |      |
| 785 |     |     |     | 790 |     |     |     |     | 795    |     |     |     |     | 800 |     |      |
| CTG | GCC | TTA | CCA | CCA | CGA | GCT | TAT | GCC | TAGTAA |     |     |     |     |     |     | 2433 |
| Leu | Ala | Leu | Pro | Pro | Arg | Ala | Tyr | Ala |        |     |     |     |     |     |     |      |
|     |     |     | 805 |     |     |     |     | 810 |        |     |     |     |     |     |     |      |

## (2) INFORMATION FOR SEQ ID NO: 50:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 809 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 50:

Met Ser Thr Asn Pro Lys Pro Gln Arg Lys Thr Lys Arg Asn Thr Asn  
 1 5 10 15  
 Arg Arg Pro Gln Asp Val Lys Phe Pro Gly Gly Gly Gln Ile Val Gly  
 20 25 30  
 Gly Val Tyr Leu Leu Pro Arg Arg Gly Pro Arg Leu Gly Val Arg Ala

| 35  |     |     |     |     | 40  |     |     |     |     | 45  |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Thr | Arg | Lys | Thr | Ser | Glu | Arg | Ser | Gln | Pro | Arg | Gly | Arg | Arg | Gln | Pro |
| 50  |     |     |     |     |     | 55  |     |     |     |     | 60  |     |     |     |     |
| Ile | Pro | Lys | Ala | Arg | Arg | Pro | Glu | Gly | Arg | Ala | Trp | Ala | Gln | Pro | Gly |
| 65  |     |     |     |     | 70  |     |     |     |     | 75  |     |     |     |     | 80  |
| Tyr | Pro | Trp | Pro | Leu | Tyr | Gly | Asn | Glu | Gly | Met | Gly | Trp | Ala | Gly | Trp |
|     |     |     |     | 85  |     |     |     |     | 90  |     |     |     |     | 95  |     |
| Leu | Leu | Ser | Pro | Arg | Gly | Ser | Arg | Pro | Ser | Trp | Gly | Pro | Thr | Asp | Pro |
|     |     |     | 100 |     |     |     |     | 105 |     |     |     |     | 110 |     |     |
| Arg | Arg | Arg | Ser | Arg | Asn | Leu | Gly | Lys | Val | Ile | Asp | Thr | Leu | Thr | Cys |
|     |     | 115 |     |     |     | 120 |     |     |     |     | 125 |     |     |     |     |
| Gly | Phe | Ala | Asp | Leu | Val | Gly | Tyr | Ile | Pro | Leu | Val | Gly | Ala | Pro | Leu |
|     | 130 |     |     |     |     | 135 |     |     |     |     | 140 |     |     |     |     |
| Gly | Gly | Ala | Ala | Arg | Ala | Leu | Ala | His | Gly | Val | Arg | Val | Leu | Glu | Asp |
| 145 |     |     |     |     | 150 |     |     |     |     | 155 |     |     |     |     | 160 |
| Gly | Val | Asn | Tyr | Ala | Thr | Gly | Asn | Leu | Pro | Gly | Cys | Ser | Phe | Ser | Ile |
|     |     |     |     | 165 |     |     |     |     | 170 |     |     |     |     | 175 |     |
| Phe | Leu | Leu | Ala | Leu | Leu | Ser | Cys | Leu | Thr | Val | Pro | Ala | Ser | Ala | Tyr |
|     |     |     | 180 |     |     |     |     | 185 |     |     |     |     |     | 190 |     |
| Glu | Val | Arg | Asn | Val | Ser | Gly | Met | Tyr | His | Val | Thr | Asn | Asp | Cys | Ser |
|     |     | 195 |     |     |     |     | 200 |     |     |     |     | 205 |     |     |     |
| Asn | Ser | Ser | Ile | Val | Tyr | Glu | Ala | Ala | Asp | Met | Ile | Met | His | Thr | Pro |
|     |     | 210 |     |     |     | 215 |     |     |     |     | 220 |     |     |     |     |
| Gly | Cys | Val | Pro | Cys | Val | Arg | Glu | Asn | Asn | Ser | Ser | Arg | Cys | Trp | Val |
| 225 |     |     |     |     | 230 |     |     |     |     | 235 |     |     |     |     | 240 |
| Ala | Leu | Thr | Pro | Thr | Leu | Ala | Ala | Arg | Asn | Ala | Ser | Val | Pro | Thr | Thr |
|     |     |     |     | 245 |     |     |     |     | 250 |     |     |     |     | 255 |     |
| Thr | Ile | Arg | Arg | His | Val | Asp | Leu | Leu | Val | Gly | Ala | Ala | Ala | Phe | Cys |
|     |     |     | 260 |     |     |     | 265 |     |     |     |     |     | 270 |     |     |
| Ser | Ala | Met | Tyr | Val | Gly | Asp | Leu | Cys | Gly | Ser | Val | Phe | Leu | Val | Ser |
|     |     | 275 |     |     |     |     | 280 |     |     |     |     | 285 |     |     |     |
| Gln | Leu | Phe | Thr | Ile | Ser | Pro | Arg | Arg | His | Glu | Thr | Val | Gln | Asp | Cys |
|     | 290 |     |     |     |     | 295 |     |     |     |     | 300 |     |     |     |     |
| Asn | Cys | Ser | Ile | Tyr | Pro | Gly | His | Ile | Thr | Gly | His | Arg | Met | Ala | Trp |
| 305 |     |     |     |     | 310 |     |     |     |     | 315 |     |     |     |     | 320 |
| Asp | Met | Met | Met | Asn | Trp | Ser | Pro | Thr | Thr | Ala | Leu | Val | Val | Ser | Gln |
|     |     |     |     | 325 |     |     |     |     | 330 |     |     |     |     | 335 |     |
| Leu | Leu | Arg | Ile | Pro | Gln | Ala | Val | Val | Asp | Met | Val | Ala | Gly | Ala | His |
|     |     |     | 340 |     |     |     |     | 345 |     |     |     |     | 350 |     |     |
| Trp | Gly | Val | Leu | Ala | Gly | Leu | Ala | Tyr | Tyr | Ser | Met | Val | Gly | Asn | Trp |
|     | 355 |     |     |     |     |     | 360 |     |     |     |     | 365 |     |     |     |
| Ala | Lys | Val | Leu | Val | Val | Met | Leu | Leu | Phe | Ala | Gly | Val | Asp | Gly | His |

| 370                                                                                | 375 | 380 |
|------------------------------------------------------------------------------------|-----|-----|
| Thr Arg Val Ser Gly Gly Ala Ala Ala Ser Asp Thr Arg Gly Leu Val<br>385 390 395 400 |     |     |
| Ser Leu Phe Ser Pro Gly Ser Ala Gln Lys Ile Gln Leu Val Asn Thr<br>405 410 415     |     |     |
| Asn Gly Ser Trp His Ile Asn Arg Thr Ala Leu Asn Cys Asn Asp Ser<br>420 425 430     |     |     |
| Leu Gln Thr Gly Phe Phe Ala Ala Leu Phe Tyr Lys His Lys Phe Asn<br>435 440 445     |     |     |
| Ser Ser Gly Cys Pro Glu Arg Leu Ala Ser Cys Arg Ser Ile Asp Lys<br>450 455 460     |     |     |
| Phe Ala Gln Gly Trp Gly Pro Leu Thr Tyr Thr Glu Pro Asn Ser Ser<br>465 470 475 480 |     |     |
| Asp Gln Arg Pro Tyr Cys Trp His Tyr Ala Pro Arg Pro Cys Gly Ile<br>485 490 495     |     |     |
| Val Pro Ala Ser Gln Val Cys Gly Pro Val Tyr Cys Phe Thr Pro Ser<br>500 505 510     |     |     |
| Pro Val Val Val Gly Thr Thr Asp Arg Phe Gly Val Pro Thr Tyr Asn<br>515 520 525     |     |     |
| Trp Gly Ala Asn Asp Ser Asp Val Leu Ile Leu Asn Asn Thr Arg Pro<br>530 535 540     |     |     |
| Pro Arg Gly Asn Trp Phe Gly Cys Thr Trp Met Asn Gly Thr Gly Phe<br>545 550 555 560 |     |     |
| Thr Lys Thr Cys Gly Gly Pro Pro Cys Asn Ile Gly Gly Ala Gly Asn<br>565 570 575     |     |     |
| Asn Thr Leu Thr Cys Pro Thr Asp Cys Phe Arg Lys His Pro Glu Ala<br>580 585 590     |     |     |
| Thr Tyr Ala Arg Cys Gly Ser Gly Pro Trp Leu Thr Pro Arg Cys Met<br>595 600 605     |     |     |
| Val His Tyr Pro Tyr Arg Leu Trp His Tyr Pro Cys Thr Val Asn Phe<br>610 615 620     |     |     |
| Thr Ile Phe Lys Val Arg Met Tyr Val Gly Gly Val Glu His Arg Phe<br>625 630 635 640 |     |     |
| Glu Ala Ala Cys Asn Trp Thr Arg Gly Glu Arg Cys Asp Leu Glu Asp<br>645 650 655     |     |     |
| Arg Asp Arg Ser Glu Leu Ser Pro Leu Leu Leu Ser Thr Thr Glu Trp<br>660 665 670     |     |     |
| Gln Ile Leu Pro Cys Ser Phe Thr Thr Leu Pro Ala Leu Ser Thr Gly<br>675 680 685     |     |     |
| Leu Ile His Leu His Gln Asn Ile Val Asp Val Gln Tyr Leu Tyr Gly<br>690 695 700     |     |     |

Val Gly Ser Ala Val Val Ser Leu Val Ile Lys Trp Glu Tyr Val Leu  
 705 710 715 720  
 Leu Leu Phe Leu Leu Leu Ala Asp Ala Arg Ile Cys Ala Cys Leu Trp  
 725 730 735  
 Met Met Leu Leu Ile Ala Gln Ala Glu Ala Ala Leu Glu Asn Leu Val  
 740 745 750  
 Val Leu Asn Ala Ala Ala Val Ala Gly Ala His Gly Thr Leu Ser Phe  
 755 760 765  
 Leu Val Phe Phe Cys Ala Ala Trp Tyr Ile Lys Gly Arg Leu Val Pro  
 770 775 780  
 Gly Ala Ala Tyr Ala Phe Tyr Gly Val Trp Pro Leu Leu Leu Leu Leu  
 785 790 795 800  
 Leu Ala Leu Pro Pro Arg Ala Tyr Ala  
 805

## (2) INFORMATION FOR SEQ ID NO: 51:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 17 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide

- (ix) FEATURE:
  - (A) NAME/KEY: Modified-site
  - (B) LOCATION: 1..17

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 51:

Ser Asn Ser Ser Glu Ala Ala Asp Met Ile Met His Thr Pro Gly Cys  
 1 5 10 15  
 Val

## (2) INFORMATION FOR SEQ ID NO: 52:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 22 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide

- (ix) FEATURE:
  - (A) NAME/KEY: Modified-site
  - (B) LOCATION: 1..22

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 52:

Gly Gly Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp  
 1                      5                      10                      15  
 Ser Pro Thr Thr Ala Leu  
                     20

## (2) INFORMATION FOR SEQ ID NO: 53:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 37 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- (ix) FEATURE:  
 (A) NAME/KEY: Modified-site  
 (B) LOCATION: 1..37

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 53:

Tyr Glu Val Arg Asn Val Ser Gly Ile Tyr His Val Thr Asn Asp Cys  
 1                      5                      10                      15  
 Ser Asn Ser Ser Ile Val Tyr Glu Ala Ala Asp Met Ile Met His Thr  
                     20                      25                      30  
 Pro Gly Cys Gly Lys  
                     35

## (2) INFORMATION FOR SEQ ID NO: 54:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 25 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- (ix) FEATURE:  
 (A) NAME/KEY: Modified-site  
 (B) LOCATION: 1..25

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 54:

Gly Gly Thr Pro Thr Val Ala Thr Arg Asp Gly Lys Leu Pro Ala Thr  
 1                      5                      10                      15  
 Gln Leu Arg Arg His Ile Asp Leu Leu  
                     20                      25

## (2) INFORMATION FOR SEQ ID NO: 55:

- (i) SEQUENCE CHARACTERISTICS:



(A) LENGTH: 25 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

```
(A) NAME/KEY: Modified-site
(B) LOCATION: 1..25
```

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 55:

Gly Gly Thr Pro Thr Leu Ala Ala Arg Asp Ala Ser Val Pro Thr Thr  
1 5 10 15  
Thr Ile Arg Arg His Val Asp Leu Leu  
20 25

(2) INFORMATION FOR SEQ ID NO: 56:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 56:

Leu Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr Gln Val Arg Asn  
1 5 10 15  
Ser Thr Gly Leu  
20

(2) INFORMATION FOR SEQ ID NO: 57:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 57:

Gln Val Arg Asn Ser Thr Gly Leu Tyr His Val Thr Asn Asp Cys Pro  
1 5 10 15  
Asn Ser Ser Ile  
20

## (2) INFORMATION FOR SEQ ID NO: 58:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 58:

Asn Asp Cys Pro Asn Ser Ser Ile Val Tyr Glu Ala His Asp Ala Ile  
1                      5                      10                      15  
Leu His Thr Pro  
                    20

## (2) INFORMATION FOR SEQ ID NO: 59:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 59:

Ser Asn Ser Ser Ile Val Tyr Glu Ala Ala Asp Met Ile Met His Thr  
1                      5                      10                      15  
Pro Gly Cys Val  
                    20

## (2) INFORMATION FOR SEQ ID NO: 60:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 19 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 60:

His Asp Ala Ile Leu His Thr Pro Gly Val Pro Cys Val Arg Glu Gly  
1                      5                      10                      15  
Asn Val Ser

## (2) INFORMATION FOR SEQ ID NO: 61:

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 61:

Cys Val Arg Glu Gly Asn Val Ser Arg Cys Trp Val Ala Met Thr Pro  
1                      5                      10                      15  
Thr Val Ala Thr  
                    20

(2) INFORMATION FOR SEQ ID NO: 62:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 62:

Ala Met Thr Pro Thr Val Ala Thr Arg Asp Gly Lys Leu Pro Ala Thr  
1                      5                      10                      15  
Gln Leu Arg Arg  
                    20

(2) INFORMATION FOR SEQ ID NO: 63:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 63:

Leu Pro Ala Thr Gln Leu Arg Arg His Ile Asp Leu Leu Val Gly Ser  
1                      5                      10                      15  
Ala Thr Leu Cys  
                    20

(2) INFORMATION FOR SEQ ID NO: 64:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 64:

Leu Val Gly Ser Ala Thr Leu Cys Ser Ala Leu Tyr Val Gly Asp Leu  
1 5 10 15  
Cys Gly Ser Val  
20

(2) INFORMATION FOR SEQ ID NO: 65:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 65:

Gln Leu Phe Thr Phe Ser Pro Arg Arg His Trp Thr Thr Gln Gly Cys  
1 5 10 15  
Asn Cys Ser Ile  
20

(2) INFORMATION FOR SEQ ID NO: 66:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 66:

Thr Gln Gly Cys Asn Cys Ser Ile Tyr Pro Gly His Ile Thr Gly His  
1 5 10 15  
Arg Met Ala Trp  
20

(2) INFORMATION FOR SEQ ID NO: 67:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 67:

Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp Ser Pro  
1 5 10 15

Thr Ala Ala Leu  
20

(2) INFORMATION FOR SEQ ID NO: 68:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 68:

Asn Trp Ser Pro Thr Ala Ala Leu Val Met Ala Gln Leu Leu Arg Ile  
1 5 10 15

Pro Gln Ala Ile  
20

(2) INFORMATION FOR SEQ ID NO: 69:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 69:

Leu Leu Arg Ile Pro Gln Ala Ile Leu Asp Met Ile Ala Gly Ala His  
1 5 10 15

Trp Gly Val Leu  
20

(2) INFORMATION FOR SEQ ID NO: 70:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 70:

Ala Gly Ala His Trp Gly Val Leu Ala Gly Ile Ala Tyr Phe Ser Met  
1 5 10 15  
Val Gly Asn Met  
20

(2) INFORMATION FOR SEQ ID NO: 71:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 71:

Val Val Leu Leu Leu Phe Ala Gly Val Asp Ala Glu Thr Ile Val Ser  
1 5 10 15  
Gly Gly Gln Ala  
20

(2) INFORMATION FOR SEQ ID NO: 72:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 72:

Ser Gly Leu Val Ser Leu Phe Thr Pro Gly Ala Lys Gln Asn Ile Gln  
1 5 10 15  
Leu Ile Asn Thr  
20

(2) INFORMATION FOR SEQ ID NO: 73:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 73:

Gln Asn Ile Gln Leu Ile Asn Thr Asn Gly Gln Trp His Ile Asn Ser  
 1 5 10 15  
 Thr Ala Leu Asn  
 20

(2) INFORMATION FOR SEQ ID NO: 74:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 21 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 74:

Leu Asn Cys Asn Glu Ser Leu Asn Thr Gly Trp Trp Leu Ala Gly Leu  
 1 5 10 15  
 Ile Tyr Gln His Lys  
 20

(2) INFORMATION FOR SEQ ID NO: 75:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 75:

Ala Gly Leu Ile Tyr Gln His Lys Phe Asn Ser Ser Gly Cys Pro Glu  
 1 5 10 15  
 Arg Leu Ala Ser  
 20

(2) INFORMATION FOR SEQ ID NO: 76:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 76:

Gly Cys Pro Glu Arg Leu Ala Ser Cys Arg Pro Leu Thr Asp Phe Asp

|                 |    |    |    |
|-----------------|----|----|----|
| 1               | 5  | 10 | 15 |
| Gln Gly Trp Gly |    |    |    |
|                 | 20 |    |    |

(2) INFORMATION FOR SEQ ID NO: 77:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 77:

Thr Asp Phe Asp Gln Gly Trp Gly Pro Ile Ser Tyr Ala Asn Gly Ser  
1 5 10 15  
Gly Pro Asp Gln  
20

(2) INFORMATION FOR SEO ID NO: 78:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 78:

Ala Asn Gly Ser Gly Pro Asp Gln Arg Pro Tyr Cys Trp His Tyr Pro  
1 5 10 15  
Pro Lys Pro Cys  
20

(2) INFORMATION FOR SEQ ID NO: 79:

(i) SEQUENCE CHARACTERISTICS:

- SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 79:

Trp His Tyr Pro Pro Lys Pro Cys Gly Ile Val Pro Ala Lys Ser Val  
1 5 10 15  
Cys Gly Pro Val



20

## (2) INFORMATION FOR SEQ ID NO: 80:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 80:

|                 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Ala             | Lys | Ser | Val | Cys | Gly | Pro | Val | Tyr | Cys | Phe | Thr | Pro | Ser | Pro | Val |
| 1               |     |     |     | 5   |     |     |     |     | 10  |     |     |     |     | 15  |     |
| Val Val Gly Thr |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
| 20              |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |

## (2) INFORMATION FOR SEQ ID NO: 81:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 81:

|                 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Pro             | Ser | Pro | Val | Val | Val | Gly | Thr | Thr | Asp | Arg | Ser | Gly | Ala | Pro | Thr |
| 1               |     |     |     | 5   |     |     |     |     | 10  |     |     |     |     | 15  |     |
| Tyr Ser Trp Gly |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
| 20              |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |

## (2) INFORMATION FOR SEQ ID NO: 82:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 82:

|                 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Gly             | Ala | Pro | Thr | Tyr | Ser | Trp | Gly | Glu | Asn | Asp | Thr | Asp | Val | Phe | Val |
| 1               |     |     |     | 5   |     |     |     |     | 10  |     |     |     |     | 15  |     |
| Leu Asn Asn Thr |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
| 20              |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |

## (2) INFORMATION FOR SEQ ID NO: 83:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 83:
- Gly Asn Trp Phe Gly Cys Thr Trp Met Asn Ser Thr Gly Phe Thr Lys  
1 5 10 15  
Val Cys Gly Ala  
20
- (2) INFORMATION FOR SEQ ID NO: 84:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 84:
- Gly Phe Thr Lys Val Cys Gly Ala Pro Pro Val Cys Ile Gly Gly Ala  
1 5 10 15  
Gly Asn Asn Thr  
20
- (2) INFORMATION FOR SEQ ID NO: 85:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 19 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 85:
- Ile Gly Gly Ala Gly Asn Asn Thr Leu His Cys Pro Thr Asp Cys Arg  
1 5 10 15  
Lys His Pro
- (2) INFORMATION FOR SEQ ID NO: 86:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids

- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 86:

Thr Asp Cys Phe Arg Lys His Pro Asp Ala Thr Tyr Ser Arg Cys Gly  
1                      5                      10                      15

Ser Gly Pro Trp  
20

(2) INFORMATION FOR SEQ ID NO: 87:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 87:

Ser Arg Cys Gly Ser Gly Pro Trp Ile Thr Pro Arg Cys Leu Val Asp  
1                      5                      10                      15

Tyr Pro Tyr Arg  
20

(2) INFORMATION FOR SEQ ID NO: 88:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 88:

Cys Leu Val Asp Tyr Pro Tyr Arg Leu Trp His Tyr Pro Cys Thr Ile  
1                      5                      10                      15

Asn Tyr Thr Ile  
20

(2) INFORMATION FOR SEQ ID NO: 89:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 89:

Pro Cys Thr Ile Asn Tyr Thr Ile Phe Lys Ile Arg Met Tyr Val Gly  
1                   5                   10                   15  
Gly Val Glu His  
20

(2) INFORMATION FOR SEQ ID NO: 90:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 90:

Met Tyr Val Gly Gly Val Glu His Arg Leu Glu Ala Ala Cys Asn Trp  
1                   5                   10                   15  
Thr Pro Gly Glu  
20

(2) INFORMATION FOR SEQ ID NO: 91:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 91:

Ala Cys Asn Trp Thr Pro Gly Glu Arg Cys Asp Leu Glu Asp Arg Asp  
1                   5                   10                   15  
Arg Ser Glu Leu  
20

(2) INFORMATION FOR SEQ ID NO: 92:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 92:

|                 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Glu             | Asp | Arg | Asp | Arg | Ser | Glu | Leu | Ser | Pro | Leu | Leu | Leu | Thr | Thr | Thr |
| 1               |     |     |     | 5   |     |     |     | 10  |     |     |     |     | 15  |     |     |
| Gln Trp Gln Val |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
| 20              |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |

(2) INFORMATION FOR SEQ ID NO: 93:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 9 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 93:

|     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Tyr | Gln | Val | Arg | Asn | Ser | Thr | Gly | Leu |
| 1   |     |     |     | 5   |     |     |     |     |

(2) INFORMATION FOR SEQ ID NO: 94:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 29 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 94:

ACGTCCGTAC GTTCGAATTA ATTAATCGA

29

(2) INFORMATION FOR SEQ ID NO: 95:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 60 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 95:

CCTCCGGACG TGCAC TAGCT CCCGTCTGTG GTAGTGGTGG TAGTGATTAT CAATTAATTG 60

(2) INFORMATION FOR SEQ ID NO: 96:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 19 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 96:

GTTTAACCAC TGCATGATG 19

(2) INFORMATION FOR SEQ ID NO: 97:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 97:

GTCCCATCGA GTGCGGCTAC 20

(2) INFORMATION FOR SEQ ID NO: 98:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 45 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 98:

CGTGACATGG TACATTCCGG ACACTTGGCG CACTTCATAA GCGGA

45

(2) INFORMATION FOR SEQ ID NO: 99:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 99:

TGCCTCATAC ACAATGGAGC TCTGGGACGA GTCGTTCGTG AC

42

(2) INFORMATION FOR SEQ ID NO: 100:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 100:

TACCCAGCAG CGGGAGCTCT GTTGCTCCCG AACGCAGGGC AC

42

(2) INFORMATION FOR SEQ ID NO: 101:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 101:

TGTCGTGGTG GGGACGGAGG CCTGCCTAGC TGCGAGCGTG GG

42

(2) INFORMATION FOR SEQ ID NO: 102:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 48 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 102:

CGTTATGTGG CCCGGGTAGA TTGAGCACTG GCAGTCCTGC ACCGTCTC

48

(2) INFORMATION FOR SEQ ID NO: 103:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 42 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 103:

CAGGGCCGTT CTAGGCCTCC ACTGCATCAT CATATCCCAA GC

42

(2) INFORMATION FOR SEQ ID NO: 104:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 104:

CCGGAATGTA CCATGTCACG AACGAC

26



## (2) INFORMATION FOR SEQ ID NO: 105:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 24 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (iii) HYPOTHETICAL: NO

- (iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 105:  
GCTCCATTGT GTATGAGGCA GCGG

24

## (2) INFORMATION FOR SEQ ID NO: 106:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 23 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (iii) HYPOTHETICAL: NO

- (iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 106:  
GAGCTCCCGC TGCTGGGTAG CGC

23

## (2) INFORMATION FOR SEQ ID NO: 107:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 25 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (iii) HYPOTHETICAL: NO

- (iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 107:  
CCTCCGTCCC CACCACGACA ATACG

25

## (2) INFORMATION FOR SEQ ID NO: 108:

- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 27 base pairs
    - (B) TYPE: nucleic acid
    - (C) STRANDEDNESS: single
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: DNA (genomic)
  - (iii) HYPOTHETICAL: NO
  - (iii) ANTI-SENSE: NO
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 108:
- CTACCCGGGC CACATAACGG GTCACCG 27
- (2) INFORMATION FOR SEQ ID NO: 109:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 24 base pairs
    - (B) TYPE: nucleic acid
    - (C) STRANDEDNESS: single
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: DNA (genomic)
  - (iii) HYPOTHETICAL: NO
  - (iii) ANTI-SENSE: NO
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 109:
- GGAGGCCTAC AACGGCCCTG GTGG 24
- (2) INFORMATION FOR SEQ ID NO: 110:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 22 base pairs
    - (B) TYPE: nucleic acid
    - (C) STRANDEDNESS: single
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: DNA (genomic)
  - (iii) HYPOTHETICAL: NO
  - (iii) ANTI-SENSE: NO
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 110:
- TTCTATCGAT TAAATAGAAT TC 22
- (2) INFORMATION FOR SEQ ID NO: 111:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 23 base pairs
    - (B) TYPE: nucleic acid

(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 111:

GCCATACGCT CACAGCCGAT CCC

## (2) INFORMATION FOR SEQ ID NO: 112:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 112:

|                 |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Tyr             | Glu | Val | Arg | Asn | Val | Ser | Gly | Ile | Tyr | His | Val | Thr | Asn | Asp | Cys |
| 1               |     |     |     | 5   |     |     |     | 10  |     |     |     |     |     | 15  |     |
| Ser Asn Ser Ser |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
| 20              |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |

## (2) INFORMATION FOR SEQ ID NO: 113:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 113:

|                     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|---------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Thr                 | Asn | Asp | Cys | Ser | Asn | Ser | Ser | Ile | Val | Tyr | Glu | Ala | Ala | Asp |
| 1                   |     |     |     | 5   |     |     |     | 10  |     |     |     |     |     | 15  |
| Met Ile Met His Thr |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
| 20                  |     |     |     |     |     |     |     |     |     |     |     |     |     |     |

## (2) INFORMATION FOR SEQ ID NO: 114:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 114:

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Ala | Ala | Asp | Met | Ile | Met | His | Thr | Pro | Gly | Cys | Val | Pro | Cys | Val |
| 1   |     |     |     | 5   |     |     |     | 10  |     |     |     |     |     | 15  |

Arg Glu Asn Asn Ser  
20

(2) INFORMATION FOR SEQ ID NO: 115:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 115:

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Pro | Cys | Val | Arg | Glu | Asn | Asn | Ser | Ser | Arg | Cys | Trp | Val | Ala | Leu |
| 1   |     |     |     | 5   |     |     |     | 10  |     |     |     |     |     | 15  |
|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
| Thr | Pro | Thr | Leu | Ala |     |     |     |     |     |     |     |     |     |     |
|     |     |     |     |     | 20  |     |     |     |     |     |     |     |     |     |

(2) INFORMATION FOR SEQ ID NO: 116:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 116:

|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Val | Ala | Leu | Thr | Pro | Thr | Leu | Ala | Ala | Arg | Asn | Ala | Ser | Val | Pro |
| 1   |     |     |     | 5   |     |     |     |     | 10  |     |     |     |     | 15  |
|     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
| Thr | Thr | Thr | Ile | Arg |     |     |     |     |     |     |     |     |     |     |
|     |     |     |     |     | 20  |     |     |     |     |     |     |     |     |     |

(2) INFORMATION FOR SEQ ID NO: 117:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 117:

```

Ser Val Pro Thr Thr Thr Ile Arg Arg His Val Asp Leu Leu Val
 1             5             10             15
Gly Ala Ala Ala Phe
                20

```

(2) INFORMATION FOR SEQ ID NO: 118:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 118:

```

Leu Leu Val Gly Ala Ala Ala Phe Cys Ser Ala Met Tyr Val Gly
 1             5             10             15
Asp Leu Cys Gly Ser
                20

```

(2) INFORMATION FOR SEQ ID NO: 119:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 119:

```

Tyr Val Gly Asp Leu Cys Gly Ser Val Phe Leu Val Ser Gln Leu
 1             5             10             15
Phe Thr Ile Ser Pro
                20

```

(2) INFORMATION FOR SEQ ID NO: 120:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 120:

Ser Gln Leu Phe Thr Ile Ser Pro Arg Arg His Glu Thr Val Gln  
1 5 10 15  
Asp Cys Asn Cys Ser  
20

(2) INFORMATION FOR SEQ ID NO: 121:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 121:

Thr Val Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His Ile Thr  
1 5 10 15  
Gly His Arg Met Ala  
20

(2) INFORMATION FOR SEQ ID NO: 122:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 122:

His Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp  
1 5 10 15  
Ser Pro Thr Thr Ala  
20

|                                          |                                                           |
|------------------------------------------|-----------------------------------------------------------|
| Applicant's or agent's<br>file reference | International application No.<br><b>PCT/EP 02 / 00219</b> |
|------------------------------------------|-----------------------------------------------------------|

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM  
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

|                                                                                                                                                                            |                                  |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------|
| A. The indications made below relate to the deposited microorganism or other biological material referred to in the description<br>on page <u>40</u> , line <u>33-37</u>   |                                  |
| B. IDENTIFICATION OF DEPOSIT <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>                            |                                  |
| Name of depositary institution<br>BCCM/LMBP (Belgian Coordinated Collections of Microorganisms/Laboratorium voor Moleculaire<br>Biologie-Plasmidencollectie)               |                                  |
| Address of depositary institution (including postal code and country)<br>Universiteit Gent<br>K.L. Ledeganckstraat 35<br>B-9000 Ghent<br>BELGIUM                           |                                  |
| Date of deposit<br><br>January 9, 2002                                                                                                                                     | Accession Number<br><br>LMBP4486 |
| C. ADDITIONAL INDICATIONS (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span> |                                  |
|                                                                                                                                                                            |                                  |
| D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)                                                                 |                                  |
|                                                                                                                                                                            |                                  |
| E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)                                                                                                      |                                  |
| The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession<br>Number of Deposit")    |                                  |
|                                                                                                                                                                            |                                  |

|                                                                                                                                                                          |                                                                                                                                                                           |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p align="center">For receiving Office use only</p> <p><input type="checkbox"/> This sheet was received with the international application</p> <p>Authorized officer</p> | <p align="center">For International Bureau use only</p> <p><input type="checkbox"/> This sheet was received by the International Bureau on:</p> <p>Authorized officer</p> |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------|